

# Comments about quantum symmetries of $SU(3)$ graphs

R. Coquereaux<sup>1</sup> \*, D. Hammaoui<sup>2</sup> †, G. Schieber<sup>1,2,3‡</sup>, E. H. Tahri<sup>2</sup> §

<sup>1</sup> *CPT - Centre de Physique Théorique - CNRS  
Campus de Luminy - Case 907  
F-13288 Marseille - France*

<sup>2</sup> *LPTP - Laboratoire de Physique Théorique et des Particules  
Département de Physique, Faculté des Sciences  
Université Mohamed I, B.P.524 Oujda 60000 - Maroc*

<sup>3</sup> *CBPF - Centro Brasileiro de Pesquisas Físicas  
Rua Dr. Xavier Sigaud, 150  
22290-180, Rio de Janeiro, Brasil*

## Abstract

For the  $SU(3)$  system of graphs generalizing the ADE Dynkin digrams in the classification of modular invariant partition functions in CFT, we present a general collection of algebraic objects and relations that describe fusion properties and quantum symmetries associated with the corresponding Ocneanu quantum groupoids. We also summarize the properties of the individual members of this system.

**Keywords:** conformal field theory, modular invariance, Coxeter-Dynkin graphs, fusion algebra, quantum symmetries, quantum groupoids

---

\*E-mail:Robert.Coquereaux@cpt.univ-mrs.fr

†E-mail:d.hammaoui@sciences.univ-oujda.ac.ma

‡Supported by a fellowship of AUF - Agence Universitaire de la Francophonie. E-mail: schieber@cbpf.br

§E-mail:tahrie@sciences.univ-oujda.ac.ma

# 1 Introduction

**The stage** Along the last fifteen years or so, investigations performed in a number of research fields belonging to theoretical physics or to mathematics suggest the existence of “fundamental objects” generalizing the usual simply laced ADE Dynkin diagrams. Let us mention a few of these fields: statistical mechanics, string theory, quantum gravity, conformal field theory, theory of bimodules, Von Neumann algebras, sector theory, (weak) Hopf algebras, modular categories, etc.

Properties of the algebraic structures associated with the choice of such a fundamental object have been analysed independently by several groups of people, with their own tools and terminology. The results obtained by these different schools are not always easy to compare, or even to apprehend, because of the required background and specificity of the language.

However, at the heart of any such fundamental object we meet a graph (or the adjacency matrix that encodes this graph). We believe that many important and useful results can be described in an elementary way obtained from the combinatorial data provided by the graph itself, or by some kind of attached modular data [23].

Roughly speaking, if we have a modular invariant (but not any kind of modular invariant), we have a (particular type of) quantum groupoid, and conversely. Now every such quantum groupoid is encoded by a graph, and this graph leads naturally to two (in general distinct) character theories: one is the so called fusion algebra, and the other is the algebra of quantum symmetries. This is the story that we want to tell. But we want to tell it in simple words, using elementary mathematics. And we want to tell it in the case of the  $SU(3)$  system of graphs, i.e., the so-called “Di Francesco - Zuber diagrams” that generalize the familiar ADE Dynkin diagrams.

As already mentioned, several groups of people (without trying to be exhaustive, we can cite [43, 23, 20, 45, 3, 12, 15, 8, 42, 6]) have investigated related topics along the past years. We believe that only A. Ocneanu has actually worked out all these examples in details, with his own language, from the point of view of the study of quantum symmetries, but his results are unfortunately unpublished and not available.

**Purpose** The purpose of this article is three-fold.

- 1) To present, in a synthetic and elementary way, a collection of algebraic objects describing fusion properties and quantum symmetries associated with graphs belonging to (higher) Coxeter-Dynkin systems.
- 2) To present a summary of results concerning members of the  $SU(3)$  system.
- 3) To make a number of comments about the various aspects of this subject, and, in some cases, to establish a distinction between what is known and what is believed to be true.

**Warning** This paper is not a review. If it is true that many results recalled here can be found in the litterature, maybe with another language or perspective, many others cannot be found elsewhere. It may well be that a number of these results have been privately worked out by several people, but, if so, they are not available. What we present here, including a good part of the terminology itself, is mostly the result of our own understanding, that has been growing up along the years.

However, this paper is not a detailed research paper either. Indeed, it is, in a sense, too short. Every single example summarized in section 6, for instance, gives rise to interesting, and, sometimes difficult, problems, and would certainly be worth a dedicated article. What we put in this section is only what we think should be remembered once all the details will have been forgotten. This, admittedly, is a partial viewpoint.

We want this paper to be used as a compendium of results, terminology, and remarks.

**Plan** The plan of this article is as follows. In the next section we summarize the properties of the  $\mathcal{A}$  system, i.e., the Weyl alcoves at level  $k$ , from the viewpoint of fusion and graph algebras. In section 3, we describe general properties associated with any member of the  $SU(3)$  system of graphs. This applies, in particular, to the  $\mathcal{A}$  graphs themselves, but they are very particular, and this is why we singled them out. In the fourth section, we describe, in plain terms, the Ocneanu quantum groupoid associated with a graph  $G$ , or, better, with a pair  $(G, \mathcal{A}_k)$ . We do not give however any information about the methods that allow one to compute the values of the corresponding cells; this is a most essential question but it should be dealt with in another publication. In the fifth section we describe the equations that allow one to recover the algebra of quantum symmetries (and sometimes the graph itself) from the data provided by a modular invariant, the leitmotiv of this section being the so-called “modular splitting technique”. Although we have used repeatedly this technique to solve several quite involved examples briefly described in section 6, we do not explicitly discuss here our method of resolution but refer to forthcoming articles (or theses) for these – important – details [27, 26, 24]. In section 6 we summarize what is known, or at least what we know, about the structure of the algebra of quantum symmetries for each member of the  $SU(3)$  series. At this point we should stress that the graphs themselves, together with their fusion properties (relations with the  $\mathcal{A}$  system) or with the associated modular invariants, have been discovered and described long ago (by Di Francesco and Zuber [17]). Several aspects related to the theory of sectors, or to the theory of bimodules have also been investigated independently by different groups of people [2, 3, 20, 18]. However we believe that only A. Ocneanu performed a detailed analysis of the algebra of quantum symmetries associated with all these diagrams and three of us remember vividly the poster describing the Cayley graph for the generators of the algebra that we call  $Oc(\mathcal{E}_9)$ , on one of the walls of the Bariloche conference lecture hall, during the January 2000 summer (!) school. However, this material was never published or even made public on the internet. Our techniques may

be sometimes clumsy but we hope that they are understandable and will draw attention of potential readers on this fascinating subject. We now return to the plan of our paper and mention that the last section (the 7th) is devoted to a set of final remarks describing possible new directions or open problems.

## 2 $\mathcal{A}_k$ graphs

### 2.1 First properties

The  $\mathcal{A}_k$  graphs are obtained as truncations of the Weyl chambers of  $SU(N)$  at some level (Weyl alcoves). They have a level  $k$  and a (generalized) Coxeter number  $\kappa = k + N$ . From now on  $N = 3$ .

**Vertices** Vertices  $\lambda$  may be labelled by Dynkin labels  $(\lambda_1, \lambda_2)$ , with  $0 \leq \lambda_1 + \lambda_2 \leq k$ , by shifted Dynkin labels  $\{\lambda_1 + 1, \lambda_2 + 1\} = (\lambda_1, \lambda_2)$ , or by Young tableaux<sup>1</sup>  $Y[p, q]$ ,  $p = \lambda_1 + \lambda_2$ ,  $q = \lambda_2$ . For instance, the unit vertex (trivial representation) is  $(0, 0) = \{1, 1\} = Y[0, 0]$ , the fundamental vertex  $(1, 0) = \{2, 1\} = Y[1, 0]$  and its conjugate  $(0, 1) = \{1, 2\} = Y[1, 1]$ . The graph  $\mathcal{A}_k$  possesses  $d_{\mathcal{A}_k} = (k + 1)(k + 2)/2$  vertices. The vector space spanned by these vertices is also called  $\mathcal{A}_k$ .

**Conjugation** The graph  $\mathcal{A}_k$  has an involution  $\star: (\lambda_1, \lambda_2) \rightarrow (\lambda_2, \lambda_1)$  called conjugation.

**Triality** Each vertex  $\lambda$  possesses a triality  $t(\lambda) = \lambda_1 - \lambda_2 \pmod{3}$ . It is equal to the number of boxes modulo 3 of the corresponding Young tableau. Conjugation leaves triality 0 invariant and interchanges 1 and 2.

**Edges** Edges are oriented. They only connect vertices of increasing triality, by step +1, i.e. we choose one of the two possible adjacency matrices (the other is its transpose, with the edges in the opposite direction).

### 2.2 Spectral properties

**Exponents and norm** The adjacency matrix of the graph  $\mathcal{A}_k$  possesses  $d_{\mathcal{A}_k}$  distinct complex eigenvalues[50]:

$$\beta(r_1, r_2) = e^{-\frac{2i\pi(2(r_1+1)+(r_2+1))}{3\kappa}} \left( 1 + e^{\frac{2i\pi(r_1+1)}{\kappa}} + e^{\frac{2i\pi((r_1+1)+(r_2+1))}{\kappa}} \right), \quad (1)$$

where  $r_1, r_2 \geq 0$  and  $r_1 + r_2 \leq k$ . Such pairs of integers  $(r_1, r_2)$  are called exponents of the graph  $\mathcal{A}_k$ . The vertices of the  $\mathcal{A}_k$  graph can be indexed by the same set of integer pairs  $(r_1, r_2)$ : they coincide with the Dynkin labels  $(\lambda_1, \lambda_2)$ . The set of eigenvalues is invariant

---

<sup>1</sup> $p$  (resp.  $q$ ) is the number of boxes in the first (resp. second) line.

under the group  $\mathbb{Z}_3$ . One of these eigenvalues  $\beta \doteq \beta(0,0)$  is real, positive, and of largest absolute value. It is called the norm of the graph, and it is equal to  $\beta = 1 + 2 \cos(2\pi/\kappa)$ .

**Class vectors, dimension vector and quantum dimensions** Normalized eigenvectors of the adjacency matrix are denoted  $c_{r_1, r_2}$ . They can be called “class vectors” in analogy with the situation that prevails for finite groups. Here “normalized” means that the first component<sup>2</sup> of each class vector, corresponding to the unit vertex, is set to 1. The normalized eigenvector associated with the biggest eigenvalue  $\beta$  is called the dimension vector, or the Perron-Frobenius vector. Its components define the quantum dimensions of the corresponding vertices of  $\mathcal{A}_k$ . The quantum dimension of a given vertex  $\lambda = (\lambda_1, \lambda_2)$  is given by the q-analog of the classical formula for dimensions of  $SU(3)$  irreps, usual numbers being replaced by quantum numbers:  $qdim(\lambda) = (1/[2]_q)([\lambda_1 + 1]_q[\lambda_2 + 1]_q[\lambda_1 + \lambda_2 + 2]_q)$ , where  $q = \exp(i\pi/\kappa)$  is a root of unity and  $[n]_q = \frac{q^n - q^{-n}}{q - q^{-1}}$ . The norm  $\beta$  itself is the quantum dimension of the fundamental vertices (1,0) and (0,1). The sum of  $qdim(\lambda)^2$  is called the order or the quantum mass of  $\mathcal{A}_k$  and denoted  $m(\mathcal{A}_k)$ .

### 2.3 Fusion algebra

The vector space  $\mathcal{A}_k$  possesses an associative (and commutative) algebra structure: it is an algebra with unity, vertex (0,0), and two generators, vertices (1,0) and (0,1), called “fundamental generators”. The graph of multiplication by the first generator (1,0) is encoded by the (oriented) graph  $\mathcal{A}_k$ : the product of a given vertex  $\lambda$  by the fundamental (1,0) is given by the sum of vertices  $\mu$  such that there is an edge going from  $\lambda$  to  $\mu$  on the graph. Equivalently, this multiplication is encoded by the adjacency matrix  $N_{(1,0)}$  of the graph. Multiplication by the other fundamental generator is obtained by reversing the arrows.

**Fusion matrices** Multiplication by generators  $\lambda = (\lambda_1, \lambda_2)$  is described by matrices  $N_\lambda$ , called fusion matrices. The identity is  $N_{(0,0)} = \mathbf{1}_{d_{\mathcal{A}}}$ . The other fusion matrices are obtained, once  $N_{(1,0)}$  is known, from the known recurrence relation for coupling of irreducible  $SU(3)$  representations (that we – of course – truncate at level  $k$ ):

$$\begin{aligned} N_{(\lambda, \mu)} &= N_{(1,0)} N_{(\lambda-1, \mu)} - N_{(\lambda-1, \mu-1)} - N_{(\lambda-2, \mu+1)} && \text{if } \mu \neq 0 \\ N_{(\lambda, 0)} &= N_{(1,0)} N_{(\lambda-1, 0)} - N_{(\lambda-2, 1)} && (2) \\ N_{(0, \lambda)} &= (N_{(\lambda, 0)})^{tr} \end{aligned}$$

where matrices  $N_{(\lambda, \mu)} = 0$  if  $\lambda = -1$  or  $k + 1$  or if  $\mu = -1$  or  $k + 1$ , and are periodic in the  $(\lambda, \mu)$  plane – the periodicity cell is a Weyl alcove and there are six of them around the origin  $\{1,1\} = (0,0)$ . These matrices have non negative integer entries  $(N_\lambda)_{\mu\nu} = N_{\lambda\mu}^\nu$  called fusion

---

<sup>2</sup>We assume that an order has been chosen on the set of vertices and that the unit vertex comes first.

coefficients. They form a faithful representation of the fusion algebra:

$$N_\lambda N_\mu = \sum_\nu N_{\lambda\mu}^\nu N_\nu . \quad (3)$$

Conjugation (operation  $\star$ ) on these matrices is obtained by transposition.

**Essential paths (also called horizontal paths)** Since fusion matrices  $N_\lambda$  have non negative integer entries, one can associate a graph to every fusion matrix. If the matrix element  $(N_\lambda)_{\mu\nu} = p$ , we introduce  $p$  oriented edges from the vertex  $\mu$  to the vertex  $\nu$ . Such an edge is called an essential path of type  $\lambda$  from  $\mu$  to  $\nu$ . Remember that these indices are themselves Young tableaux. The graph associated with the fundamental generator  $(1,0)$  is the  $\mathcal{A}_k$  graph itself.

## 2.4 Modular considerations

The graphs  $\mathcal{A}_k$  support a representation of the group  $SL(2, Z)$ . This group is generated by two transformations  $S$  and  $T$  satisfying  $S^2 = (ST)^3 = C$ , with  $C^2 = 1$ . The modular group itself, called  $PSL(2, Z)$  is the quotient of this group by the relation  $C = 1$ .

**The modular generator  $S$**  The adjacency matrix of  $\mathcal{A}_k$  can be diagonalized by a matrix constructed from the set of eigenvectors (all eigenvalues are distinct). As fusion matrices  $N_\lambda$  commute, this matrix therefore diagonalizes all fusion matrices. Each line of this matrix is given by a (renormalized) class vector. We renormalize the lines in order that each line is of norm 1. We therefore divide each class vector by its norm. The obtained diagonalizing matrix is then unitary but not a priori symmetric, and not necessarily related to the generator of the modular group. To write such an unitarizing matrix, one has first to choose an order on the set of eigenvalues (this fixes the ordering of line vectors), and also an order on the set of vertices of the graph (this fixes the ordering of the components for each line). One member of this family of unitarizing matrices gives the modular generator  $S$ . The point is that vertices of the graph  $\mathcal{A}_k$  have to be indexed by the same set of integers as the eigenvalues themselves<sup>3</sup>. So, whatever the order we choose on the set of vertices, we decide to choose the *same* order on the set of eigenvalues. This procedure determines – for each ordering of the vertices – a particular unitarizing matrix which can be identified with the modular generator  $S$ . It coincides with the expression explicitly given by the formula [28, 22]:

$$S_{\lambda\mu} = \frac{-i}{\sqrt{3\kappa}} (e_\kappa[2\lambda_1\mu_1 + \lambda_1\mu_2 + \lambda_2\mu_1 + 2\lambda_2\mu_2] - e_\kappa[-\lambda_1\mu_1 + \lambda_1\mu_2 + \lambda_2\mu_1 + 2\lambda_2\mu_2] \\ - e_\kappa[2\lambda_1\mu_1 + \lambda_1\mu_2 + \lambda_2\mu_1 - \lambda_2\mu_2] + e_\kappa[-\lambda_1\mu_1 + \lambda_1\mu_2 - 2\lambda_2\mu_1 - \lambda_2\mu_2] \\ + e_\kappa[-\lambda_1\mu_1 - 2\lambda_1\mu_2 + \lambda_2\mu_1 - \lambda_2\mu_2] - e_\kappa[-\lambda_1\mu_1 - 2\lambda_1\mu_2 - 2\lambda_2\mu_1 - \lambda_2\mu_2]) ,$$

---

<sup>3</sup>We thank O. Ogievetsky for this remark.

where  $e_\kappa[x] := \exp[\frac{-2i\pi x}{3\kappa}]$  and where the vertices are labelled by shifted Dynkin labels  $\lambda = \{\lambda_1, \lambda_2\}$ ,  $\mu = \{\mu_1, \mu_2\}$ . This  $d_{\mathcal{A}_k}^2$  matrix  $S$ , obtained as a –properly normalized and ordered – quantum “character table” , defines the quantum analogue of a Fourier transform for the graphs  $\mathcal{A}_k$ . The matrix  $S$  is symmetric and such that  $S^4 = 1$ . In the opposite direction, the well known Verlinde formula [49] expresses fusion matrices  $N_\lambda$  in terms of coefficients of  $S$ :

$$\mathcal{N}_{\lambda\mu}^\nu = \sum_{\beta \in \mathcal{A}_k} \frac{S_{\lambda\beta} S_{\mu\beta} S_{\nu\beta}^*}{S_{0\beta}}, \quad (4)$$

where  $\lambda = 0 = (0, 0)$  is the trivial representation. In the present paper we prefer to obtain the  $S$  matrix from the combinatorial data provided by the graph.

**The modular generator  $T$**  The modular generator  $T$  is diagonal in the basis defined by vertices. Its eigenvalue associated with a vertex of shifted coordinates  $\lambda = \{\lambda_1, \lambda_2\}$  is equal to [28]:

$$T_{\lambda\lambda} = \exp \left[ 2i\pi \left( \frac{[\lambda_1^2 + \lambda_1\lambda_2 + \lambda_2^2] - \kappa}{3\kappa} \right) \right]. \quad (5)$$

The square bracket in the numerator of the argument of  $\exp$  can be simply read from the coordinates of the chosen vertex since it is the corresponding eigenvalue for the quadratic Casimir of the Lie group  $SU(3)$ . We call “modular exponent” the whole numerator (i.e., the difference between the Casimir and the generalized Coxeter value  $\kappa$ ) taken modulo  $3\kappa$ . The  $T$  operator is therefore essentially (up to a trivial geometric phase) obtained as the exponential of the quadratic Casimir: the values for the shift ( $-\kappa$ ) and multiplicative constant ( $3\kappa$ ) can indeed be fixed by imposing that the  $SL(2, \mathbb{Z})$  relation  $(ST)^3 = S^2$  hold.

**The  $SL(2, \mathbb{Z})$  representation defined by  $\mathcal{A}_k$**  Matrices  $S$  and  $T$  provide therefore a representation of the group  $SL(2, \mathbb{Z})$  for each alcove of  $SU(3)$ . Actually, one obtains moreover the identity  $T^{3\kappa} = 1$  so that this representation factorizes through the finite group  $SL(2, \mathbb{Z}/3\kappa\mathbb{Z})$ .

## 2.5 Symmetry and automorphism

**The  $Z_3$  action** Rotations of angle  $0, 2\pi/3$  or  $4\pi/3$  around the center of the equilateral triangle associated with the graph  $\mathcal{A}_k$  define a  $Z_3$  action – that we denote by  $z$  – on the set of vertices and therefore an endomorphism of the algebra (its cube is the identity). Its action on the irreps labelled by Dynkin labels  $(\lambda_1, \lambda_2)$  is given by:

$$z(\lambda_1, \lambda_2) = (k - \lambda_1 - \lambda_2, \lambda_1). \quad (6)$$

**The Gannon automorphism  $\rho$**  It is defined on the vertices, as [22]

$$\rho = z^{kt}, \quad (7)$$

where  $t$  is the triality and  $k$  is the level of the graph. We found the following result [25]: if vertices  $v_1$  and  $v_2$  are such that  $v_2 = \rho[v_1]$ , then  $T[v_1] = T[v_2]$ . The proof is given in [25].

### 3 General properties of the $SU(3)$ system of graphs

This is a collection of graphs. As it will be discussed later, each graph  $G$  gives rise to a weak Hopf algebra (a quantum groupoid)  $\mathcal{B}G$ , and each graph  $G$  is also associated with a given  $\widehat{su}(3)$  modular invariant  $Z$ . At the moment, we suppose that the collection of graphs (also called the “Coxeter-Dynkin system of type  $SU(3)$ ”) is given and we list several of their properties. Several graphs (the orbifolds of the A series) were obtained by Kostov [31] but the full list of graphs for this system was obtained by Di Francesco and Zuber [17, 16]. Later, A. Ocneanu, at the Bariloche school 2000 [40], explained why one member of their original list had to be removed.

#### 3.1 First properties

**Vertices and edges** Vertices of  $G$  are denoted  $a, b, c, \dots$ . Edges are oriented. In some cases there are multiple edges between two vertices.

**Spectral properties of the graph  $G$**  A graph  $G$  belonging to the  $SU(3)$  system is characterized by an adjacency matrix. Its biggest eigenvalue is called  $\beta = 1 + 2 \cos(2\pi/\kappa)$ . The Coxeter number  $\kappa$  is read from  $\beta$ . The level is defined as  $k = \kappa - 3$ . The set of eigenvalues of the graph  $G$  is a subset of the eigenvalues of the graph  $\mathcal{A}_k$  with same level. They are of the form  $\beta(r_1, r_2)$  in Eq.(1), with possible multiplicities. The pairs of integers  $(r_1, r_2)$  are called the exponents of the graph  $G$ .

**The associated modular invariant**  $SU(3)$  graphs have been proposed as graphs associated to  $\widehat{su}(3)$  modular invariant partition functions. These partition functions  $Z$  are sesquilinear forms on the characters labelled by irreps of  $\widehat{su}(3)_k$ . The correspondance is such that diagonal terms of  $Z$  match the set of exponents for the corresponding graph  $G$ . The interpretation for the off diagonal terms of  $Z$  was found by A. Ocneanu [39, 38]. We shall come back to this later.

**Quantum dimensions and order of  $G$**  One of the vertices of the graph  $G$ , denoted  $\mathbf{0}$ , is called the unit vertex. It is defined from the eigenvector corresponding to  $\beta$  as the vertex associated to the smallest component<sup>4</sup>. The components of the normalized eigenvector associated with  $\beta$  (the dimension vector) define the quantum dimensions of the corresponding vertices – normalisation is obtained by setting to 1 the quantum dimension of the unit vertex<sup>5</sup>. When there is only one arrow leaving (and going to) the unit vertex<sup>6</sup>  $\mathbf{0}$ , the quantum dimensions of

---

<sup>4</sup>If the graph possesses some (classical) symmetry, there can be several vertices associated to the smallest component. In those cases, we just choose one of them.

<sup>5</sup>It plays indeed the role of a unit when the graph  $G$  has self-fusion (see later), otherwise it is only a vertex whose quantum dimension is 1.

<sup>6</sup>This is for instance not so for  $\mathcal{D}_k^*$ .



its two neighbours (denoted as  $\mathbf{1}$  and  $\mathbf{1}^*$ ) are both equal to  $\beta$ . The sum of the squares of the quantum dimensions of vertices is called the order or the quantum mass of  $G$ , and denoted  $m(G)$ .

### 3.2 The two representation theories associated with the bialgebra $\mathcal{B}G$

A quantum groupoid  $\mathcal{B}G$  is associated with any graph  $G$  of the  $SU(3)$  system. It is both semi-simple and co-semi-simple. We present several basic properties here; more details will be given in Section 4.

**The fusion algebra  $A(G)$**  The algebra  $\mathcal{B}G$  endowed with its associative product is a direct sum of matrix algebras labelled by the index  $\lambda$  (i.e., by vertices of the  $\mathcal{A}_k$  graph with same level). Its representation theory (algebra of characters)  $A(G)$  is isomorphic to the fusion algebra of  $\mathcal{A}_k$ . Matrix representatives of the generators  $\lambda$  of  $\mathcal{A}_k$  have been already introduced: they correspond to the fusion matrices  $N_\lambda$ .

**The algebra of quantum symmetries  $Oc(G)$**  The dual algebra  $\widehat{\mathcal{B}}G$  endowed with its associative product is also a direct sum of matrix algebras labelled by an index  $x$ . Its representation theory (algebra of characters) is called the “algebra of quantum symmetries” of  $G$  and denoted  $Oc(G)$ . We call  $d_O$  the dimension of  $Oc(G)$ . It is an algebra with a unit (denoted 0) and, for  $SU(3)$  graphs, with – in general but not always – two algebraic generators (called chiral left and chiral right generators and denoted as  $1_L$  and  $1_R$ ), together with their conjugates  $1_L^*$  and  $1_R^*$ . The Cayley graph of multiplication by the two generators  $1_L$  and  $1_R$  (two types of lines) is called the Ocneanu graph of  $G$ . The graph corresponding to the conjugated generators  $1_L^*$  and  $1_R^*$  is obtained from the (oriented) Ocneanu graph by reversing the arrows.  $Oc(G)$  has also another conjugation, called the chiral conjugation, that permutes the two algebraic generators  $1_L$  and  $1_R$ . Another way of displaying the Cayley graph is to draw only the graph of multiplication by one chiral generator, say  $1_L$ , and to associate (for example using dashed lines) each basis element with its chiral conjugate. Multiplication of a vertex  $x$  by the chiral generator  $1_R$  is obtained as follows: we start with  $x$ , follow the dashed lines to find its chiral vertex  $y$ , then use the multiplication by  $1_L$  and finally pull back using the dashed lines to obtain the result. Linear generators of  $Oc(G)$  (i.e., vertices of the Ocneanu graph) that are identical with their chiral conjugates are called self-dual. The two subalgebras generated by the chiral generators are called chiral subalgebras. The intersection of these two subalgebras is called the ambichiral subalgebra, and its generators are the ambichiral generators (they are self-dual).  $Oc(G)$ , like  $A(G) \simeq \mathcal{A}_k$ , is not only an algebra but an algebra that comes with a particular basis (the vertices of the Ocneanu graph), for which structure constants are non negative integers. The multiplication between vertices reads  $xy = \sum_z O_{xy}^z z$ , where  $O_{xy}^z$ , called quantum symmetry coefficients, are non negative integers. Matrix representatives of these linear generators  $x$  of  $Oc(G)$  are called “Ocneanu matrices” and denoted  $O_x$ , with

elements  $(O_x)_{yz} = O_{xy}^z$ . They form an anti-representation of the Ocneanu algebra:

$$O_x O_y = \sum_z O_{yx}^z O_z . \quad (8)$$

If  $Oc(G)$  is commutative - which is not always so - then  $O_{xy}^z = O_{yx}^z$  and the Ocneanu matrices form a representation of the Ocneanu algebra:  $O_x O_y = \sum_z O_{xy}^z O_z$ . The structure of  $Oc(G)$  is very much case dependent. One of the purpose of this paper is actually to present the corresponding results (for the  $SU(3)$  system) in a synthetic way. In many cases  $Oc(G)$  can be written as the direct sum of a chiral subalgebra and one or several modules over this subalgebra. Knowledge of the Ocneanu graph (i.e., the action of  $1_L$  and  $1_R$ ) may sometimes be insufficient to encode the full structure (like for the  $D_4$  case of the  $SU(2)$  system). Matrices  $O_{1_L}$  and  $O_{1_R}$  are the adjacency matrices of the Ocneanu graph. The two dimension vectors (normalized eigenvectors associated with the largest eigenvalue for each adjacency matrix) allow one to attribute – unambiguously – quantum dimensions to all the linear generators of  $Oc(G)$ . Actually, the two chiral generators have dimension  $\beta$  and the whole list of quantum dimensions can be read directly from the Ocneanu graph by using the fact that this property is multiplicative  $qdim(xy) = qdim(x)qdim(y)$ . The sum of their squares is called the order or the quantum mass of  $Oc(G)$ , denoted  $m(Oc(G))$ : it is equal to the order of  $m(\mathcal{A}_k)$  of  $\mathcal{A}_k = A(G)$ . This property generalizes the usual group theory result.

### 3.3 $G$ as a module over $A(G) = \mathcal{A}_k$

Call also  $G$  the vector space spanned by the vertices of a graph  $G$ . Call  $r$  the number of vertices of the graph. This vector space is a module for the action of the fusion algebra associated with  $\mathcal{A}_k$ , where  $k$  is the level of  $G$  (Coxeter number minus 3). The action is defined by the relation  $\lambda a = \sum_b F_{\lambda a}^b b$ , where  $F_{\lambda a}^b$  are non negative integers called fused or annular coefficients. In some cases, the same graph  $G$  may also be a module over some other graph of type  $A$  with a different Coxeter value, but we are not interested in this phenomenon.

**Annular matrices** This action is encoded by a set of matrices  $F_\lambda$  called annular matrices or fused (not fusion !) matrices, defined by  $(F_\lambda)_{ab} = F_{\lambda a}^b$ . From the module property  $\lambda(\mu a) = (\lambda\mu) a$ , the annular matrices satisfy:

$$F_\lambda F_\mu = \sum_\nu N_{\lambda\mu}^\nu F_\nu . \quad (9)$$

They form a representation of the fusion algebra (usually of different dimension since  $r \neq d_{\mathcal{A}_k}$ ). They are obtained by the same recurrence relation (2) as the fusion matrices but with  $F_{(0,0)} = \mathbf{1}_{r \times r}$  and  $F_{(1,0)} = Ad(G)$ , where  $Ad(G)$  is the adjacency matrix of  $G$ . We obtain in this way  $d_A$  matrices of size  $r \times r$ . As before  $d_A$  is the number of vertices of the associated  $\mathcal{A}_k$  graph, the index  $\lambda$  of  $F_\lambda$  is a Young tableau.

**Essential paths (also called horizontal paths)** Since annular matrices  $F_\lambda$  have non negative integer entries, one can associate a graph to every such matrix. If the matrix element of  $(F_\lambda)_{ab} = p$ , we introduce  $p$  oriented edges between vertices  $a$  and  $b$  of  $G$ . Such an edge is called an essential path of type  $\lambda$  from  $a$  to  $b$ . This graph will be called the horizontal graph of type  $\lambda$ . Remember that the  $\lambda$  index is a Young tableau (a vertex of the corresponding  $\mathcal{A}_k$  diagram). The graph associated with the generator  $F_{(1,0)}$  is the graph  $G$  itself.

**Essential matrices (or horizontal matrices)** Essential matrices have the same information contents as the annular matrices, however, they are rectangular rather than square. They are defined as follows

$$(E_a)_{\lambda b} \doteq (F_\lambda)_{ab} . \quad (10)$$

We have therefore one essential matrix  $E_a$  for each vertex  $a$  of the graph  $G$ . The integer  $(E_a)_{\lambda b}$  gives the number of horizontal paths of type  $\lambda$  from  $a$  to  $b$ . The property (9) can be written as follows using essential matrices:

$$N_\lambda E_a = E_a F_\lambda . \quad (11)$$

In particular we have  $N_{(1,0)} E_0 = E_0 F_{(1,0)}$ . The essential matrix  $E_0$  associated with the unit 0 of the graph  $G$  intertwines the adjacency matrices of the graphs  $G$  and  $\mathcal{A}$ : it is also called the  $(\mathcal{A}_k, G)$  intertwiner.

**Restriction-induction coefficients** Non-zero entries of the first line of  $F_\lambda$  (ie relative to the unit vertex of  $G$ ) are called restriction coefficients. They define a restriction from  $\mathcal{A}_k$  to  $G$  (like irreps of a group versus irreps of a subgroup). The branching rules are given by:

$$\lambda \hookrightarrow \sum_b (F_\lambda)_{1b} b = \sum_b (E_0)_{\lambda b} b . \quad (12)$$

The line indices corresponding to the non-zero entries of the column  $b$  of the matrix  $E_0$  are called induction coefficients associated with the vertex  $b$ . They give the vertices  $\lambda$  for which  $b$  appears in their branching rules. The line indices (Young tableaux) corresponding to the non-zero entries of the first column of the matrix  $E_0$  are called degrees of the family of would-be quantum invariants tensors by analogy with the situation that prevails for finite subgroups of Lie groups (for instance, when  $G$  is the fusion graph by the fundamental representation of binary polyhedral groups, these non-zero entries of the first column of  $E_0$  reflect the existence of invariant symmetric tensors and therefore give the degrees of the Klein invariant polynomials for symmetry groups of Platonic bodies).

### 3.4 $G$ as a module over $Oc(G)$

The vector space  $G$  is also a module for the action of the algebra of quantum symmetries  $Oc(G)$ . Call  $x$  the elements of  $Oc(G)$ . The action is defined by the relation  $x a = \sum_b S_{xa}^b b$ , where  $S_{xa}^b$  are non negative integers called dual annular coefficients.

**Dual annular matrices** The action can be encoded in a set of matrices  $S_x$  that we call the dual annular matrices, defined by  $(S_x)_{ab} = S_{xa}^b$ . From the module property  $x(ya) = (xy)a$ , the dual annular matrices satisfy:

$$S_x S_y = \sum_z O_{yx}^z S_z . \quad (13)$$

They satisfy the same relations as the Ocneanu matrices  $O_x$  (they form an anti-representation of the quantum symmetry algebra). We obtain in this way  $d_O$  matrices of size  $r \times r$ . As before  $d_O$  is the number of vertices of the associated Ocneanu graph.

**Vertical paths** Since dual annular matrices  $S_x$  have non negative integer entries, one can associate a graph to every such matrix. If the matrix element  $(S_x)_{ab} = p$ , we introduce  $p$  oriented edges between vertices  $a$  and  $b$  of  $G$ . Such an edge is called a vertical path of type  $x$  from  $a$  to  $b$ . This graph will be called the vertical graph of type  $x$ . The vertical graphs associated with the two chiral generators of  $Oc(G)$  coincide with  $G$  itself.

**Vertical matrices** Vertical matrices have the same information content as the dual annular matrices, however, they are rectangular rather than square. They are defined as follows:

$$(R_a)_{xb} \doteq (S_x)_{ab} . \quad (14)$$

We have therefore one vertical matrix  $R_a$  for each vertex  $a$  of the graph  $G$ . The integer  $(R_a)_{xb}$  gives the number of vertical paths of type  $x$  from  $a$  to  $b$ .

### 3.5 Self-fusion

$\mathcal{A}_k$  diagrams have self-fusion (the fusion algebra). A graph  $G$  has self-fusion when the vector space spanned by its vertices is not only a module over the corresponding  $A(G)$  fusion algebra but when it possesses an associative algebra structure encoded by the graph itself (its adjacency matrix), with non negative integral structure constants, compatible with the already known  $A(G)$  action. If  $a, b, c, \dots$  are vertices of a graph  $G$  with self-fusion, we have  $ab = \sum_c G_{ab}^c c$ , where the coefficients are non negative integers. The unit  $\mathbf{0}$  of the graph is the identity for the multiplication. The multiplication of some chosen vertex by the special vertex  $\mathbf{1}$  (resp.  $\mathbf{1}^*$ ) is given by the sum of vertices  $a$  such that there is an edge of  $G$  from the chosen vertex to  $a$  (resp. from  $a$  to the chosen vertex). The compatibility condition between self-fusion and module structure reads  $\lambda(ab) = (\lambda a)b$ .

**Conjugation** Conjugation is defined for all self-fusion graphs. It is compatible with the conjugation already defined for  $A$  graphs. We call  $a^*$  the conjugate of  $a$  in  $G$ . The compatibility condition is understood as follows: all vertices of  $\mathcal{A}_k$  appearing in the induction list associated with  $a^*$  should be the conjugated vertices (taken in  $\mathcal{A}_k$ ) of those associated with

$a$ . When these two sets are equal, then  $a^* = a$ . This provides a method for determining the conjugation of the  $G$  vertices. We have  $(\lambda a)^* = \lambda^* a^*$ , thus the annular coefficients should satisfy  $(F_{\lambda^*})_{a^* b^*} = (F_{\lambda})_{ab}$ .

**Triality** Triality is also defined for all graphs with self-fusion. It is compatible with the triality already defined for  $A$  graphs. This compatibility condition is understood as follows: if the level of the graph  $G$  is  $k$ , then all the vertices of  $\mathcal{A}_k$  appearing in the induction list associated with a given vertex of  $G$  should have the same triality. This provides a method for determining the triality of the  $G$  vertices.

**Graph matrices** The fusion of  $G$  vertices can be encoded in a set of matrices  $G_a$  with non negative integer coefficients  $(G_a)_{bc} = G_{ab}^c$ , called graph matrices. We have  $G_0 = F_{(0,0)}$ ,  $G_1 = F_{(1,0)}$  and  $G_{1^*} = F_{(0,1)}$ . The compatibility condition for graphs with self-fusion (cf supra) reads  $G_a F_{\lambda} = F_{\lambda} G_a$ . In particular, using essential matrices  $E_a$  defined in Eq.(10) one can get  $E_a = E_0 G_a$ .

**Remark** Some of the graphs belonging to a Coxeter-Dynkin system have self-fusion, others don't. For example, in the  $SU(2)$  system, the diagrams  $A_n$ ,  $D_{even}$ ,  $E_6$  and  $E_8$  have self-fusion, this is not the case for  $D_{odd}$  and  $E_7$ . In the  $SU(3)$  system, diagrams  $\mathcal{A}_k$ ,  $\mathcal{D}_{3n}$ ,  $\mathcal{E}_5$ ,  $\mathcal{E}_9$  and  $\mathcal{E}_{21}$  have self-fusion. The others don't.

**Flatness** We believe that self-fusion is equivalent to flatness, as defined for instance in [35, 36] or [29]. The two notions look a priori very different but it seems that all known graphs with self-fusion are also flat (and reciprocally). We are not aware of any formal proof relating the two concepts.

### 3.6 Coxeter-Dynkin systems of graphs, self-connections and Kuperberg spiders

A graph that is a member of a Coxeter-Dynkin system gives rise to a particular kind of quantum groupoid. Such a graph is associated with some modular invariant, but sometimes more than one graph can be associated with the same invariant. Moreover, a member of a Coxeter-Dynkin system has also to be compatible, in a sense that should be precised, with a given Lie group (here  $SU(3)$ ). Being a module over the graph algebra of a Weyl alcove at some level is a necessary but not sufficient condition. A condition, using the notion of self-connections on graphs, was given by A. Ocneanu in Bariloche (2000) [40] and this lead him to discard one of the graphs of the original Di Francesco - Zuber list. We believe that the appropriate algebraic concept can be phrased in terms of Kuperberg spiders [30] but we have no rigorous proof that the two concepts are the same.

## 4 The quantum groupoid associated to a pair $(G_1, G_2)$

If  $G_1$  has self-fusion and if  $G_2$  is a module over  $G_1$ , one can associate a bialgebra  $\mathcal{B}(G_1, G_2)$  to this pair of graphs [39]. This bialgebra is a particular type of weak Hopf algebra (or quantum groupoid) (see for instance [37, 4, 5, 34, 33]). We call it the ‘‘Ocneanu quantum groupoid’’ associated with the chosen pair. In particular if  $G_2 = G$  and  $G_1 = \mathcal{A}_k$ , with  $k$  the level of  $G$ , we just denote  $\mathcal{B}G \doteq \mathcal{B}(\mathcal{A}_k, G)$ , or simply  $\mathcal{B}$  if the choice of  $G$  is clear from the context. In what follows we consider mostly bialgebras of that type.

### 4.1 The vector spaces $\mathcal{B}$ and $\widehat{\mathcal{B}}$

**Admissible triangles** To every essential (i.e. horizontal) path of type  $\lambda$  between  $a$  and  $b$  one associates a triangle with one horizontal edge labelled by  $\lambda$  and two edges labelled by  $a$  and  $b$ . Such triangles (with 1 line of type  $A$  and 2 lines of type  $G$ ) are called admissible triangles. By duality, they can also be drawn as  $(GGA)$  vertices. The vector space spanned by such triangles is called  $EssPath(G)$  or  $Hpaths(G)$ , it is graded by  $\lambda$ :  $Hpaths(G) = \sum_{\lambda} Hpaths_{\lambda}(G)$ .

To every vertical path of type  $x$  between  $a$  and  $b$  one associates a triangle with one vertical edge labelled by  $x$  and two edges labelled by  $a$  and  $b$ . Such triangles (with 1 line of type  $Oc$  and 2 lines of type  $G$ ) are also called admissible triangles. By duality, they can also be drawn as  $(GGO)$  vertices. The vector space spanned by such triangles is called  $Vpaths(G)$ , it is graded by  $x$ :  $Vpaths(G) = \sum_x Vpaths_x(G)$ .

**Double triangles** We call  $\mathcal{B}$  the graded vector space  $\sum_{\lambda} Hpaths_{\lambda}(G) \otimes Hpaths_{\lambda}(G)$ . It is spanned by double triangles  $GGAGG$  (two triangles of type  $(GGA)$  sharing a common edge of type  $A$ ). By duality they can also be drawn as diffusion diagrams (like in Figure 1).




Figure 1: A double triangle of type  $GGAGG$  of  $\mathcal{B}$ .

We call  $\widehat{\mathcal{B}}$  the graded vector space  $\sum_x Vpaths_x(G) \otimes Vpaths_x(G)$ . It is spanned by double triangles  $GGOGG$  (two triangles of type  $(GGO)$  sharing a common edge of type  $O$ ). By duality they can also be drawn as diffusion diagrams (like in Figure 2).

### 4.2 The multiplications

**The multiplication  $\circ$  on the vector space  $\mathcal{B}$**  This algebra structure on  $\mathcal{B}$  is obtained by choosing the set of double triangles of type  $(GGAGG)$  as a basis of matrix units  $e_{IJ}$  for an asso-




Figure 2: A double triangle of type GGOGG of  $\widehat{\mathcal{B}}$ .

ciative product that we call  $\circ$ , and such that multi-indices are like  $\{I, J\} = \{(\lambda, a, b), (\lambda, c, d)\}$ , i.e. with same  $\lambda$ .

**The multiplication  $\hat{\circ}$  on the dual vector space  $\widehat{\mathcal{B}}$**  This algebra structure on  $\widehat{\mathcal{B}}$  is obtained by choosing the set of double triangles of type (GGOGG) as a basis of matrix units  $\epsilon^{AB}$  for an associative product that we call  $\hat{\circ}$ , and such that multi-indices are like  $\{A, B\} = \{(x, a, b), (x, c, d)\}$ , i.e. with same  $x$ .

**Comultiplications and compatibility : Ocneanu cells** Since we have a product  $\circ$  in  $\mathcal{B}$  we have a coproduct  $\hat{\Delta}$  in  $\widehat{\mathcal{B}}$ . Since we have a product  $\hat{\circ}$  in  $\widehat{\mathcal{B}}$  we have a coproduct  $\Delta$  in  $\mathcal{B}$ . In order to have a bialgebra structure, we need a compatibility condition for the coproducts (homomorphism property). In order to ensure this, it is not possible to assume that the two bases of double triangles that we have used in  $\mathcal{B}$  and in  $\widehat{\mathcal{B}}$  are dual bases. At the contrary, the fact that there exists a non trivial pairing (between these two bases) such that the compatibility conditions holds is the main non trivial part of the claim that  $\mathcal{B}$  is actually a bialgebra. This non trivial pairing is determined by the family of Ocneanu cells or inverse cells  $\langle \epsilon^{AB}, e_{IJ} \rangle$ , labelled with tetrahedra  $a, b, \lambda, d, c, x$  (in some cases there is more than one path – horizontal or vertical –with fixed  $\lambda$  or  $x$  and given endpoints, so that cells may depend of other indices). Explicit determination of these numerical coefficients is not studied in the present paper.

For an arbitrary graph  $G$ , there are actually several (five) sets of such coefficients generalizing the Racah-Wigner  $6j$  symbols; they obey orthogonality relations and several types (five) of mixed pentagonal relations. Their proper definition involves non-trivial normalization choices.

**Scalar product and convolution product** Making a particular choice for a scalar product in  $\mathcal{B}$ , it is possible to trade the associative product  $\hat{\circ}$ , defined on the dual vector space  $\widehat{\mathcal{B}}$  against an associative product  $*$  (convolution product) in the vector space  $\mathcal{B}$ . The situation is self-dual so that we can also find a scalar product in  $\widehat{\mathcal{B}}$  in order to trade the associative product  $\circ$  defined on  $\mathcal{B}$  against an associative product  $\hat{*}$  in the dual vector space  $\widehat{\mathcal{B}}$ .

### 4.3 Properties of $B$

It is a finite dimensional semi-simple algebra and co-semi-simple coalgebra (equivalently, its dual  $\widehat{B}$  is also a finite dimensional semi-simple algebra and co-semi-simple coalgebra).

**Quadratic sum rules** We call  $d_\lambda = \dim(HPath_\lambda)$  the dimensions of the blocks labelled by  $\lambda$ , associated with the first algebra structure, and  $d_x = \dim(VPath_x)$  the dimensions of those labelled by  $x$ , associated with the other algebra structure. Since the underlying vector space is the same, and since both algebra structures are semi-simple, we can calculate the dimension  $d_B$  of  $B$  in two possible ways and check the identity:

$$d_B = \sum_{\lambda} d_\lambda^2 = \sum_x d_x^2. \quad (15)$$

The dimensions  $d_\lambda$  and  $d_x$  can be calculated from the annular and dual annular matrices:  $d_\lambda = \sum_{a,b} (F_\lambda)_{ab}$ ,  $d_x = \sum_{a,b} (S_x)_{ab}$ .

**Linear sum rules** Call  $d_H = \sum_{\lambda} d_\lambda$  and  $d_V = \sum_x d_x$ . It happens that, in many cases, the relation  $d_H = d_V$  holds, and when it does not, one knows how to correct it. Existence of this linear sum rule (first observed in [45]) is an observational fact. Its origin is not understood.

**$B$  is not a Hopf algebra but a weak Hopf algebra (a quantum groupoid)** The main difference with the quantum group case is that the coproduct of the unit is not equal to the tensor square of the unit. What replaces it can be written  $\sum \mathbf{1}_{(1)} \otimes \mathbf{1}_{(2)}$ . The terms appearing in this sum also show up in the axioms defining weak Hopf algebras (see for instance [4]). In particular the appropriate tensor product for the category of representations is not  $\otimes$  but  $\otimes \circ \Delta \mathbf{1}$ .

**Available references** The fact that a quantum groupoid is associated with every member of a Coxeter-Dynkin system is not phrased as such in [39] but the two multiplicative structures are described there in quite general terms<sup>7</sup>. The correspondance between ADE graphs and particular weak Hopf algebras is also strongly suggested in [45]. Nowadays the fact that any member of a Coxeter-Dynkin system is associated with a quantum groupoid (as defined by [4]) belongs to the folklore (see [42, 6] for the description of this situation in the language of fusion categories and module categories). They are actually quantum groupoids of a very particular kind (so they should better be called ‘‘Ocneanu quantum groupoids’’). In the case of the  $SU(2)$  system, elementary proofs, based on axiomatic properties of Ocneanu cells, are now available in published form [14]; several explicit examples have also been worked out (for instance in [10] or [47]). In the case of the  $SU(3)$  system, general proofs are not available. Our attitude in this paper is however to take the above property for granted.

<sup>7</sup>This description is clearly related to the concept of (Ocneanu) paragroups introduced a long time before the notion of quantum groupoid.



## 5 The double fusion algebra and the modular splitting

### 5.1 Bimodule properties

**Toric matrices and double annular matrices** The Ocneanu quantum groupoids  $\mathcal{B}G$  are of a very special kind. In particular, we have the following property involving simultaneously the two representation theories associated with the bialgebra  $\mathcal{B}G$  – the fusion algebra  $A(G)$  and the quantum symmetries algebra  $Oc(G)$  :  $Oc(G)$  is an  $A(G)$  bimodule, i.e., an  $A(G) - A(G)$  module. This comes from the fact that in all cases,  $Oc(G)$  can be written as the tensor square (maybe twisted or quotiented) of some graph algebra on which  $A(G)$  acts. We write this action  $\lambda x \mu = \sum_y (V_{\lambda\mu})_{xy} y$ . The  $V_{\lambda\mu}$  are  $d_O \times d_O$  matrices with non negative integer coefficients, called double annular matrices. The same information can be encoded in  $d_{\mathcal{A}_k} \times d_{\mathcal{A}_k}$  matrices  $W_{xy}$  called toric matrices, with non negative integer coefficients defined by  $(W_{xy})_{\lambda\mu} \doteq (V_{\lambda\mu})_{xy}$ .

**Double fusion equation** The bimodule associativity property  $(\lambda\lambda')x(\mu\mu') = \lambda(\lambda'x\mu)\mu'$  leads to the following equation, called the double fusion equation:

$$V_{\lambda\mu} V_{\lambda'\mu'} = \sum_{\lambda''\mu''} N_{\lambda\lambda''}^{\lambda'} N_{\mu\mu''}^{\mu'} V_{\lambda''\mu''} . \quad (16)$$

This equation taken at  $\mu = \mu' = 0$ , at  $\lambda = \lambda' = 0$  and at  $\lambda' = \mu = 0$  leads to:

$$V_{\lambda 0} V_{\lambda' 0} = \sum_{\lambda''} N_{\lambda\lambda''}^{\lambda'} V_{\lambda'' 0} \quad (17)$$

$$V_{0\mu} V_{0\mu'} = \sum_{\mu''} N_{\mu\mu''}^{\mu'} V_{0\mu''} \quad (18)$$

$$V_{\lambda\mu'} = V_{\lambda 0} V_{0\mu'} = V_{0\mu'} V_{\lambda 0} . \quad (19)$$

Each set of matrices  $V_{\lambda 0}$  or  $V_{0\mu}$  gives therefore a representation of dimension  $d_O \times d_O$  of the fusion algebra and  $V_{00}$  is the identity matrix. They can be determined by the same recurrence relation as the fusion matrices  $N_\lambda$ , once the fundamental generators  $V_{(1,0),(0,0)}$  and  $V_{(0,0),(1,0)}$  are known.

**Other properties of  $V_{\lambda\mu}$  matrices** The action is central. Writing  $\lambda(x y)\mu = x(\lambda y\mu) = (\lambda x\mu)y$  leads to:

$$O_x V_{\lambda\mu} = V_{\lambda\mu} O_x = \sum_z (V_{\lambda\mu})_{xz} O_z . \quad (20)$$

**The Ocneanu graph** With the set of relations satisfied by  $V_{\lambda\mu}$  matrices and with the help of the known recurrence relations of irreps of  $SU(3)$ , all the coefficients  $(V_{\lambda\mu})_{xy}$  can be simply determined from the fundamental matrices  $V_{(1,0),(0,0)}$  and  $V_{(0,0),(1,0)}$ . These matrices are the adjacency matrices of the Ocneanu graph:

$$V_{(1,0),(0,0)} = O_{1_L} \quad V_{(0,0),(1,0)} = O_{1_R} . \quad (21)$$

The Ocneanu graph determines (and is determined by) these two matrices.

**Generalized partition functions** In the boundary conformal field theory associated to the given graph, the partition function on a torus with defect lines labelled by  $x$  and  $y$  is given by  $Z_{xy} = \bar{\chi} W_{xy} \chi$  where  $\chi$  is the vector of characters of affine  $su(3)$  [44].

**The modular matrix  $M$**  In particular, when there are no defect lines ( $x = y = 0$ ), we recover the modular invariant partition function  $Z = \bar{\chi} M \chi$ , since the modular invariant matrix  $M = W_{00}$  commutes with the modular generators  $S$  and  $T$  in the representation of  $SL(2, \mathbb{Z})$  associated with the Weyl alcove at this level. In contrast, the  $V_{00}$  matrix is the identity matrix.

**The double intertwining relation** From the fact that a graph  $G$  with level  $k$  is an  $\mathcal{A}_k$  module we deduced the intertwining relation given in Eq.(11), written in terms of essential matrices  $E_a$  attached to each vertex of the graph  $G$ . By analogy, let us introduce here the “essential tensor”  $K_x$ , with components  $(K_x)_{\lambda\mu y} = (V_{\lambda\mu})_{xy}$ , associated to each vertex  $x$  of  $Oc(G)$ . It can be written as a rectangular matrix of size  $d_A^2 \times d_O$  (call it double essential matrix). From the fact that  $Oc(G)$  is an  $A(G)$  bimodule, the double fusion equation (16) can be written using  $K_x$ , leading to the following double intertwining relation:

$$\tau \circ (N_\lambda \otimes N_\mu) K_x = K_x V_{\lambda\mu}, \quad (22)$$

where  $\tau$  gives a flip on tensor components:  $\tau \circ (T_{(\lambda'\lambda'')(\mu'\mu'')}) = T_{(\lambda'\mu')(\lambda''\mu'')}$ .

**Other useful formulae** We already recalled the graph interpretation for the diagonal entries of  $M$  in terms of exponents of the graph. More generally we have the following result [39, 38]. The number of vertices  $d_O$  of the Ocneanu graph (also called “number of irreducible quantum symmetries”) is equal to the sum of square of entries of the modular matrix. Moreover, the algebra of quantum symmetries is isomorphic to a direct sum of finite dimensional matrix algebras of the form  $\bigoplus_{m,n} Mat_{M_{mn}}(C)$  where  $M_{mn}$  are the entries of the modular matrix. In other words these entries give the dimensions of the irreducible representations of this algebra.

Another interpretation for these numerical entries can be given in terms of higher quantum Klein invariants (cf supra).

The above result was stated, by A. Ocneanu, for the  $SU(2)$  system. It can also be checked explicitly for all members of the  $SU(3)$  system. In the framework of the theory of sectors, such a decomposition has been proved in theorem 6.8 of [3] in a completely general setting (theorem 5.3 of the same paper shows that it is equivalent to Ocneanu’s graphical method). A nice graphical way to encode the modular matrix  $M$  associated with a graph  $G$  is provided by the “modular diagram”: it is a picture of the Weyl chamber at the given level, with

arcs connecting the vertices associated with non-zero entries  $M_{mn}$ . The degrees of quantum invariant tensors can also be read from this diagram: they correspond to those vertices that belong to the arc going through the origin  $(0,0)$ . For instance figure 3 shows these results for the  $\mathcal{D}_3$  case.




Figure 3: The modular diagram and the modular invariant associated to the  $\mathcal{D}_3$  graph

The first part of the previous theorem can be written  $d_O = Tr(M M^\dagger)$ . When the modular splitting technique (see the next section) is used to determine explicitly the  $W_{xy}$  and the algebra  $Oc(G)$  itself, the above result<sup>8</sup> provides a numerical check.

## 5.2 Modular splitting

The double fusion equation (16) at  $x = y = 0$  leads to the following equation, written in terms of  $W$  matrices, called the modular splitting equation :

$$\sum_z (W_{0z})_{\lambda\mu} (W_{z0})_{\lambda'\mu'} = \sum_{\lambda''\mu''} (N_\lambda)_{\lambda'\lambda''} (N_\mu)_{\mu'\mu''} M_{\lambda''\mu''} . \quad (23)$$

The double fusion equation (16) at  $y = 0$  leads to the following equation, written in terms of  $W$  matrices, called the generalized modular splitting equation:

$$\sum_z (W_{xz})_{\lambda\mu} (W_{z0})_{\lambda'\mu'} = \sum_{\lambda''\mu''} (N_\lambda)_{\lambda'\lambda''} (N_\mu)_{\mu'\mu''} (W_{x0})_{\lambda''\mu''} . \quad (24)$$

**Modular splitting technique I : from the modular matrix  $M$  to the toric matrices  $W_{x0}$**  In many cases, the graph  $G$  itself is not known (see comments in the last section) and the only knowledge that we have is the modular matrix  $M$ . It is possible to use the modular

<sup>8</sup>Here – and in the whole paper – we have in mind the simply laced cases (the  $ADE$  diagrams) or their generalizations.

splitting equation to determine the toric matrices. This was certainly the road followed by A. Ocneanu but a general method of resolution was first described in [11], many more details and examples can be found in [27].

One starts from the modular splitting equation (23). The fusion matrices  $N_\lambda$  and the modular matrix  $M$  are known. The right hand side of (23) is thus known: it can be seen as a matrix, called  $K$  (the “fused modular matrix”), of size  $d_A^2 \times d_A^2$ . Toric matrices that appear on the left hand side are integer entries matrices  $d_A \times d_A$  to be determined. The number of *distinct* toric matrices with one twist is equal to the rank of  $K$ . In simple cases, the number  $d_O = \text{Tr}(MM^\dagger)$  of Ocneanu generators  $O_x$  is precisely equal to the rank of  $K$ . In more complicated cases the rank of  $K$  is strictly smaller (which means that several toric matrices associated with distinct generators  $O_x$  may coincide). The explicit method leading to the determination of toric matrices (i.e., the technique used to solve the modular splitting equation) is not explained in the present paper. It is described (for a particular example) in one section of [11]. A detailed study of this method together with several  $SU(3)$  examples will be given in [27].

**Modular splitting technique II : from the toric matrices  $W_{x0}$  to the Ocneanu generators  $O_x$**  Once we have determined the toric matrices with one twist  $W_{x0}$ , we have to determine the toric matrices  $W_{xy}$ . The right hand side of the generalized modular splitting equation(24) is known. Toric matrices  $W_{xy}$  appearing on the left hand side can then be calculated. This is equivalent to solve the double intertwining relation (22) in the particular case  $x = 0$  (this is a set of linear equations that involves only the already determined toric matrices with only one twist). This leads therefore to the determination of the double annular matrices and in particular of the two chiral generators  $O_{1_L}$  and  $O_{1_R}$ . The other Ocneanu generators  $O_x$  can be determined solving Eq. (20).

**Remark** Once the algebra (or graph) of quantum symmetries  $Oc(G)$  has been obtained, we can determine the generalized Dynkin diagram  $G$  as a module graph on  $Oc(G)$ . Sometimes there is not unicity of the result and two different graphs may be associated with the same initial modular invariant. See also our comments in the last section.

**Relative modular splitting formula and relative double fusion algebra** Often, the algebra  $Oc(G)$  is not only a bimodule over  $A(G)$  but also a bimodule over the graph algebra of  $H$  where  $H$  is a graph with self-fusion on which  $A(G)$  acts. In the cases where  $G$  admits self-fusion, it is often so that  $H$  is  $G$  itself. In those cases we have a relative modular splitting formula: fusion matrices are still the same but the relative modular matrix  $M^{rel}$  is written in terms of the  $G$  graph (so it is of size  $d_G^2$  rather than  $d_A^2$ );  $M = E_0 M^{rel} E_0^T$ , where  $E_0$  is the first essential matrix (intertwiner). In the same way, toric matrices  $W$  of size  $d_A^2$  are replaced by relative toric matrices  $W^{rel}$  of size  $d_G^2$ . The modular splitting technique can be applied

as before, with the advantage that the size of tensors is greatly reduced. Once the relative matrices are found, we can retrieve the others by the relation  $W_{xy} = E_0 W^{rel}_{xy} E_0^T$ . Such an example is worked out in the last section of reference [11]

### 5.3 A dual bimodule structure?

Axioms for quantum groupoids are certainly self-dual, but the objects that we have at hand are not generic : they are quite special. In particular, if it is clear that  $Oc(G)$  is an  $A(G)$  bimodule, there is no obvious reason for  $A(G)$  to be an  $Oc(G)$  bimodule. If it were so, this action would be defined by a set of coefficients  $P_{xy}$ , with  $x \lambda y = \sum_{\mu} (P_{xy})_{\lambda\mu} \mu$ . The  $P_{xy}$  being of dimension  $d_{\mathcal{A}_k} \times d_{\mathcal{A}_k}$  and the bimodule associativity property  $(xx')\lambda(yy') = x(x'\lambda y)y'$  would lead to a double quantum symmetry equation:  $P_{x'y} P_{xy'} = \sum_{x''y''} O_{xx'}^{x''} O_{yy'}^{y''} P_{x''y''}$ . This equation taken at  $y = y' = 0$ , at  $x = x' = 0$  and at  $x = y = 0$  would itself lead to:  $P_{x'0} P_{x0} = \sum_{x''} O_{xx'}^{x''} P_{x''0}$ ,  $P_{0y} P_{0y'} = \sum_{y''} O_{yy'}^{y''} P_{0y''}$ ,  $P_{x'y'} = P_{x'0} P_{0y'} = P_{0y'} P_{x'0}$  and each set of matrices  $P_{x0}$  or  $P_{0y}$  would give respectively an anti-representation and a representation of dimension  $d_{\mathcal{A}_k} \times d_{\mathcal{A}_k}$  of the quantum symmetry algebra. Now, what could these  $P_{xy}$  matrices be? One obvious candidate is to set them equal to the toric matrices  $W_{xy}$ . The problem is that this choice cannot work since, as it can be checked on simple examples,  $W_{x'y'}$  is not equal to  $W_{x'0} W_{0y'}$  in general. Existence of a dual bimodule structure is not excluded, but if it exists, it cannot be defined by the toric matrices alone. Supposing the existence of such dual bimodule structure, it should also satisfy some compatibility conditions, like  $(\lambda(x(\mu(ya)))) = ((\lambda x \mu)(ya)) = (\lambda(x \mu y)a)$ , leading to the following set of relations:

$$S_y F_{\mu} S_x F_{\lambda} = \sum_z (V_{\lambda\mu})_{xz} S_y S_z = \sum_{\nu} (P_{xy})_{\mu\nu} F_{\nu} F_{\lambda}. \quad (25)$$

### 5.4 Realization of the Ocneanu quantum symmetries

In many cases  $Oc(G)$  can be written in terms of the tensor square of the graph algebras of some related graph  $K$  with self fusion, with the tensor product taken over a subalgebra, called the modular subalgebra  $J$ . In the simplest cases, i.e., when  $G$  has self fusion,  $K$  is  $G$  itself. The set of elements of  $J$  is determined by modular properties [9, 12, 13, 47]. Each vertex of an  $\mathcal{A}_k$  graph has a fixed modular operator value  $T$ . The vector space spanned by vertices of a  $G$  graph is a module over  $\mathcal{A}_k$ , and one can try to define a modular operator value on vertices of  $G$ . Suppose that the vertex  $a$  of  $G$  appears both in the branching rules (restriction map from  $\mathcal{A}_k$  to  $G$ ) of vertices  $\lambda$  and  $\mu$  of  $\mathcal{A}_k$ . The vertex  $a$  will have a well-defined modular operator value if the two values  $T(\lambda)$  and  $T(\mu)$  are equal. The set of vertices having this property is a subalgebra of the graph algebra of  $G$ , denoted  $J$ .

As already commented, non trivial multiplicities in the modular matrix lead to non commutativity for  $Oc(G)$ . This happens whenever  $G$  possesses classical symmetries<sup>9</sup>. In those

<sup>9</sup>By this we mean that, the unit vertex being chosen, the graph still contains a classical symmetry, making

cases, the algebraic realization of  $Oc(G)$  involves not only a tensor square over some sub-algebra but a cross product by an appropriate discrete group algebra [47]. The bimodule structure of  $Oc(G)$  over  $\mathcal{A}_k \otimes \mathcal{A}_k$  is thus related to the module structure of  $G$  over  $\mathcal{A}_k$ .

## 6 The $SU(3)$ system of graphs and their quantum symmetries

Starting with the complete list of modular invariants [22], the list of graphs was found by [17], slightly amended by [40]. We believe that a determination of the graph of quantum symmetries associated with the above was worked out in 2000 or before by A. Ocneanu (unpublished). We now present a compendium of results concerning not only these quantum symmetries but also several other results that use the concepts introduced in previous sections. In particular we give in most cases an algebraic realization of  $Oc(G)$  that allows one to perform calculations without having to use the graph of quantum symmetries. A detailed study of several cases has already been made available in the literature [13, 47] and details concerning the others will be published elsewhere [25, 27, 24]. Several graphs are displayed in figures 4 and 5.

### 6.1 The $\mathcal{A}$ series and its conjugated series

#### 6.1.1 The $\mathcal{A}$ series (graphs with self-fusion)

The  $\mathcal{A}_k$  graphs are the Weyl alcoves of  $SU(3)$  at level  $k$ . We have  $A(\mathcal{A}_k) = \mathcal{A}_k$ , so the annular matrices coincide with the fusion matrices:  $F_\lambda = N_\lambda$ . The algebra of quantum symmetries is realized as  $Oc(\mathcal{A}_k) = \mathcal{A}_k \dot{\otimes} \mathcal{A}_k$  where the tensor product is taken over  $\mathcal{A}_k$  with the identification  $\lambda \dot{\otimes} \mu \equiv \lambda\mu^* \dot{\otimes} 0$ . A basis of  $Oc(\mathcal{A}_k)$  is  $x = \lambda \dot{\otimes} 0$  and the dimension  $d_O = d_{\mathcal{A}_k}$ . The dual annular matrices are  $S_x = F_\lambda = N_\lambda$  and the double annular matrices are  $V_{\lambda\mu} = N_\lambda N_{\mu^*}$ . The modular invariant associated to the  $\mathcal{A}_k$  graph is diagonal  $M_{\lambda\mu} = \delta_{\lambda\mu}$ . We can easily check that  $(V_{\lambda\mu})_{00} = M_{\lambda\mu}$ . The two algebras  $\mathcal{BA}_k$  and  $\widehat{\mathcal{BA}}_k$  are isomorphic. We have  $d_x = d_\lambda$ , the quadratic and linear sum rules are trivially satisfied. In the  $SU(2)$  system, i.e. for  $ADE$  diagrams, the value of  $d_H = \sum_\lambda d_\lambda$  has been obtained independently, for all diagrams, by A. Ocneanu (unpublished) and by [45] and [9, 12]. It is easy to see, for instance, that for  $A_{k+1} = \mathcal{A}_k$  graphs, the following formula holds :  $d_H = \frac{(k+1)(k+2)(k+3)}{6}$ . Actually, setting  $r = k + 1$  (the number of vertices) and  $\kappa = k + 2$  (the usual Coxeter number), this formula also works for  $D$ -graphs and for exceptionals, when it is written as  $d_H = r\kappa(\kappa + 1)/6$ . It is interesting to notice that the same formula also gives the dimension of the Gelfand - Ponomarev preprojective algebra associated with the chosen graph (see [32]). For the  $SU(3)$  system of graphs (now  $\kappa = k + 3$ ) we observe that the dimension  $d_H = \sum_\lambda d_\lambda$  of graphs  $\mathcal{A}_k$  is given by the formula

$$d_H = \frac{(k+1)(k+2)(k+3)(k+4)(k+5)(k^2+6k+14)}{1680}. \quad (26)$$

---

impossible a direct computation of the table of multiplication.

### 6.1.2 The $\mathcal{A}^*$ modules (no self-fusion)

The  $\mathcal{A}_k^*$  graphs are the conjugated graphs of  $\mathcal{A}_k$ . Their vertices are the real vertices of  $\mathcal{A}_k$  (see for example [21, 46, 1]). We have  $A(\mathcal{A}_k^*) = \mathcal{A}_k$ . The algebra of quantum symmetries is realized as  $Oc(\mathcal{A}_k^*) = \mathcal{A}_k \dot{\otimes} \mathcal{A}_k$  where the tensor product is again taken over  $\mathcal{A}_k$  but now with the identification  $\lambda \dot{\otimes} \mu \equiv \lambda \mu \dot{\otimes} 0$ . A basis of  $Oc(\mathcal{A}_k^*)$  is again  $x = \lambda \dot{\otimes} 0$ , and we have  $d_O = d_{\mathcal{A}_k}$ . The dual annular matrices are  $S_x = F_\lambda$  and the double annular matrices are  $V_{\lambda\mu} = N_\lambda N_\mu$ . The modular invariant is  $M_{\lambda\mu} = \delta_{\lambda\mu^*}$ . The two algebras  $\mathcal{BA}_k$  and  $\widehat{\mathcal{BA}}_k$  are isomorphic. We have  $d_x = d_\lambda$ , the quadratic and linear sum rules are trivially satisfied.

## 6.2 The $\mathcal{D}$ series and the conjugated $\mathcal{D}^*$ series

The  $\mathcal{D}_k = \mathcal{A}_k/3$  graphs are orbifold graphs of the  $\mathcal{A}_k$  graphs. They are obtained from the action of the geometrical  $\mathbb{Z}_3$ -automorphism  $z$  (see Eq.(6)) on irreps of the  $\mathcal{A}_k$  graphs [31, 19, 17]. Vertices of  $\mathcal{A}_k$  that belong to the same orbit lead to a single vertex in the orbifold graph  $\mathcal{D}_k$ . When there is a fixed vertex under  $z$  (this happens when  $k \equiv 0 \pmod{3}$ ), this vertex is triplicated on the orbifold graph. Among all orbifold graphs  $\mathcal{D}_k$ , the  $\mathcal{D}_{3n}$  are the only ones that have self-fusion.

### 6.2.1 The $\mathcal{D}_k$ orbifold modules for $k \not\equiv 0 \pmod{3}$ (no self-fusion)

For  $k \not\equiv 0 \pmod{3}$ , the  $\mathcal{D}_k$  graphs have  $(k+1)(k+2)/6$  vertices. One can define a graph algebra with non negative integer structure constants for these graphs, but it is not compatible with the  $\mathcal{A}_k$  action. Therefore these graphs don't have self-fusion. The Ocneanu algebra is realized as  $Oc(\mathcal{D}_k) = \mathcal{A}_k \dot{\otimes} \mathcal{A}_k$  where the tensor product is again taken over  $\mathcal{A}_k$  but with the identification  $\lambda \dot{\otimes} \mu \equiv \lambda \rho(\mu^*) \dot{\otimes} 0$ , where  $\rho$  is the Gannon twist (see Eq.(7)). A basis of  $Oc(\mathcal{D}_k)$  is  $x = \lambda \dot{\otimes} 0$ , and we have  $d_O = d_{\mathcal{A}_k}$ . The dual annular matrices are  $S_x = F_\lambda$  and the double annular matrices are  $V_{\lambda\mu} = N_\lambda N_{\rho(\mu^*)}$ . The associated modular invariant is  $M_{\lambda\mu} = \delta_{\lambda\rho(\mu^*)}$ . The two algebras  $\mathcal{BD}_k$  and  $\widehat{\mathcal{BD}}_k$  are isomorphic. We have  $d_x = d_\lambda$ , the quadratic and linear sum rules are trivially satisfied. The dimensions  $d_\lambda$  of the blocks labelled by  $\lambda$  (or by  $x$ , which is the same here) satisfy  $d_\lambda(\mathcal{D}_k) = d_\lambda(\mathcal{A}_k)/3$ . The dimensions therefore satisfy  $\dim(\mathcal{BD}_k) = \dim(\mathcal{BA}_k)/9$ .

### 6.2.2 The $\mathcal{D}_k^*$ conjugated orbifold modules $k \not\equiv 0 \pmod{3}$ (no self-fusion)

The conjugated orbifold graphs  $\mathcal{D}_k^*$  are the unfolded (i.e. triplicated) graphs of the  $\mathcal{A}_k^*$  ones [17], i.e. their adjacency matrices are such that  $Ad(\mathcal{D}_k^*) = \sigma_{123} \otimes Ad(\mathcal{A}_k^*)$ , where  $\sigma_{123} = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}$  is the permutation matrix. These graphs are modules over the fusion algebras  $\mathcal{A}_k$ . The Ocneanu algebra is realized as  $Oc(\mathcal{D}_k^*) = \mathcal{A}_k \dot{\otimes} \mathcal{A}_k$  where the tensor product is again taken over  $\mathcal{A}_k$  but with the identification  $\lambda \dot{\otimes} \mu \equiv \lambda \rho(\mu) \dot{\otimes} 0$ , where  $\rho$  is the Gannon twist defined in Eq. (7). A basis of  $Oc(\mathcal{D}_k^*)$  is again  $x = \lambda \dot{\otimes} 0$ , and we have  $d_O = d_{\mathcal{A}_k}$ . The dual annular

matrices are  $S_x = F_\lambda$  and the double annular matrices are  $V_{\lambda\mu} = N_\lambda N_{\rho(\mu)}$ . The associated modular invariant is  $M_{\lambda\mu} = \delta_{\lambda\rho(\mu^*)}$ . The two algebras  $\mathcal{BD}_k^*$  and  $\widehat{\mathcal{BD}}_k^*$  are isomorphic. We have  $d_x = d_\lambda$ , the quadratic and linear sum rules are trivially satisfied. The dimensions  $d_\lambda$  of the blocks labelled by  $\lambda$  (or by  $x$ , which is the same here) satisfy  $d_\lambda(\mathcal{D}_k^*) = 3 d_\lambda(\mathcal{A}_k^*)$ . The dimensions therefore satisfy  $\dim(\mathcal{BD}_k^*) = 9 \dim(\mathcal{BA}_k^*)$ .

### 6.2.3 The $\mathcal{D}_k$ orbifolds for $k = 0 \bmod 3$ (self-fusion)

For  $k = 0 \bmod 3$ , the  $\mathcal{A}_k$  graphs have a central vertex  $\mathbb{Z}_3$ -invariant, which is triplicated on the orbifold graph. In this case  $\mathcal{D}_k$  graphs have  $(\frac{(k+1)(k+2)}{2} - 1)/3 + 3$  vertices, and they possess self-fusion. The associated modular invariant partition function is:

$$\mathcal{Z}(\mathcal{D}_k) = \frac{1}{3} \sum_{\lambda|t(\lambda)=0} |\chi_\lambda^k + \chi_{z(\lambda)}^k + \chi_{z^2(\lambda)}^k|^2 \quad (27)$$

The special vertex  $z$ -invariant on the  $\mathcal{A}_k$  graph leads to the presence of a coefficient equal to 3 in the modular invariant. Therefore the algebra of quantum symmetries of  $\mathcal{D}_{3n}$  is non-commutative. A realization is given by a semi-direct product  $Oc(\mathcal{D}_{3n}) = \mathcal{D}_{3n} \rtimes \mathbb{Z}_3$  (see [48]). The Ocneanu graph is made of 3 copies of the  $\mathcal{D}_{3n}$  graph, its dimension is  $d_O = (k+1)(k+2)/2 + 8$ . The quadratic sum rule is satisfied but the linear sum rule does not hold  $d_H \neq d_V$  (it may be recovered by introducing appropriate symmetry factors).

### 6.2.4 The $\mathcal{D}_k^*$ conjugate orbifolds for $k = 0 \bmod 3$ (no self-fusion)

The conjugate orbifold graphs  $\mathcal{D}_k^*$  are the unfolded (i.e. triplicated) graphs of the  $\mathcal{A}_k^*$  ones [17]. These graphs are modules over the fusion algebras  $\mathcal{A}_k$ . For  $k = 0 \bmod 3$ , the associated modular invariant partition function is

$$\mathcal{Z}(\mathcal{D}_k^*) = \frac{1}{3} \sum_{\lambda|t(\lambda)=0} (\chi_\lambda^k + \chi_{z(\lambda)}^k + \chi_{z^2(\lambda)}^k) (\overline{\chi_\lambda^k} + \overline{\chi_{z(\lambda)}^k} + \overline{\chi_{z^2(\lambda)}^k}) \quad (28)$$

Its algebra of quantum symmetries is also non-commutative, and can be realized as a conjugated version of semi-direct product  $Oc(\mathcal{D}_{3n}) = \mathcal{D}_{3n} \rtimes \mathbb{Z}_3$  (see [48]). Its dimension is  $d_O(\mathcal{D}_k^*) = d_O(\mathcal{D}_k)$ . The quadratic sum rule is satisfied but the linear sum rule does not hold  $d_H \neq d_V$  (it may be recovered by introducing appropriate symmetry factors).

## 6.3 Exceptional graphs with self-fusion and their modules

In the  $SU(3)$  family, we have three exceptional graphs with self-fusion, namely  $\mathcal{E}_5$ ,  $\mathcal{E}_9$  and  $\mathcal{E}_{21}$ . Diagrams  $\mathcal{E}_5$  and  $\mathcal{E}_{21}$  are generalizations of the two Dynkin diagrams  $E_6$  and  $E_8$ . We have also the module graphs  $\mathcal{E}_5^* = \mathcal{E}_5/3$  and  $\mathcal{E}_9^* = \mathcal{E}_9/3$  (they don't have self-fusion). Finally we have the exceptional graph  $\mathcal{D}_9^t$  obtained from the exceptional twist of the  $\mathcal{D}_9$  graph (a generalization of the  $E_7$  Dynkin diagram), together with the conjugated exceptional graph  $\mathcal{D}_9^{t*}$ .



### 6.3.1 The exceptional $\mathcal{E}_5$ graph (self-fusion)

The  $\mathcal{E}_5$  graph has self-fusion and has 12 vertices denoted  $1_i$  and  $2_j$  where  $i, j = 1, 2, \dots, 6$ . The unit vertex is  $1_0$  and the fundamental conjugated generators are  $2_1$  and  $2_2$  (for more details see [13] and [47]). Its quantum mass is  $m(\mathcal{E}_5) = 12(2 + \sqrt{2})$ . The associated modular invariant partition functions is:

$$\begin{aligned} \mathcal{Z}(\mathcal{E}_5) &= |\chi_{(0,0)}^5 + \chi_{(2,2)}^5|^2 + |\chi_{(0,2)}^5 + \chi_{(3,2)}^5|^2 + |\chi_{(2,0)}^5 + \chi_{(2,3)}^5|^2 \\ &+ |\chi_{(2,1)}^5 + \chi_{(0,5)}^5|^2 + |\chi_{(3,0)}^5 + \chi_{(0,3)}^5|^2 + |\chi_{(1,2)}^5 + \chi_{(5,0)}^5|^2. \end{aligned}$$

The modular subalgebra is  $J = \{1_i, i = 1, \dots, 6\}$  and a realization of the Ocneanu algebra is given by  $Oc(\mathcal{E}_5) = \mathcal{E}_5 \dot{\otimes}_J \mathcal{E}_5$ , with the identifications  $a \dot{\otimes}_J u b \equiv a u^* \dot{\otimes}_J b$ , for all  $u \in J$  and  $a, b \in \mathcal{E}_5$ . Conjugation on  $\mathcal{E}_5$  is defined as:  $1_0^* = 1_0$ ,  $1_5^* = 1_1$ ,  $1_4^* = 1_2$ ,  $1_3^* = 1_3$ ,  $2_0^* = 2_3$ ,  $2_1^* = 2_2$  and  $2_5^* = 2_4$  (it corresponds to the symmetry with respect to the vertical axis joining vertices  $1_0$  and  $1_3$  of the diagram  $\mathcal{E}_5$  given on Figure 4). Its dimension is 24 and a basis of  $Oc(\mathcal{E}_5)$  is given by  $a \dot{\otimes}_J 1_0$  and  $b \dot{\otimes}_J 2_0$ , for  $a, b \in \mathcal{E}_5$ . The chiral generators are  $2_1 \dot{\otimes}_J 1_0$  and  $1_0 \dot{\otimes}_J 2_1 \equiv 1_5 \dot{\otimes}_J 2_0$ . The left and right chiral subalgebras are  $L = \{a \dot{\otimes}_J 1_0\}$  and  $R = \{1_0 \dot{\otimes}_J a\}$ , and the ambichiral subalgebra is  $A = \{1_i \dot{\otimes}_J 1_0 \equiv 1_0 \dot{\otimes}_J 1_i^*\}$ . The quantum mass of  $Oc(\mathcal{E}_5)$  is  $m[Oc(\mathcal{E}_5)] = \frac{m[\mathcal{E}_5]m[\mathcal{E}_5]}{m[J]} = m[\mathcal{A}_5] = 48(3 + \sqrt{2})$ . The linear and quadratic sum rules hold and read  $d_H = d_V = 720$ ,  $\dim(\mathcal{BE}_5) = 29376$ , respectively.

### 6.3.2 The exceptional module of the $\mathcal{E}_5$ graph (no self-fusion)

The  $\mathcal{E}_5^* = \mathcal{E}_5/3$  is the  $\mathbb{Z}_3$ -orbifold graph of  $\mathcal{E}_5$ , it has 4 vertices. It is a module over  $\mathcal{A}_5$  and over  $\mathcal{E}_5$ . In particular it has the same norm  $\beta = [3]_q = 1 + \sqrt{2}$  as  $\mathcal{A}_5$  and  $\mathcal{E}_5$ . Its quantum mass is  $m(\mathcal{E}_5^*) = m(\mathcal{E}_5)/3 = 4(2 + \sqrt{2})$ . The associated modular invariant partition function is:

$$\begin{aligned} \mathcal{Z}(\mathcal{E}_5^*) &= |\chi_{(0,0)}^5 + \chi_{(2,2)}^5|^2 + |\chi_{(3,0)}^5 + \chi_{(0,3)}^5|^2 + (\chi_{(0,2)}^5 + \chi_{(3,2)}^5)(\overline{\chi_{(2,0)}^5 + \chi_{(2,3)}^5}) \\ &+ (\chi_{(2,0)}^5 + \chi_{(2,3)}^5)(\overline{\chi_{(0,2)}^5 + \chi_{(3,2)}^5}) + (\chi_{(1,2)}^5 + \chi_{(5,0)}^5)(\overline{\chi_{(0,5)}^5 + \chi_{(2,1)}^5}) \\ &+ (\chi_{(2,1)}^5 + \chi_{(0,5)}^5)(\overline{\chi_{(1,2)}^5 + \chi_{(5,0)}^5}). \end{aligned}$$

The Ocneanu algebra is  $Oc(\mathcal{E}_5^*) = \mathcal{E}_5 \dot{\otimes}_J \mathcal{E}_5$  where the tensor product is taken over the modular subalgebra  $J$  of  $\mathcal{E}_5$  but with the identifications  $a \dot{\otimes}_J u b \equiv a u \dot{\otimes}_J b$ , for all  $u \in J$  and  $a, b \in \mathcal{E}_5$ . The two algebras  $Oc(\mathcal{E}_5)$  and  $Oc(\mathcal{E}_5^*)$  are isomorphic but their realization in terms of tensor products are different. Here the right chiral generator is  $1_0 \dot{\otimes}_J 2_1 \equiv 1_1 \dot{\otimes}_J 2_0$ . The quantum mass is  $m(Oc(\mathcal{E}_5^*)) = m(Oc(\mathcal{E}_5))$ . The dimensions of the blocks labelled by  $\lambda$  and  $x$  satisfy  $d_\lambda(\mathcal{E}_5^*) = d_\lambda(\mathcal{E}_5)/3$  and  $d_x(\mathcal{E}_5^*) = d_x(\mathcal{E}_5)/3$ . The linear and quadratic sum rules hold and read  $d_H = d_V = 720/3 = 240$  and  $\dim(\mathcal{BE}_5^*) = \dim(\mathcal{BE}_5)/9 = 3264$ .

### 6.3.3 The exceptional $\mathcal{E}_9$ graph (self-fusion)

The  $\mathcal{E}_9$  graph has self-fusion and possesses 12 vertices denoted  $0_i, 1_i, 2_i$  and  $3_i$  where  $i = 0, 1$  or  $2$ . Its quantum mass is  $m(\mathcal{E}_9) = 36(2 + \sqrt{3})$ . The associated modular invariant partition function is:

$$\mathcal{Z}(\mathcal{E}_9) = |\chi_{(0,0)}^9 + \chi_{(0,9)}^9 + \chi_{(9,0)}^9 + \chi_{(1,4)}^9 + \chi_{(4,1)}^9 + \chi_{(4,4)}^9|^2 + 2|\chi_{(2,2)}^9 + \chi_{(2,5)}^9 + \chi_{(5,2)}^9|^2$$

The presence of the factor 2 in the second term of the modular invariant indicates that the Ocneanu algebra  $Oc(\mathcal{E}_9)$  is non commutative. It is isomorphic to a direct sum of 36 one-dimensional blocks of  $\mathbb{C}$  and of 9 copies of 2-dimensional matrices  $M_2(\mathbb{C})$ , its dimension is 72. The modular subalgebra is  $J = \{0_0, 1_0, 2_0\}$  and the Ocneanu algebra  $Oc(\mathcal{E}_9)$  involves  $\mathcal{E}_9 \otimes_J \mathcal{E}_9$  and a non commutative matrix complement (see [27] for more details). The Ocneanu graph is made of  $12 \times 6 = 72$  vertices, corresponding to 3 copies of the  $\mathcal{E}_9$  graph and 3 copies of its module graph  $\mathcal{E}_9/3$ . The quantum mass is  $m(Oc(\mathcal{E}_9)) = \frac{m(\mathcal{E}_9)m(\mathcal{E}_9)}{m(J)} = m(\mathcal{A}_9) = 432(7 + 4\sqrt{3})$ , where  $m(J) = 3$ . Note that the quadratic sum rule can be checked ( $\dim(\mathcal{B}\mathcal{E}_9) = \sum_{\lambda} d_{\lambda}^2(\mathcal{E}_9) = \sum_x d_x^2(\mathcal{E}_9) = 518976$ ) but the linear sum rule does not hold:  $d_H = 4656$  but  $d_V = 5448$ .

### 6.3.4 The exceptional module of the $\mathcal{E}_9$ graph (no self-fusion)

The  $\mathcal{E}_9^* = \mathcal{E}_9/3$  graph is a module over the graph algebra  $\mathcal{A}_9$  and over the graph algebra  $\mathcal{E}_9$ . It has the same norm  $\beta = [3]_q = 1 + \sqrt{3}$  as  $\mathcal{A}_9$  and  $\mathcal{E}_9$ . The  $\mathcal{E}_9^*$  graph is associated to the same modular invariant as  $\mathcal{E}_9$ . Furthermore, the Ocneanu algebra  $Oc(\mathcal{E}_9^*)$  is isomorphic to  $Oc(\mathcal{E}_9)$ . But the module structures of  $\mathcal{E}_9^*$  over  $\mathcal{A}_9$  and over  $Oc(\mathcal{E}_9) \equiv Oc(\mathcal{E}_9^*)$  are not the same as for  $\mathcal{E}_9$ : the annular matrices  $F_{\lambda}$  and dual annular matrices  $S_x$  differ from those of  $\mathcal{E}_9$ . The quadratic sum rule hold and read  $\dim(\mathcal{B}\mathcal{E}_9^*) = 754272$ , but the linear sum rule does not hold:  $d_H = 5616$  but  $d_V = 6552$ .

### 6.3.5 The exceptional $\mathcal{E}_{21}$ graph (self-fusion)

The  $\mathcal{E}_{21}$  graph has self-fusion and possesses 24 vertices denoted  $0, 2, \dots, 23$ . The unit vertex is 0, the conjugated generators are 1 and 2. Complex conjugation corresponds to the symmetry with respect to the horizontal axis joining vertices 0 and 21 of the  $\mathcal{E}_{21}$  graph given on figure 4. Triality is equal to the labels taken modulo 3. The norm of the  $\mathcal{E}_{21}$  graph is  $\beta = \frac{1}{2}(1 + \sqrt{2} + \sqrt{6})$ . Actually all quantum dimensions are of the kind  $(a, b, c, d) = a + b\sqrt{2} + c\sqrt{3} + d\sqrt{6}$ , for appropriate values of  $a, b, c, d$ . The quantum mass is  $m(\mathcal{E}_{21}) = 24(18 + 10\sqrt{3} + \sqrt{6(97 + 56\sqrt{3})})$ . The associated modular invariant partition function is:

$$\begin{aligned} \mathcal{Z}(\mathcal{E}_{21}) &= |\chi_{(0,0)}^{21} + \chi_{(4,4)}^{21} + \chi_{(6,6)}^{21} + \chi_{(10,10)}^{21} + \chi_{(0,21)}^{21} + \chi_{(21,0)}^{21} + \chi_{(1,10)}^{21} + \chi_{(10,1)}^{21} + \chi_{(4,13)}^{21} \\ &+ \chi_{(13,4)}^{21} + \chi_{(6,9)}^{21} + \chi_{(9,6)}^{21}|^2 + |\chi_{(0,6)}^{21} + \chi_{(6,0)}^{21} + \chi_{(0,15)}^{21} + \chi_{(15,0)}^{21} + \chi_{(4,7)}^{21} + \chi_{(7,4)}^{21} \\ &+ \chi_{(4,10)}^{21} + \chi_{(10,4)}^{21} + \chi_{(6,15)}^{21} + \chi_{(15,6)}^{21} + \chi_{(7,10)}^{21} + \chi_{(10,7)}^{21}|^2 \end{aligned}$$

The modular subalgebra is  $J = \{0, 21\}$  and a realization of the Ocneanu algebra is  $Oc(\mathcal{E}_{21}) = \mathcal{E}_{21} \dot{\otimes}_J \mathcal{E}_{21}$ , with the identifications  $a \dot{\otimes}_J u b \equiv a u^* \dot{\otimes}_J b$ , for all  $u \in J$  and  $a, b \in \mathcal{E}_{21}$ . The Ocneanu graph involves 12 copies of  $\mathcal{E}_{21}$ . The dimension of  $Oc(\mathcal{E}_{21})$  is 288 (see [13, 47] for more details). Its quantum mass is given by  $m[Oc(\mathcal{E}_{21})] = \frac{m[\mathcal{E}_{21}]m[\mathcal{E}_{21}]}{m[J]} = m[\mathcal{A}_{21}]$ , where  $m[J] = 2$ . Numerically  $m[Oc(\mathcal{E}_{21})] = 1728(201 + 142\sqrt{2} + 116\sqrt{3} + 82\sqrt{6})$ . The linear and quadratic sum rules hold and read  $d_H = d_V = 288\,576$ ,  $\dim(\mathcal{B}\mathcal{E}_{21}) = 480\,701\,952$ , respectively.

### 6.3.6 The twisted exceptional $\mathcal{D}_9^t$ (no self-fusion)

The  $\mathcal{D}_9^t$  graph is a module over the graph algebra  $\mathcal{A}_9$  and over the graph algebra  $\mathcal{D}_9$ . It is associated to the following modular invariant partition function:

$$\begin{aligned} \mathcal{Z}(\mathcal{D}_9^t) &= |\chi_{(0,0)}^9 + \chi_{(9,0)}^9 + \chi_{(0,9)}^9|^2 + |\chi_{(3,0)}^9 + \chi_{(6,3)}^9 + \chi_{(0,6)}^9|^2 + |\chi_{(0,3)}^9 + \chi_{(6,0)}^9 + \chi_{(3,6)}^9|^2 \\ &+ |\chi_{(2,2)}^9 + \chi_{(5,2)}^9 + \chi_{(2,5)}^9|^2 + |\chi_{(4,4)}^9 + \chi_{(4,1)}^9 + \chi_{(1,4)}^9|^2 + 2|\chi_{(3,3)}^9|^2 \\ &+ \left[ (\chi_{(1,1)}^9 + \chi_{(7,1)}^9 + \chi_{(1,7)}^9) \overline{\chi_{(3,3)}^9} + h.c. \right] \end{aligned}$$

The graph  $\mathcal{D}_9^t$  appears as a module of its own algebra of quantum symmetries (calculated from the modular splitting equation). It is a generalization of the  $E_7$  graph<sup>10</sup> of the  $SU(2)$  system. Its quantum mass is  $m(\mathcal{D}_9^t) = 72(2 + \sqrt{3})$ .  $Oc(\mathcal{D}_9^t)$  is obtained via an anti-automorphism called the exceptional ambichiral twist  $\xi$ , which acts on vertices of the modular subalgebra  $J = \{0_0, 2_0, 3_0, 3_0^1, 4_0, 5_0, \alpha_0^1, \alpha_0^2, \alpha_0^3\}$  of  $\mathcal{D}_9$  (see Figure 4), such that  $\xi(2_0) = \alpha_0^2$ ,  $\xi(\alpha_0) = 2_0$  and  $\xi(u) = u$  for all others  $u \in J$ . The Ocneanu algebra  $Oc(\mathcal{D}_9^t)$  involves  $\mathcal{D}_9 \dot{\otimes}_J \mathcal{D}_9$  and a non commutative matrix complement. We identify  $a \dot{\otimes}_J u b \equiv a \xi(u^*) \dot{\otimes}_J b$  for all  $u \in J$  and  $a, b \in \mathcal{D}_9$ . Its dimension is 55 and the quantum mass is  $m(Oc(\mathcal{D}_9^t)) = m(\mathcal{A}_9) = 432(7 + 4\sqrt{3})$ . The dimension is  $\dim(\mathcal{B}\mathcal{D}_9^t) = 1\,167\,355$ .

### 6.3.7 The twisted conjugate exceptional $\mathcal{D}_9^{t*}$ (no self-fusion)

The  $\mathcal{D}_9^{t*}$  graph is a module graph over the graph algebras  $\mathcal{A}_9$ ,  $\mathcal{D}_9$  and also  $\mathcal{D}_9^t$ . The modular invariant partition function associated to this graph is:

$$\begin{aligned} \mathcal{Z}(\mathcal{D}_9^{t*}) &= |\chi_{(0,0)}^9 + \chi_{(9,0)}^9 + \chi_{(0,9)}^9|^2 + |\chi_{(2,2)}^9 + \chi_{(5,2)}^9 + \chi_{(2,5)}^9|^2 + |\chi_{(4,4)}^9 + \chi_{(4,1)}^9 + \chi_{(1,4)}^9|^2 \\ &+ 2|\chi_{(3,3)}^9|^2 + \left[ (\chi_{(0,3)}^9 + \chi_{(6,0)}^9 + \chi_{(3,6)}^9) (\overline{\chi_{(3,0)}^9} + \overline{\chi_{(6,3)}^9} + \overline{\chi_{(0,6)}^9}) + h.c. \right] \\ &+ \left[ (\chi_{(1,1)}^9 + \chi_{(7,1)}^9 + \chi_{(1,7)}^9) \overline{\chi_{(3,3)}^9} + h.c. \right] \end{aligned}$$

The  $\mathcal{D}_9^{t*}$  graph appears as a module of its own algebra of quantum symmetries, which is also obtained via the exceptional ambichiral twist  $\xi$  acting on vertices of  $J \subset \mathcal{D}_9$ . The Ocneanu algebra  $Oc(\mathcal{D}_9^{t*})$  involves also  $\mathcal{D}_9 \dot{\otimes}_J \mathcal{D}_9$  and a non commutative matrix complement, but with the identifications  $a \dot{\otimes}_J u b = a \xi(u) \dot{\otimes}_J b$  for all  $u \in J$  and  $a, b \in \mathcal{D}_9$ . Its dimension

<sup>10</sup>The  $E_7$  graph should better be called  $\mathcal{D}_{16}^t$ .




Figure 4: Some graphs with self-fusion: The  $\mathcal{A}_k$  series,  $\mathcal{D}_9$ ,  $\mathcal{E}_5$ ,  $\mathcal{E}_9$  and  $\mathcal{E}_{21}$ .

is 55 and the quantum mass is  $m(\text{Oc}(\mathcal{D}_9^{t*})) = m(\text{Oc}(\mathcal{D}_9^t)) = m(\mathcal{A}_9)$ . The dimension is  $\dim(\mathcal{BD}_9^{t*}) = 531\,435$ .

## 7 Comments

**Overall features of quantum groupoids and graphs associated to higher Coxeter-Dynkin systems** For an  $SU(n)$  system of graphs, one expects the following pattern. The family of  $\mathcal{A}_k$  graphs is easily obtained by truncation of the Weyl chambers at level  $k$ ; such  $\mathcal{A}_k$  graphs involve several types of oriented lines (one for each fundamental representation of  $SU(n)$ ). Then one can obtain several other families by using the existence of automorphisms such as complex conjugacy (leading to the  $\mathcal{A}_k^*$  series),  $\mathbb{Z}_p$  symmetries (leading to the orbifold  $\mathcal{D}_k[p] = \mathcal{A}_k/p$  series), or a combination of these two automorphisms (leading to the  $\mathcal{D}_k^*[p]$  series). From our experience with small values of  $n$ , we expect rather different families of  $\mathcal{D}$  graphs, depending on whether  $n$  is even or odd. For  $SU(2)$ , orbifold graphs  $\mathcal{D}_k[2] = \mathcal{D}_{\frac{k}{2}+2}$  exist if  $k = 0, 2 \pmod{4}$ , and they have self-fusion whenever  $k = 0 \pmod{4}$ . For  $SU(3)$ , orbifold




Figure 5: Some module graphs without self-fusion:  $\mathcal{A}_4^*$ ,  $\mathcal{D}_4$ ,  $\mathcal{D}_4^*$ ,  $\mathcal{E}_5^*$ ,  $\mathcal{E}_9^*$ ,  $\mathcal{D}_9^t$ , and  $\mathcal{D}_9^{t*}$ .

graphs  $\mathcal{D}_k[3]$  exist for all  $k$ , and they have self-fusion whenever  $k = 0 \pmod 3$ . For  $SU(4)$ , and according to [40], we have orbifold graphs of type  $\mathcal{D}_k[2]$  for all  $k$ , and they have self-fusion whenever  $k = 0 \pmod 2$ , but we have also orbifold graphs of type  $\mathcal{D}_k[4]$  for  $k = 0, 2, 6 \pmod 8$ , and they have self-fusion whenever  $k = 0 \pmod 8$ .

For  $\mathcal{A}_k$  and  $\mathcal{A}_k^*$  series, the algebra of quantum symmetries can be determined from the tensor square of the graph algebra  $\mathcal{A}_k$ , suitably quotiented. When  $\mathcal{D}_k$  does not have self-fusion, its algebra of quantum symmetries can also be determined from the tensor square of the graph algebra  $\mathcal{A}_k$ , suitably quotiented with the help of appropriate generalizations of the Gannon twist. This is also the case for its corresponding conjugated series. When  $\mathcal{D}_k$  graph has self-fusion, its algebra of quantum symmetries (which, in this case, is non commutative) can be obtained as a cross-product of the graph algebra of  $\mathcal{D}_k$  by the cyclic group  $\mathbb{Z}_p$ ; this is also the case for the corresponding conjugate series. In any of these cases, the associated modular invariant is easy to obtain from the  $\mathcal{A}$  modular invariant at same level.

For a given system, it seems that one can always find a (unique) exceptional graph  $\mathcal{D}^t$ , without self-fusion, whose algebra of quantum symmetries is equal to the quotient of the tensor square of a particular  $\mathcal{D}$  graph by an exceptional automorphism (this generalizes the  $(E_7, D_{10})$  situation of the  $SU(2)$  family). The graph  $\mathcal{D}^t$  itself is then recognized as a module over its algebra of quantum symmetries. Determination of this automorphism can be found by looking at the values of the modular operator  $T$  on vertices of the corresponding  $\mathcal{A}$  graph and the induction-restriction rules from  $\mathcal{A}$  to  $\mathcal{D}$  [13]. Same discussion for the corresponding conjugated graph  $\mathcal{D}^{t*}$ .

We are then left with the other exceptional graphs. They may admit self-fusion or not. When they don't, they are orbifolds of those exceptionals that enjoy self-fusion. Graphs with self-fusion are called “quantum subgroups” by A. Ocneanu, the others being only “quantum modules”. Those exceptional subgroups are  $E_6 \equiv \mathcal{E}_{10}$  and  $E_8 \equiv \mathcal{E}_{28}$  for the  $SU(2)$  system,

$\mathcal{E}_5$ ,  $\mathcal{E}_9$  and  $\mathcal{E}_{21}$  for the  $SU(3)$  system and  $\mathcal{E}_4$ ,  $\mathcal{E}_6$  and  $\mathcal{E}_8$  for the  $SU(4)$  system. Their algebra of quantum symmetries may be commutative or not. Non commutativity can be deduced, either from the presence of integer entries bigger than 1 in the modular invariant, or from the existence of non trivial classical symmetries in the graph itself (see footnote in section 5.4). When the algebra of quantum symmetries  $Oc(G)$  is commutative, like for  $E_6$  and  $E_8$  in the  $SU(2)$  system, or like for  $\mathcal{E}_5$ ,  $\mathcal{E}_{21}$  in the  $SU(3)$  system, it is easy to obtain the corresponding toric matrices and  $Oc(G)$  itself without having to solve the modular splitting equation, because, in these cases, one obtains  $Oc(G)$  as a tensor square of  $G$  itself over the modular subalgebra  $J$  which can be determined by using the properties of the modular generator  $T$  under restriction-induction (see [13]). Of course, it is always advisable to check that the obtained result satisfies the modular splitting equation. If, however, the algebra of quantum symmetries of this exceptional graph with self-fusion is non commutative (like for the  $\mathcal{E}_9$  case), the determination of  $Oc(G)$  becomes quite involved and the only method we can think of is again to use the modular splitting technique.

Once the exceptional graphs with self-fusion are known, it is not too difficult to obtain the exceptional modules : they are quotients or orbifolds of the former and often appear as particular subspaces of  $Oc(G)$ .

Finally, let us mention that when the graph  $G$  is a priori known, and whenever the vertex  $x$  of  $Oc(G)$  can be written as  $a \dot{\otimes} b$ , with  $a, b \in G$ , it is usually possible to obtain (or recover) the toric matrices  $W_{x0}$  from the annular or essential matrices, see for instance [12] or [47]. This method, first presented in [9], is particularly easy to implement when one considers generalizations of the exceptional graphs with self-fusion  $E_6$  and  $E_8$  (i.e.,  $\mathcal{E}_5$  and  $\mathcal{E}_{21}$  for the  $SU(3)$  system), since  $Oc(G) = G \dot{\otimes}_J G$ , in those cases. One obtains  $W_{x0} = \sum_{c \in J} (F_\lambda)_{ac} (F_\lambda)_{bc} = E_a \cdot ((E_b)^{red})^T$ , where the reduced essential matrices  $E_b^{red}$  are obtained from the  $E_b$  by keeping the matrix elements of those columns corresponding to the modular subalgebra  $J$  and putting all others entries to zero.

**Graphs from modular invariants.** One possibility is to rely on a given classification of the modular invariants. Such a classification exists for  $SU(2)$  [7] and  $SU(3)$  [22] but is not available for  $SU(n)$  when  $n > 3$ . However there are arguments showing that the level of exceptionals cannot be too high [41], so that it is enough to explore a sizeable list of possibilities. Once a modular invariant is known, one can use the modular splitting technique and find the algebra  $Oc(G)$ . Generically, the Ocneanu graph involves one or several copies of the graph  $G$  itself and of its modules; this may not be so in special cases, for instance the  $D_{odd}$  case of the  $SU(2)$  system or in the conjugated series of the  $SU(3)$  system, but then, other techniques of determination can be used (cf the above discussion). Once the graph  $G$  is obtained, one has still to check that the obtained result gives rise to a “good” theory of representations (here  $SU(3)$ ); otherwise, it should be discarded. We believe that the precise meaning of this sentence is that the obtained graph should give rise to a Kuperberg spider

[30]; another possibility is to use the existence of a self-connection, as defined by A. Ocneanu in [40]. As already mentioned, we believe that the two notions coincide but it is clear that some more work is needed in this direction. The list of graphs expected to provide an answer to the  $SU(4)$  classification problem is given in [40].

**Conformal embeddings** Another possibility leading to interesting candidates for graphs  $G$  of higher Coxeter-Dynkin systems is to use the existence of conformal embeddings of affine algebras – a subject that we did not touch in this paper. One should be aware that 1) List of modular invariants, 2) List of conformal embeddings, 3) List of graphs belonging to higher Coxeter-Dynkin systems (or defining Ocneanu quantum groupoids) are distinct problems.

It happens that, for  $SU(2)$  and  $SU(3)$ , all exceptional graphs with self-fusion correspond to particular conformal embeddings, but other such embeddings lead to orbifolds or to members (with small level) of the  $\mathcal{D}$  series. In the case of  $SU(4)$ , it seems that there is one exceptional graph with self-fusion not associated with any conformal embedding.

Conformal embeddings of affine algebras at level  $k$  of the type  $\widehat{su}(n)_k \subset \hat{g}_1$ , where  $g$  is a simple Lie algebra, simply laced or not, can be associated with graphs that are candidates to become members, at level  $k$ , of the Coxeter-Dynkin system of  $SU(n)$ . The condition to be conformal imposes equality of the central charges :

$$\frac{(n^2 - 1)k}{k + n} = \frac{\dim(g)}{1 + \kappa(g)} \quad (29)$$

where  $\dim(g)$  is the dimension of  $g$  and  $\kappa(g)$  its *dual* Coxeter number. This equation is easy to solve for all  $SU(n)$  systems. In the case  $n = 2$  there are three non trivial solutions:  $E_6$  ( $\equiv \mathcal{E}_{10}$ ), for  $g = B_2 = spin(5)$ , then  $E_8$  ( $\equiv \mathcal{E}_{28}$ ) for  $g = G_2$  and finally  $D_4$  ( $\equiv \mathcal{D}_4$ ), for  $g = A_2 = su(3)$ . In the case  $n = 3$  there are many more solutions; let us just mention those that give rise to exceptional graphs with self-fusion :  $\mathcal{E}_5$  for  $g = A_5 = su(6)$ , then  $\mathcal{E}_9$  for  $g = A_6 = su(7)$  and finally  $\mathcal{E}_{21}$  for  $g = E_7$ .

**Other generalizations.** The algebra of quantum symmetries described in the previous section refers to quantum groupoids for which a basis of matrix units, for the vertical product, is made of double triangles of type  $GGAGG$ , where  $G$  is any graph of the system ( $A$ -type,  $D$ -type, exceptional type etc. ). However one may replace these double triangles by others, of type  $GGKGG$ , whenever  $G$  is a  $K$  module. This was apparently not studied.

**About the definitions of  $Oc(G)$**  The most pleasant definition of  $Oc(G)$  is to take it as the algebra of characters (or irreps) for the horizontal product on  $\widehat{\mathcal{B}}G$ . This amounts to consider the center of  $\widehat{\mathcal{B}}G$  (for the horizontal multiplication  $\widehat{\circ}$ ) and analyse its structure when endowed with a product inherited from the vertical multiplication on  $\mathcal{B}G$ . However, to determine it in this way requires a priori the calculation of several sets (finite but huge) of generalized  $6J$  symbols. It seems that nobody ever did it this way (the family of  $6J$  symbols is not

even known for the exceptional cases of the  $SU(2)$  system !). Rather, the generators  $O_x$  were obtained as explained in step II of the modular splitting technique. A clear discussion relating these two types of concepts would be welcome.

**Frontiers.** The possibility of associating higher order algebraic systems (somehow generalizing universal enveloping algebras and their root systems) to graphs that are members of higher Coxeter-Dynkin families is certainly a fascinating perspective, which was not discussed in this paper.

**Conclusion.** The quantum groupoid aspects of these systems are still largely under-studied. As already stated previously, and in agreement with popular wisdom, every graph  $G$  belonging to an  $SU(n)$  system should give rise, and conversely, to an “Ocneanu quantum groupoid”. All together these objects constitute a particular family of finite dimensional weak Hopf algebras. However, many general properties still need clarification and every single particular diagram should deserve more study – for instance the explicit determination of the different types of cells (generalized  $6J$  symbols), is an open problem.

**Acknowledgements** D.H. and E.H.T. are grateful to FRUNAM and CIRM (Marseille) for supporting their stay at CIRM where part of this work was done. R. C. is grateful to the pole RENAPT to support his stay at LPTP in Oujda. G.S. would like to thank AUF – Agence Universitaire de la Francophonie – for financial support.

## References

- [1] R. E. Behrend, D. E. Evans, *Integrable lattice models for conjugate  $A_n^{(1)}$* , J. Phys. **A 37** (2004) 2937–2948, hep-th/0309068.
- [2] J. Böckenhauer, D. E. Evans, *Modular invariants, graphs and  $\alpha$ -induction for nets of subfactors I*, Commun. Math. Phys. **197** (1998) 361–386, hep-th/9801171; *II*, Commun. Math. Phys. **200** (1999) 57–103, hep-th/9805023; *III*, **205** (1999) 183–200, hep-th/9812110.
- [3] J. Böckenhauer, D. E. Evans, Y. Kawahigashi, *On  $\alpha$ -induction, chiral generators and modular invariants for subfactors*, Commun. Math. Phys. **208** (1999) 429–487, math.OA/9904109.
- [4] G. Böhm, K. Szlachányi, *A coassociative  $C^*$  quantum group with non-integral dimensions*, Lett. Math. Phys. **38** (1996) 437–456, q-alg/9509008.
- [5] G. Böhm, F. Nill, K. Szlachányi, *Weak Hopf Algebras I. Integral theory and  $C^*$  structure*, J. Algebra **221** (1999) 385–438, math.QA/9805116.



- [6] D. Calaque, P. Etingof, *Lectures on tensor categories*, math.QA/0401246.
- [7] A. Cappelli, C. Itzykson, J.-B. Zuber, *The ADE classification of minimal and  $A_1^{(1)}$  conformal invariant theories*, Commun. Math. Phys. **113** (1987) 1–26.
- [8] C.H. Otto Chui, C. Mercat, P. A. Pearce, *Integrable and Conformal Twisted Boundary Conditions for  $sl(2)$  A-D-E Lattice Models*, J.Phys. **A36** (2003) 2623–2662, hep-th/0210301.
- [9] R. Coquereaux, *Notes on the quantum tetrahedron*, Moscow Math. J. **2**, no.1 (2002) 41–80, math-ph/0011006.
- [10] R. Coquereaux, *The  $A_2$  Ocneanu quantum groupoid*, from the talk: "Quantum groupoids and Ocneanu Bialgebras for Coxeter-Dynkin systems" given at the XV Colloquio Latinoamericano de Algebra, Cocoyoc, Mexico, July 20th-26th, 2003. Contemporary Mathematics **376** (2005), hep-th/0311151.
- [11] R. Coquereaux, E. Isasi, *On quantum symmetries of the non-ADE graph  $F_4$* , Adv. Theor. Math. Phys. **8** (2004) 955–985, hep-th/0409201.
- [12] R. Coquereaux, G. Schieber, *Twisted partition functions for ADE boundary conformal field theories and Ocneanu algebra of quantum symmetries*, J. of Geom. and Phys. **781** (2002) 1–43, hep-th/0107001.
- [13] R. Coquereaux, G. Schieber, *Determination of quantum symmetries for higher ADE systems from the modular  $T$  matrix*, J. of Math. Physics **44** (2003) 3809–3837, hep-th/0203242.
- [14] R. Coquereaux, R. Trinchero, *On quantum symmetries of ADE graphs*, Adv. Theor. Math. Phys. **8** (2004) 189–216, hep-th/0401140.
- [15] R. Coquereaux, M. Huerta, *Torus structure on graphs and twisted partition functions for minimal and affine models*, J. of Geom. and Phys. **48** (2003) 580–634, hep-th/0301215.
- [16] P. Di Francesco, *Integrable lattice models, graphs and modular invariants conformal field theories*, Int. J. Mod. Phys. **A7** (1992) 407–500.
- [17] P. Di Francesco, J.-B. Zuber,  *$SU(N)$  Lattice integrable models associated with graphs*, Nucl. Phys **B338** (1990) 602–646;  *$SU(N)$  Lattice Integrable Models and Modular Invariance*, Recent Developments in Conformal Field Theories, Trieste Conference (1989), S. Randjbar-Daemi, E. Sezgin, J.-B. Zuber eds., World Scientific (1990).
- [18] D.E. Evans, P.R. Pinto, *Subfactor realization of modular invariants*, Commun. Math. Phys. **237** (2003) 309–363, math.OA/0309174.

- [19] P. Fendley, P. Ginsparg, *Non-critical orbifolds*, Nucl. Phys. **B324** (1989) 549–580.  
P. Fendley, *New exactly solvable models*, J. Phys. **A22** (1989) 4633–4642.
- [20] J. Fuchs, I. Runkel, C. Schweigert, *TFT construction of RCFT correlators I: Partition functions*, Nucl. Phys. **B646** (2002) 353–497, hep-th/0204148; *II: Unoriented world sheets*, Nucl. Phys. **B678** (2004) 511–637, hep-th/0306164; *III: Simple currents*, Nucl. Phys. **B694** (2004) 277–353, hep-th/0403157; *IV: Structure constants and correlation functions*, Nucl. Phys. **B715** (2005) 539–638, hep-th/0412290.
- [21] M. R. Gaberdiel, T. Gannon, *Boundary states for WZW models*, Nucl. Phys. **B639** (2002) 471–501, hep-th/0202067.
- [22] T. Gannon, *The classification of affine  $su(3)$  modular invariant partition functions*, Commun. Math. Phys. **161** (1994) 233–263, hep-th/9212060; *The classification of  $SU(3)$  modular invariants revisited*, Annales de l’Institut Poincaré, Phys. Theor. **65** (1996) 15–55, hep-th/9404185.
- [23] T. Gannon, *Modular data: the algebraic combinatorics of conformal field theory*, math.QA/0103044.
- [24] D. Hammaoui, PhD thesis, in preparation, LPTP, Université Mohamed I, Oujda, Maroc.
- [25] D. Hammaoui, G. Schieber, E.H. Tahri, *Higher Coxeter graphs associated to affine  $su(3)$  modular invariants*, J. Phys **A38** (2005) 8259–8286, hep-th/0412102.
- [26] D. Hammaoui, G. Schieber, E.H. Tahri, *Quantum symmetries of higher Coxeter graphs associated to affine  $su(3)$  modular invariants*, in preparation.
- [27] E. Isasi, G. Schieber, *From the modular invariant to graphs: the modular splitting method*, in preparation.
- [28] V. G. Kăc, D. H. Peterson, *Infinite-dimensional Lie algebras, theta functions and modular forms.*, Adv. Math. **53** (1984) 125–264.
- [29] V. Kodiyalam, V. S. Sunder, *Flatness and fusion coefficients*, Pacific J. of Math. **201** (2001).
- [30] G. Kuperberg, *Spiders for rank 2 Lie algebras*, Commun. Math. Phys. **180** (1996) 109–151, q-alg/9712003.
- [31] I. Kostov, *Free field presentation of the  $A_n$  coset model on the torus*, Nucl. Phys. **B300** (1988) 559–587.
- [32] A. Malkin, V. Ostrik, M. Vybornov, *Quiver varieties and Lusztig’s algebra*, math.RT/0403222.

- [33] D. Nikshych, L. Vainerman, *Finite quantum groupoids and their applications*, in New directions in Hopf algebras, 211-262, Math. Sci. Res. Inst. Publ. **43**, Cambridge Univ. Press, Cambridge, 2002, math.QA/0006057.
- [34] F. Nill, *Axioms for Weak Bialgebras*, Preprint Inst. Theor. Phys. FU, Berlin, math.QA/9805104.
- [35] A. Ocneanu, *Quantized groups, string algebras and Galois theory for algebras*, Operator algebras and Appl., Vol. 2, London Math. Soc. Lecture Notes Ser., **136**, Cambridge Univ. Press (1988) 119–172.
- [36] A. Ocneanu, (Lecture Notes written by Y. Kawahigashi), *Quantum Symmetry, Differential Geometry of Finite Graphs and Classification of Subfactors*, Univ. of Tokyo Seminar Notes (1990).
- [37] A. Ocneanu, *Quantum symmetries, operator algebras and invariant for manifolds*, Talk given at the First Caribbean Spring School of Mathematical and Theoretical Physics, Saint-François-Guadeloupe, 1993.
- [38] A. Ocneanu, *Operator Algebras, Topology and subgroups of quantum symmetries*, Tanaguchi conference on Mathematics, Nara, Japan, 1998. Notes by S Goto and N. Sato. Advances Studies in Pure Mathematics **31**, Math. Soc. of Japan (2001) 235–263.
- [39] A. Ocneanu, *Paths on Coxeter diagrams: from Platonic solids and singularities to minimal models and subfactors*, Notes taken by S. Goto, AMS Fields Institute Monographs **13** (1999), Rajarama Bhat et al eds.
- [40] A. Ocneanu, *The Classification of subgroups of quantum  $SU(N)$* , Lectures at Bariloche Summer School 2000, Argentina, AMS Contemp. Math. **294**, R. Coquereaux, A. García and R. Trinchero eds.
- [41] A. Ocneanu, *Higher Coxeter systems*, Talk given at MSRI, <http://www.msri.org/publications/ln/msri/2000/subfactors/ocneanu>.
- [42] V. Ostrik, *Module categories, weak Hopf algebras and modular invariants*, Transform. Groups **8** (2003) 177-206.
- [43] V. Pasquier, *Operator content of the ADE lattice models*, J. Phys. **A 20** (1987) 5707–5717; *Two-dimensional critical systems labelled by Dynkin diagrams*, Nucl.Phys. **B285** (1987) 162–172.
- [44] V.B. Petkova, J.-B. Zuber, *Generalised twisted partition functions*, Phys. Lett. **B504** (2001) 157–164, hep-th/0011021.

- [45] V.B. Petkova, J.B. Zuber, *The many faces of Ocneanu cells*, Nucl. Phys. **B603** (2001) 449–496, hep-th/0101151.
- [46] V.B. Petkova, J.-B. Zuber, *Verlinde Nim-reps for charge conjugate  $sl(N)$  WZW theory*, in A. Cappelli and G. Mussardo eds. Statistical Field Theories (Kluwer, 2002) 161–170; *Boundary conditions in charge conjugate  $sl(N)$  WZW theories*, hep-th/0201239.
- [47] G. Schieber, *L’algèbre des symétries quantiques d’Ocneanu et la classification des systèmes conformes à 2D*, PhD thesis (available in french and in portuguese, UP (Marseille) and UFRJ (Rio de Janeiro), Sept. 2003, math-ph/0411077.
- [48] G. Schieber, *Quantum symmetries of self-fusion orbifold graphs*, in preparation.
- [49] E. Verlinde, *Fusion rules and modular transformations in 2-D conformal field theory*, Nucl. Phys. **B300** (1988) 360–376.
- [50] J.-B. Zuber, *Generalized Dynkin diagrams and root systems and their folding*. YITP International Workshop on Recent Developments in QCD and Hadron Physics, Kyoto , Japan, 16-18 Dec 1996; *On Dubrovin Topological Field Theories*, Mod. Phys. Lett. **A9** (1994) 749–760.