

# Some comments on quantum magnetic monopoles

LUIZ C.L. BOTELHO

Departamento de Matemática Aplicada,  
Instituto de Matemática, Universidade Federal Fluminense,  
24220-140, Niterói, Rio de Janeiro, Brazil  
e-mail: botelho.luiz@ig.com.br

## Abstract

In this paper we intend to present some path-integral studies in the problem of confinement in the presence of fermionic and scalar magnetic monopole fields through:

1 - A Wilson Loop Path-Integral Evaluation associated to an effective second-quantized electromagnetic field generated by chiral abelian point-like monopole magnetic field current at its large mass London asymptotic limit.

2 - A Path-Integral Bosonization analysis of Quarks fields interacting with Kalb-Ramond fields considered as an effective Disorder Field Theory of a Q.C.D. vacuum of heavier monopoles.

3 - Improvements on the Wilson Loops evaluations in the well-known ADHM Antonov-Ebert model for Cooper pairs of point-like fermionic magnetic monopoles.

**Key-words:** Magnetic monopoles; Confinement; Path integral.

# 1 Introduction

The question of the existence of Magnetic Monopoles has been a fruitful research path on modern theoretical physics since the appearance of the seminal work of P.M. Dirac ([1]) on the subject ([1]). In the modern framework of Non-Abelian Gauge theories, most of the relevant dynamical questions about the physical modeling of particles interactions are transferred to the difficult and more subtle mathematical analysis of special gauge-field configurations (instantons, merons, strings, magnetic monopoles, etc...) which are expected to constitute the non-perturbative vacuum structure of the underlying Bosonic Yang-Mills Gauge theory. Among those special field configurations, the Magnetic Monopole has been considered as one of the basic hypothetical non-perturbative excitation expected to be connected to practically all non-trivial charge confining dynamical effects occurring on non-abelian Gauge theories. This fact is due to the hope that Magnetic Monopoles are the best candidates for explain naturally the (electrical) charge confinement ([2]). However magnetic monopoles by themselves should not be observed in the particle spectrum as a physical excitation. Note that this last constraint on monopole confinement makes the use of the standard Quantum Field techniques to handle magnetic monopoles dynamics a very difficult task ([3], [4]).

In this paper we address to these dynamical questions on Magnetic Monopole theory by path integrals analysis, specially the technique of four-dimensional chiral bosonization path-integral as earlier proposed by this author ([5]).

This paper is organized as follows

In Section 2, we show how to obtain by a direct evaluation, the area behavior for an abelian Wilson Loop phase Factor in the presence of an effective second quantized electromagnetic field generated by an (condensate) second quantized monopole fermion field, as much as envisaged as an dynamical mechanism in the famous Nambu-Mandelstam propose for the existence of a Meissner effect for magnetic monopoles vacuum condensation in Yang-Mills theory in order to explain the quark-gluon confinement. As a new result of our study, we claim, thus, to have produced a well-defined path integral procedure to prove the electric charge confining in the presence of a quantum dynamics of magnetic monopoles, with a Fermi-Dirac statistics.

In Section 3, we exactly analyze by path-integrals techniques the quantum field dynamics of (massless) fermions field interacting with Kalb-Ramond tensor fields, expected to represent dynamically quark fields interacting with rank-two tensor field, with the later field representing

the disorder field of a vacuum structure formed by condensation of magnetic monopoles ([3]). We show, thus, that it is ill-defined to associated physical observables LSZ interpolating fields for the fermion fields in the theory as consequence of the explicitly Bosonized structure formulae obtained for the matter excitations interacting with rank-two tensor fields through a spin orbit coupling with the Kalb Ramond field strenght, which by its turn provides another support for electrical charge confining in the presence of magnetic monopoles.

Finally in Section 4, we present some improvements on the Wilson Loops evaluation in the context of the Antonov-Ebert dual path integral associated to the dual Abelian Higgs Model of third reference in [3].

## 2 The abelian confinement in presence of magnetic monopoles, a Wilson Loop Gauge invariant path-integral evaluation

Let us start this section by considering the Euclidean path integral average associated to a  $U(1)$ -abelian field  $A_\mu(x)$  whose dual strength field intensity has a second quantized magnetic monopole as a (chiral) electromagnetic source

$$\begin{aligned} \langle W[C_{(R,T)}] \rangle &= \int D^F[A_\mu] D^F[\Omega] D^F[\bar{\Omega}] \delta^{(F)}[\partial_\mu^* F^{\mu\nu}(A) - (g\bar{\Omega}\gamma^\nu\gamma^5\Omega)] \\ &\times \exp\left(-\frac{1}{2} \int d^4x (\Omega, \bar{\Omega}) \begin{bmatrix} 0 & i \not{\partial} + M \\ (i \not{\partial} + M)^* & 0 \end{bmatrix} \begin{pmatrix} \Omega \\ \bar{\Omega} \end{pmatrix}\right) \\ &\times \exp\left(ie \oint_{C_{(R,T)}} A_\mu(x^\alpha) dX_\mu\right) \end{aligned} \quad (1-a)$$

Here  $(\Omega, \bar{\Omega})(x)$  are the Euclidean Fermion (second-quantized) point-like fundamental monopole fields with  $g$  denoting the magnetic charge which by its turn is supposed to be related to the  $U(1)$ -electric charge  $e$  by the Dirac quantization relation  $eg = \frac{n}{4}$  (with  $n \in \mathbb{Z}$ ).  $M$  denotes the magnetic monopole mass and  $W[C] = \exp\{ie \oint_{C_{(R,T)}} A_\mu dX_\mu\}$  is the  $U(1)$ -Wilson Loop phase factor defined by the (Euclidean) space-time trajectory of two static electric carrier external charges interacting with the fluctuating  $A_\mu(x)$  field generated by the (fluctuating) second quantized magnetic monopole fermionic source (see the constraint on eq.(1-a)). Note that

$C_{(R,T)}$  is the boundary of the square  $S_{(R,T)}$  below

$$C_{(R,T)} = \partial S_{(R,T)}; \quad S_{(R,T)} = \left\{ (x_0, x_1) \in R^2; -\frac{T}{2} \leq x_0 \leq +\frac{T}{2}; -\frac{R}{2} \leq x_1 \leq +\frac{R}{2} \right\} \subset R^4. \quad (1-b)$$

It is worth call the reader attention that the above written quantum Wilson Loop associated to static quarks charges can be physically replaced by the complete Generating functional of the second quantized Quark fields interacting with the Monopole Generated Electromagnetic field, namely

$$\begin{aligned} Z[\eta, \bar{\eta}] = & \int D^F[A_\mu] D^F[\Omega] D^F[\bar{\Omega}] [\delta(*\partial_\mu F^{\mu\nu}(A) - (g\bar{\Omega}\gamma^\nu\gamma^5\Omega)] \\ & \times \exp \left( -\frac{1}{2} \int d^4x(\Omega, \bar{\Omega}) \begin{bmatrix} 0 & i \not{\partial} + M \\ (i \not{\partial} + M)^* & 0 \end{bmatrix} \begin{pmatrix} \Omega \\ \bar{\Omega} \end{pmatrix} \right) \\ & \times \exp \left( -\frac{1}{2} \int d^4x(\psi, \bar{\psi}) \begin{bmatrix} 0 & i \not{\partial} + \not{A} \\ (i \not{\partial} + \not{A})^* & 0 \end{bmatrix} \begin{pmatrix} \psi \\ \bar{\psi} \end{pmatrix} \right) \\ & \times \exp \left( i \int d^4x(\psi, \bar{\psi}) \begin{pmatrix} \bar{\eta} \\ \eta \end{pmatrix} \right) \end{aligned} \quad (1-c)$$

For static charges eq.(1-c) reduces to eq.(1-a) as it is showed in first ref. [6].

In order to evaluate the path-integral eq.(1-a) from the physical point of view of an effective field theory ([5]), we should consider firstly the magnetic monopole field as a London large mass excitation in the fermonic path-integral weight of the Wilson Loop path integral average eq.(1-a). The reason why we should evaluate our Wilson Loop average in this context can be related to the fact that very heavy monopoles (but with small quantum fluctuations) are expected to populating the non-perturbative vacuum phase of any non-abelian Gauge Theory (at least in its confining phase) ([2], [3]). Let us, thus, re-write the magnetic monopole axial current constraint in eq.(1) by means of an axial-vectorial Lagrange multiplier field  $\lambda_\mu(x)$ , namely:

$$\begin{aligned} \langle W[C_{(R,T)}] \rangle = & \left\{ \int D^F[A_\mu] D^F[\Omega] D^F[\bar{\Omega}] D^F[\lambda_\mu] \right. \\ & \times \exp \left( i \int d^4x [\lambda_\nu (\partial_\mu^* F^{\mu\nu}(A) - g\bar{\Omega}\gamma^\nu\gamma^5\Omega)](x) \right) \\ & \times \exp \left[ -\frac{1}{2} \int d^4x(\Omega, \bar{\Omega}) \begin{bmatrix} 0 & i \not{\partial} + M \\ (i \not{\partial} + M)^* & 0 \end{bmatrix} \begin{pmatrix} \Omega \\ \bar{\Omega} \end{pmatrix} \right] \times \exp \left( ie \int_{C_{(R,T)}} A_\mu dX_\mu \right) \Big\} \end{aligned} \quad (2)$$

At this point we follow well known studies in the literature in order to give a correct meaning for the effective field theory associated to very heavier magnetic monopoles London large mass limit in the monopoles Fermionic determinants ([5]). It is a standard result in the subject that the (mathematical) leading limit of (renormalized) magnetic monopole large mass should be given by the auxiliary Gauge field mass term, (see refs. [5] for the calculational details an this London limit for Fermion determinants)

$$\begin{aligned} \lim_{M_{\text{ren}} \rightarrow \infty} |\det(i \not{\partial} + M_{\text{ren}} + g\gamma^5 \not{\lambda}_\mu)|^2 \\ \cong \exp \left\{ -\frac{1}{2} (\Lambda_{QCD} \cdot g^2) \int d^4x (\lambda_\mu(x))^2 \right\} + O(1/M_{\text{ren}}) \end{aligned} \quad (3)$$

Note that the appearance [through the phenomenological  $QCD$  vacuum scale  $\Lambda_{QCD} = (M_{\text{ren}})^{+2}$ ] of a mass term for the auxiliary vector field  $\lambda_\mu(x)$  which by its turn, should signals the expected dynamical breaking of the  $U(1)$ -axial gauge invariance (with opposite parity ([4], [5]) of this (non-physical) vectorial field by the phenomenon of dimensional transmutation on the adimensional  $g$ -coupling constant. This result indicates strongly the dynamical breaking of the  $U(1)$ -axial symmetry of the fermionic magnetic monopole second quantized field  $\{\Omega(x), \overline{\Omega}(x)\}$ .

After inserting eq.(3) into eq.(2) and by realizing the Gaussian  $\lambda_\mu$ -field path integral, we are led to consider the effective fourth-order Wilson Loop path integral average for eq.(1) as the leading London limit on the magnetic monopole mass  $M$ , namely:

$$\begin{aligned} \langle W[C_{(R,T)}] \rangle = & \left\{ \int D^F[A_\mu(x)] \delta^{(F)}(\partial_\mu A_\mu) \right. \\ & \times \exp \left( -\frac{1}{2(g^2 \Lambda_{QCD})} \int d^4x (A_\mu [(-\partial^2)^2] A_\mu)(x) \right) \\ & \left. \times \exp \left( ie \int_{C_{(R,T)}} A_\mu dX_\mu \right) \right\} + O(M^{-1}) \end{aligned} \quad (4)$$

The static inter-quark linear risen potential can be obtained from eq.(4) by using the dimensional regularization scheme of Bollini-Giambiagi for evaluating the Feynman-diagrams integrals as it is exposed in details on refs. ([6]). It yields the expected linear raising confining potential

$$\begin{aligned} V(R) &= (e^2 \cdot g^2) (\Lambda_{QCD}) R \\ &= \overline{A} \frac{n^2}{16} (\Lambda_{QCD}) \cdot R = \overline{A} \left( \frac{n^2}{2\pi\alpha'} \right) R = \alpha_{\text{eff}}(N^2) R \end{aligned} \quad (5)$$

Here  $\overline{A}$  is a model-calculational positive adimensional constant, which details will not be needed for our study, and  $\alpha'$  denotes the Regge Slope parameter associated to the non-perturbative vacuum scale  $\Lambda_{QCD} \sim (\frac{1}{2\pi\alpha'})$ . It is worth call the reader attention that we have obtained somewhat the infinite quantized number of parallel Regge trajectories from the Dirac topological quantization rule for electric and magnetic charges as it is suggested in the effective Regge slope parameter  $\alpha_{\text{eff}}(n^2) = n^2/2\pi\alpha'$ .

Thus we see that the effective path integral eq.(1) for the Wilson Loop in the presence of an electromagnetic field generated by a heavy quantum monopole leads naturally to a dynamics of Wilson Loop area behavior for the electrical charges in the theory, a result obtained by us explicitly through an exactly gauge invariant path-integral evaluation.

### 3 Monopoles interacting with Kalb-Ramond fields through spin-orbit coupling

In the last years, Kalb-Ramond field theory has been widely studied as an alternative dynamical quantum field scheme to the Higgs mechanism, as well as in relation to the dynamics of strings in the problem of string representation for Q.C.D. at large number of colors as a dynamical disorder field representing the effects of existence of magnetic monopoles ([2], [3]). The basic formalism used to analyze such Kalb-Ramond non-perturbative quantum dynamics has been the path-integral formalism, which has shown itself to be a very powerful procedure to understand correctly the different phases of the associated Kalb-Ramond Quantum Field Theory [7].

One important problem in those Path-integral studies, still missing in the literature, is that one related to the presence of interacting dynamical fermions (simulating second quantized matter fields) in the Kalb-Ramond Gauge theory. In this Section 3 we shall describe the extension of previous path-integral dualization-bosonization studies [8] to the case of Fermionic matter coupling through a spin-orbit field quantum interaction as it is expected to be relevant to describe the interacting physics of quarks and magnetic monopoles.

Let us start by considering the Abelian Kalb-Ramond first order action but now in the presence of massless dynamical fermions in the four-dimensional Euclidean world.

$$S[H, B, \psi, \overline{\psi}] = \int_{R^4} d^4x \left\{ \frac{1}{12} H_{\lambda\mu\nu} H^{\lambda\mu\nu} - \frac{1}{6} H^{\lambda\mu\nu} \partial_{[\lambda} B_{\mu\nu]} + \overline{\psi} (i \not{\partial} + ig \gamma^\alpha \gamma^\beta \gamma^\mu H_{\alpha\beta\mu}) \psi \right\}. \quad (6)$$

Here the dynamical fields are the independent three-form  $H$ , the KR gauge field  $B$  and the Dirac fermion fields  $(\psi, \bar{\psi})$ .

We shall apply the bosonization procedure in the path-integral framework through the following theory's generating functional (normalized to unity)

$$\begin{aligned} Z[J, \eta, \bar{\eta}] &= \int D^F[H] D^F[B] D^F[\psi] D^F[\bar{\psi}] \\ &\times \exp\{-S[H, B, \psi, \bar{\psi}]\} \\ &\times \exp\left\{-i \int_{R^4} d^4x (\bar{\eta}\psi + \bar{\psi}\eta + J_{\mu\nu}B^{\mu\nu})(x)\right\}. \end{aligned} \quad (7)$$

It is worth call the reader attention that the Path-integral eq.(7) is invariant under the KR gauge symmetry, provide the external source corrent  $J_{\mu\nu}$  is chosen to be divergence free and our proposed action term related to the direct interaction of the quantum fermionic matter with the Kalb-Ramond gauge field through its strenght three-form  $H$  – the spin orbit fermion interaction. (see eq.(6)).

The Path-Integral Bosonization analysis proceeds as usually by integrating exactly out the Kalb-Ramond gauge potential field which produces as a result the delta functional [8].

$$\begin{aligned} Z[J, \eta, \bar{\eta}] &= \int D^F[H] D^F[\psi] D^F[\bar{\psi}] \delta^{(F)}(\partial_\lambda H^{\lambda\mu\nu} - J^{\mu\nu}) \\ &\times \exp\left\{-\int_{R^4} d^4x \left[\frac{1}{12} H_{\lambda\mu\nu} H^{\lambda\mu\nu} + \bar{\psi}(i \not{\partial} + ig\gamma^\alpha \gamma^\beta \gamma^\mu H_{\alpha\beta\mu})\psi\right](x)\right\}. \end{aligned} \quad (8)$$

Let us note that the delta functional integrand inside of the path integral eq.(8) imposes the classical equations of motion on the three-form Kalb-Ramond strenght  $H$  which by its turn can be exactly solved by the Rham-Hodge theorem in terms of the effective dual scalar axion (zero-form) dynamical degree of freedom in the KR theory defined in a space-time topologically trivial as considered in our path integral eq.(8)

$$H_{\lambda\mu\nu} = g\varepsilon^{\lambda\mu\nu\rho} \partial_\rho \vartheta + \partial^{[\lambda} \frac{1}{\partial^2} J^{\mu\nu]}. \quad (9)$$

At this point we re-write the effective action eq.(8) in a four-dimensional bosonized chiral

action [9]

$$\begin{aligned}
Z[J, \eta, \bar{\eta}] = & \int D^F[\vartheta] \\
& \times \exp \left\{ -\frac{1}{2} \int_{R^4} d^4x \left[ g^2 \partial_\mu \vartheta \partial^\mu \vartheta + \frac{1}{2} J^{\mu\nu} \left( -\frac{1}{\partial^2} \right) J_{\mu\nu} \right] (x) \right\} \\
& \times \int D^F[\psi] D^F[\bar{\psi}] \exp \left\{ -\frac{1}{2} \int_{R^4} d^4x (\bar{\psi} e^{ig\gamma_5 \vartheta} \not{\partial} e^{ig\gamma_5 \vartheta} \psi)(x) \right\} \\
& \times \exp \left\{ -\frac{1}{2} \int_{R^4} d^4x \left( ig \bar{\psi} \left[ \gamma^\alpha \gamma^\beta \gamma^\rho \partial^{[\alpha} \frac{1}{\partial^2} J^{\beta\rho]} \right] \right) \psi \right\} (x) \\
& \times \exp \left\{ -i \int_{R^4} d^4x (\psi \bar{\eta} + \bar{\psi} \eta)(x) \right\}. \tag{10}
\end{aligned}$$

After considering the chiral-fermion field variable change on the fermionic path-integral term of eq.(10)

$$\bar{\psi} = \bar{\chi} e^{-ig\gamma_5 \vartheta} \tag{11-a}$$

$$\psi = e^{-ig\gamma_5 \vartheta} \chi \tag{11-b}$$

$$\begin{aligned}
D[\psi] D[\bar{\psi}] &= D[\chi] D[\bar{\chi}] \frac{\det[e^{ig\gamma_5 \vartheta} \not{\partial} e^{ig\gamma_5 \vartheta}]}{\det[\not{\partial}]} \\
&= D[\chi] D[\bar{\chi}] J[\vartheta], \tag{11-c}
\end{aligned}$$

we obtain the exactly bosonized path-integral representation for the KR first order theory as given by eq.(7), namely:

$$\begin{aligned}
Z[J, \eta, \bar{\eta}] = & \int D^F[\vartheta] D[\chi] D[\bar{\chi}] J[\vartheta] \\
& \times \exp \left\{ -\int_{R^4} d^4x \left[ \frac{g^2}{2} \partial_\mu \vartheta \partial^\mu \vartheta - \frac{1}{2} J^{\mu\nu} (\partial^2)^{-1} J_{\mu\nu} \right] (x) \right\} \\
& \times \exp \left\{ -\frac{1}{2} \int_{R^4} d^4x (\bar{\chi} \not{\partial} \chi)(x) \right\} \\
& \times \exp \left\{ -\frac{1}{2} ig \int_{R^4} d^4x \left( \bar{\chi} \left( \gamma^\alpha \gamma^\mu \gamma^\nu \partial^{[\alpha} \frac{1}{\partial^2} J^{\mu\nu]} \right) \chi \right) (x) \right\} \\
& \times \exp \left\{ -i \int_{R^4} d^4x (\bar{\chi} e^{-ig\gamma_5 \vartheta} \eta + \bar{\eta} e^{-ig\gamma_5 \vartheta} \chi)(x) \right\}, \tag{12}
\end{aligned}$$

here the functional Fermion Jacobian eq.(11-c) has been exactly evaluated in refs. [9]:

$$\begin{aligned}
J_\varepsilon[\vartheta] = & \exp \left\{ \frac{g^2}{4\pi^2 \varepsilon} \int_{R^4} d^4x (\partial_\mu \vartheta)^2(x) \right\} \\
& \times \exp \left\{ -\frac{g^2}{4\pi^2} \int_{R^4} d^4x (\partial^2 \vartheta)(\partial^2 \vartheta)(x) \right\} \\
& \times \exp \left\{ \frac{g^4}{12\pi^2} \int_{R^4} d^4x [\vartheta (\partial_\mu \vartheta)^2 (-\partial^2 \vartheta)](x) \right\}. \tag{13}
\end{aligned}$$



As a first remark to be made on the above written result we note that its first term has the effect of formally inducing a renormalisation of the  $g$ -charge after the cutt-off removing  $\varepsilon \rightarrow 0$  on the complete result eq.(7), namely

$$g_{\text{bare}}^2(\varepsilon) \left( 1 + \frac{1}{4\pi^2\varepsilon} \right) = g_{\text{ren}}^2. \quad (9)$$

By secondly, we point out the appearance of the fourth-order kinetic term for the scalar effective KR field  $\vartheta(x)$ , a very important result for the model ultra-violet finiteness.

An another important physical result coming from the set eq.(12)–eq.(14) is the explicitly fermionic matter asymptotic freedom as can be see directly from the factorized – decoupled form of the full interacting matter fermionic propagator, namely

$$\frac{1}{(i)^3} \frac{\delta Z[\eta, \bar{\eta}, J]}{\delta \eta_\alpha(x) \delta \eta_\beta(y)} \Big|_{\substack{J=0 \\ \eta=\bar{\eta}=0}} = S_{\alpha\beta}(x-y) \times F(x,y) \quad (15)$$

with  $S_{\alpha\beta}(x-y)$  denoting the free fermion propagator and the (decoupled) Kalb-Ramond form factor being given exactly by the (perturbative finite) fourth-order  $\vartheta$ -path integral as remarked above.

$$\begin{aligned} F(x, y) = & \int D^F[\vartheta] e^{-\frac{1}{2} g_{\text{ren}}^2 \int_{R^4} (\partial_\mu \vartheta)^2(x) d^4x} \\ & \times e^{-\frac{g_{\text{ren}}^2}{4\pi^2} \int_{R^4} (\partial_\mu^2 \vartheta)^2(x) d^4x} \\ & \times e^{+\frac{g_{\text{ren}}^2}{4\pi^2} \int_{R^4} [\vartheta (\partial_\mu \vartheta)^2 (-\partial_\mu^2 \vartheta)](x) d^4x} \\ & \times \{ (\exp -ig_{\text{ren}} \gamma_5 \vartheta(x)) (\exp -ie_{\text{ren}} \gamma_5 \vartheta(y)) \} \end{aligned} \quad (16)$$

which goes to 1 in the high energy limit of  $|x-y| \rightarrow 0$  as a result of the path-integral super renormalizability associated to the effective axion scalar dual Kalb-Ramond theory eq.(6)) [the well-known phenomenon of asymptotic freedom in confining gauge theories]. A low energy study of the form-factor eq.(16) has been carried out in refs. [9] (Appendix). There, we have suggested that these bosonized fermionic fields do not possesses LSZ interpolating fields, since the associated two-point Euclidean correlation function eq.(15) defines Wightman functions which are ultra-distributions in Jaffe Distributional Spaces and not in the usual Schwartz Tempered Distributional Spaces naturally associated to the existence of LSZ interpolating fields (a well defined Scattering Matrix) in the quantum field theory eq.(7).

A calculational remark to be made at this point of our paper is related to the straightforward exactly solubility for the Macroscopic radiative corrections evaluations of the Kalb-Ramond

gauge potential propagator

$$\begin{aligned} \frac{1}{i^2} \frac{\delta^2[J, \eta, \bar{\eta}]}{\delta J_{\mu\nu}(x) \delta J_{\alpha\beta}(y)} \Big|_{\substack{\eta=\bar{\eta}=0 \\ J=0}} &= \langle B_{\mu\nu}(x) B_{\alpha\beta}(y) \rangle \\ &= (-\partial^2)^{-1}(x, y) + e_{\text{ren}}^2 \int d^4z d^4z' (-\partial^2)^{-1}(z-x) (-\partial^2)^{-1}(z-y) \\ &\quad \times \partial_z^{[\lambda} \partial_{z'}^{\lambda']} \langle (\bar{\chi}(z) (\gamma^\lambda \gamma^{[\mu} \gamma^{\nu]}) \chi(z)) (\bar{\chi}(z') (\gamma^{\lambda'} \gamma^{[\alpha} \gamma^{\beta]}) \chi(z')) \rangle^{(0)}, \end{aligned} \quad (17)$$

here  $\langle \rangle^{(0)}$  denotes the free fermion average path integral

$$\langle \rangle^{(0)} = \int D(\chi) D[\bar{\chi}] e^{-\frac{1}{2} \int_{R^4} d^4x (\bar{\chi} \not{\partial} \chi)(x)}. \quad (18)$$

The exactly evaluation of the quantum correction eq.(17) is standard and can be easily obtained by just using the well-known Dirac matrixes relationship

$$\gamma^\lambda \gamma^\mu \gamma^\nu = (S_{\lambda\mu\nu\sigma} + \varepsilon_{\lambda\mu\nu\sigma} \gamma_5) \gamma^\sigma \quad (19)$$

$$S_{\lambda\mu\nu\sigma} = (\delta_{\lambda\mu} \delta_{\nu\sigma} + \delta_{\mu\nu} \delta_{\lambda\sigma} - \delta_{\lambda\nu} \delta_{\mu\sigma}). \quad (20)$$

The above exposed results concludes our Section 3 these path-integral method studies on the four-dimensional exactly path-integral Bosonization of our abelian interacting KR field.

## 4 Some comments on the path-integral Wilson Loop evaluation in the Dual Abelian Higgs Model of Antonov-Ebert

In this last section of our study, we intend to present some calculational improvements on the Wilson Loop path integral evaluation in the context of the usual Abelian Higgs Model through the framework of path integral duality transformation as exposed in details on the Antonov & Ebert paper in ref. [3].

Let us briefly describe the path integral duality of the extended Dual Abelian Higgs Model of Antonov-Ebert.

As a first step in such analysis, one starts from the following phenomenological expression for the partition functional of the model

$$\begin{aligned} Z(\lambda) = \int |\Phi| D^F |\Phi| D^F B_\mu D^F \theta \exp \left\{ - \int_{R^4} \left[ \frac{1}{4} (F_{\mu\nu} - F_{\mu\nu}^E)^2 + \frac{1}{2} |D_\mu \Phi|^2 \right. \right. \\ \left. \left. + \lambda (|\Phi|^2 - \eta)^2 \right] \right\} \end{aligned} \quad (21-a)$$

where  $\Phi(x) = |\Phi(x)| \exp |\theta(x)|$  is an disorder scalar Higgs field of the Magnetic Monopoles “Cooper pairs”  $(\Omega\overline{\Omega})(x)$  (see eq.(1-a)) and  $F_{\mu\nu}^E(x)$  is the dual electromagnetic field generated by the external static “quarks” source (the loop  $X_\mu(\sigma)$  on eq.(1-a)).

It is thus argued in details on the above cited paper of Antonov & Ebert that the partition functional eq.(21) has the following stringy representation in the London phenomenological  $\lambda \rightarrow +\infty$  limit (the equivalent of our London large mass  $M$  limit - eq.(2)-eq.(3). Namely

$$Z^{\text{eff}}(\infty) \sim \int_{\partial X^\mu(\xi)=C^\mu} D^F X_\mu(\xi) \exp \left\{ -\pi^2 \int_\Sigma d\sigma_{\lambda\nu}(x) \int_\Sigma d\sigma_{\mu\rho}(y) D^{\lambda\nu,\mu\rho}(|x-y|) \right\} \quad (21-b)$$

Here  $X^\mu(\xi)$  parametrizes the string world-sheet  $\Sigma$  possessing as boundary the quark source loop  $C^\mu \equiv X_\mu(\sigma)$  and the Antonov-Ebert propagator of the Kalb-Ramond field duality in this London limit is exactly given in momentum space by

$$D^{\lambda\nu,\mu\rho}(|x-y|) = D_{(1)}^{\lambda\nu,\mu\rho}(|x-y|) + D_{(2)}^{\lambda\nu,\mu\rho}(|x-y|) \quad (21-c)$$

where

$$\begin{aligned} D_{(1)}^{\lambda\nu,\mu\rho}(|x|) &= (\delta_{\lambda\mu}\delta_{\nu\rho} - \delta_{\mu\nu}\delta_{\lambda\rho}) \frac{C_1}{|x|} \\ D_{(2)}^{\lambda\nu,\mu\rho}(|x|) &= \frac{C_2}{|x|^2} \left\{ \left[ \frac{K_1(m|x|)}{|x|} + \frac{m}{2}(K_0 + K_1)(m|x|) \right] \right. \\ &\quad \times (\delta_{\lambda\mu}\delta_{\nu\rho} - \delta_{\mu\nu}\delta_{\lambda\rho}) \\ &\quad + \frac{1}{2|x|} \left[ 3\left(\frac{m^2}{4} + \frac{1}{|x|^2}\right) K_1(m|x|) + \frac{3m}{2|x|}(K_0 + K_2)(m|x|) + \frac{m^2}{4} K_3(m|x|) \right] \\ &\quad \left. \times (\delta_{\lambda\rho}x_\mu x_\nu + \delta_{\mu\nu}x_\lambda x_\rho - \delta_{\mu\lambda}x_\nu x_\rho - \delta_{\nu\rho}x_\mu x_\lambda) \right\} \end{aligned} \quad (21-e)$$

with  $K_i$  denoting the relevant usual Modified Bessel functions,  $m$  is the mass of the dual gauge bosons generated by the Higgs mechanism and  $C_1$  e  $C_2$  are model calculational constants.

At this point, we argue that all the above pointed out duality derivation holds true only for small string world-sheet deviations from the minimal surface  $C^\mu = \partial X^\mu(\xi)$  since we have Frozen the radial part of the monopole disorder field to its fixed v.e.v  $\eta$ . A very important and direct consequence of this remark of ours is that one can safely substitute the somewhat formal Feynman path measure on the string vector position effective partition functional eq.(21-b) by the so-called extrinsic space-times vorticity tensor current defined as

$$\begin{aligned} \Sigma_{\mu\nu}(x) &\equiv \int_\Sigma d\sigma_{\mu\nu}(x) \delta(x - x(\xi)) \\ &\equiv \int_\Sigma d^2\xi J^{\mu\nu}(\xi) \delta(x - X(\xi)) \sqrt{g(X(\xi))}, \quad \text{where } J^{\mu\nu}(\xi) = \varepsilon^{ab}(\partial_a X^\mu \partial_b X^\nu)(\xi) \end{aligned} \quad (21-f)$$

is the (non-normalized to unity) string world-sheet extrinsic orientation area tensor.

The argument for the validity of such path-integral dynamical degree replacement is the following.

For small deviations from the minimal area string world-sheet  $X_{cl}^\mu(\xi)$ , we have the usual functional metric decomposition

$$(X^\mu(\xi) = X_{cl}^\mu(\xi) + \varepsilon Y^\mu(\xi) + O(\varepsilon))$$

$$\int_{\Sigma} d^2\xi \sqrt{g(x_{cl}^\mu)} (\delta J^{\mu\nu}(\delta J^{\mu\nu}))(\xi) \sim \int_{\Sigma} d^2\xi \sqrt{g(X_{cl}^\mu)} \left\{ [\varepsilon^{ab}(\partial_a X_{cl}^\mu)(\partial_b \delta Y^\nu) + \varepsilon^{ab}(\partial_a \delta Y^\mu)(\partial_b X_{cl}^\mu)] \right. \\ \left. [\varepsilon^{ab}(\partial_a X_{cl}^\mu)(\partial_b \delta Y^\nu) + \varepsilon^{ab}(\partial_a \delta Y^\mu)(\partial_b X_{cl}^\mu)] \right\} + O(\varepsilon^4) \quad (22)$$

which straightforwardly leads to the following volume element functional change

$$D^F[J^{\mu\nu}(\xi)] \equiv \prod_{(\xi)} (\delta J^{\mu\nu}(\xi)) = \left( \prod_{(\xi)} \delta x^\mu(\xi) \right) \cdot \det_{(\xi, \mu, \nu)}^{\frac{1}{2}} [\mathcal{L}^{\mu\nu}(x_{cl}^\mu)] \quad (23)$$

where  $\mathcal{L}^{\mu\nu}(X_{cl}^\mu)$  is a second-order elliptic operator depending only on the classical minimal-area string configuration  $X_{cl}^\mu(\xi)$ , so cancelling itself when one is realising path-integral averages with the partition functional eq.(21-b).

Now it is straightforward to see that one can replace on basis of the above expected small string world-sheet deviations the average over the string vector position by the string vorticity degree of freedom, namely

$$\int_{R^4} (\delta \Sigma^{\mu\nu}(x) \cdot \delta \Sigma_{\mu\nu}(x)) d^4x \sim \overline{C} \int_{\Sigma} (\delta J^{\mu\nu}(\xi) \cdot \delta J_{\mu\nu}(\xi)) d^2\xi \quad (24)$$

with  $\overline{C}$  denoting an over-all (cut-off dependent) constant which cancels with itself when evaluating path-integral averages ([6]).

The important consequence of the above analyzed variable change of the string vector position variable by the string extrinsic vorticity field is affording the exactly path-integral solubility of the generating functional of the strenght of the usual gauge field  $A_\mu$  in the Abelian Higgs Model with the following result

$$Z[S_{\alpha\beta}] = \exp \left( - \int d^4x S_{\mu\nu}^2 \right) \\ \times \int D^F[\Sigma^{\gamma\zeta}(x)] \times \exp \left( -4\pi i e \int d^4x (S_{\mu\nu} \Sigma^{\mu\nu})(x) \right) \\ \times \exp \left\{ - \int d^4x d^4y \left( \pi \Sigma_{\lambda\nu}(x) - \frac{i}{e} S_{\lambda\nu}(x) \right) D^{\lambda\nu, \mu\rho}(|x-y|) \right. \\ \left. \times \left( \pi \Sigma_{\mu\rho}(y) - \frac{i}{e} S_{\mu\rho}(y) \right) \right\} \quad (25)$$

For instance:

$$\begin{aligned}
& \left\langle \left( \frac{1}{2} \varepsilon^{\lambda\nu\alpha\beta} F_{\alpha\beta} \right) (x) \left( \frac{1}{2} \varepsilon^{\mu\rho\alpha'\beta'} F_{\alpha'\beta'} \right) (x) \right\rangle \\
& \equiv \frac{1}{Z(0)} \frac{\delta^2 Z[S_{\gamma\lambda}]}{\delta S_{\lambda\nu}(x) \delta S_{\mu\rho}(y)} \Big|_{S_{\gamma\lambda} \equiv 0} \\
& = (\delta_{\lambda\mu} \delta_{\nu\rho} - \delta_{\lambda\rho} \delta_{\mu\nu}) \delta^{(4)}(x-y) + \frac{2}{e^2} D^{\lambda\nu, \mu\rho}(|x-y|) \\
& \quad - 4\pi^2 \left\langle \left[ 2e \Sigma_{\lambda\nu}(x) - \left( \frac{1}{e} \int d^4 z \Sigma_{\alpha\beta}(x) D^{\alpha\beta, \lambda\nu}(z-x) \right) \right] \right. \\
& \quad \times \left. \left[ 2e \Sigma_{\mu\rho}(y) - \frac{1}{e} \int d^4 u \Sigma_{\gamma\zeta}(u) D^{\gamma\zeta, \mu\rho}(|u-y|) \right] \right\rangle_{\Sigma} \quad (26)
\end{aligned}$$

where the normalized Gaussian path-integral average  $\langle \cdot \rangle_{\Sigma}$  is defined explicitly by

$$\begin{aligned}
\langle \cdot \rangle_{\Sigma} &= \frac{1}{2} \int D^F[\Sigma_{\alpha\beta}(x)] (\dots) \exp \left[ -\pi^2 \int_{R^4} dx dy \Sigma_{\alpha\beta}(x) \right. \\
& \quad \times \left. D^{\alpha\beta, \gamma\zeta}(|x-y|) \Sigma_{\gamma\zeta}(y) \right] \quad (27-a)
\end{aligned}$$

with

$$Z = \int D^F[\Sigma_{\alpha\beta}(x)] \exp \left[ -\pi^2 \int_{R^4} dx dy \Sigma_{\alpha\beta}(x) D^{\alpha\beta, \gamma\zeta}(|x-y|) \Sigma_{\gamma\zeta}(y) \right]. \quad (27-b)$$

After our remarks as expressed by eq.(26)–eq.(27) on the Antonov-Ebert paper ([3]), we now pass on to the Wilson Loop evaluation of eq.(1-a) on the dual “stringy effective” path integral eq.(26)–eqs.(27).

The main point for evaluation of eq.(1-a) in terms of an effective string theory is to re-write it in terms of the strenght field by means of the Stokes Theorem followed obviously by the string path integral average eqs.(27).

We have thus the following string functional integral representation for the Wilson Loop

$$\langle W[C_{(R,T)}] \rangle = \left\langle \exp \left\{ ie \int d^4 x (F_{\alpha\beta}(A) \cdot J_{\alpha\beta}(C_{R,T})) \right\} \right\rangle_{\Sigma} \quad (28)$$

Here the boundary’s rectangle Loop current  $J_{\alpha\beta}(C_{R,T})(x^\mu) = \int_{C_{(R,T)}} \delta(x^\mu - X^\mu(s)) (X_\alpha dX_\beta)(s)$ , with  $X_\mu(s)$  for  $0 \leq s \leq 1$  denoting a parametrization of the rectangle’s boundary  $C_{(R,T)}$ . At this point of our study, we propose to use a cumulant expansion for evaluating eq.(28) in the “stringy” DAHM model, which in generic form reads

$$\begin{aligned}
& \langle W[C_{R,T}] \rangle \\
& = \langle \exp \{ ie \int F \cdot J \} \rangle \\
& = \exp \left\{ \langle ie \int F \cdot J \rangle_{\Sigma} + \frac{1}{2} \left[ \langle (ie \int F \cdot J)^2 \rangle_{\Sigma} - \left\langle \left( ie \int F \cdot J \right)^2 \right\rangle_{\Sigma} \right] + \dots \right\}. \quad (29)
\end{aligned}$$

Extensive calculations of eq.(29), including spin degrees of freedom (see second reference on ref. [6] and refs. [10]) will be reported elsewhere.

## References

- [1] P.A.M. Dirac, Phys. Rev. 74, 817 (1948)
- [2] S. Mandelstam, Phys. Lett. B 53, 476 (1975)
  - K.I. Kondo, Phys. Rev. D57, 7467, (1998)
  - W. Ellwanger, Nucl. Phys. B 531, 593 (1998)
  - Z. Ezawa, A. Iwazaki, Phys. Rev. D25, 2681 (1982)
  - Y. Nambu, Phys. Rev. D10, 4246 (1974)
- [3] A.M. Polyakov, Part. Phys. B486, 23, (1997).
  - L.C.L. Botelho, Mod. Phys. Lett 20A, 12, (2005).
  - D. Antonov, D. Ebert, Eur. Phys. J. C8, 343–351 (1999).
- [4] Chan Hung-Mo, Tsou sheung, Tsun and, Phys. Rev. D56, 3646 (1997)
- [5] Luiz C.L. Botelho, Int. J. Mod. Phys. A 15 (5): 755-770 (2000)
  - M. Faber et al, Eur. Phys. J. C7, 685–695 (1999)
- [6] Luiz C.L. Botelho, Phys. Rev. D70, 045010 (2004)
  - Luiz C.L. Botelho, Eur. Phys. J. C44, 267–276 (2005).
- [7] Orland P. Nucl. Phys. B205, 107 (1982).
  - Aurilia A., Takahashi, Y., Prog. Theor. Phys. 66, 69 (1981).
  - Savit R., Rev. Mod. Phys. 52, 453 (1980).
  - Luiz C.L. Botelho, J. Math. Phys. 30 (9): 2160 (1989).
- [8] Smailagic A. and Spallucci, E., Phys. Rev. D61, 067701 (2000).

- [9] Luiz C.L. Botelho, Phys. Rev. D**39**(10): 3051–3054 (1989).  
Damgard P.H., Nielsen, H.B., Sollacher, P. Nucl. Phys. B**385**, 227–250 (1992).
- [10] A.I. Karamikas, C.N. Ktorides, N.G. Stefanis - Eur. Phys. J. C**26**, 445–455 (2003).