

PARAMETRIZATION OF SINGULARITIES OF THE DEMIAŃSKI-NEWMAN SPACETIMES

J. Gariel^{1*}, G. Marilhacy¹, N. O. Santos^{2†}
and R. Colistete, Jr.^{1‡}

¹Université Paris VI, CNRS/UPRESA 7065

Gravitation et Cosmologie Relativistes

Tour 22-12, 4ème étage, Boîte 142, 4 place Jussieu

75005 Paris, France.

²Centro Regional Sul de Pesquisas Espaciais - INPE/MCT,

Cidade Universitária, CEP. 97105-900, Santa Maria, RS, Brazil

and

LAFEX, Centro Brasileiro de pesquisas Físicas - CBPF,

Rua Dr. Xavier Sigaud 150, Urca, CEP: 22290-180, Rio de Janeiro – RJ, Brazil

November 28, 2000

Abstract

We propose a new presentation of the Demiański-Newman solution by introducing a new dimensionless parameter s . We ascribe to s the role of parametrizing the singular rings on the inner ergosphere.

Key-words: Theory of Gravitation; Axially symmetric systems; Singularities.

Pacs numbers : 04.70.Bw, 04.20.Dw, 04.20.Jb

*Email address: gariel@ccr.jussieu.fr

†E-mail: nos@lacesm.ufsm.br and nos@lafex.cbpf.br

‡Email address: coliste@ccr.jussieu.fr

1 Introduction

The Kerr metric can be easily obtained from the Ernst equation [1],

$$(\xi\xi^* - 1)\nabla^2\xi = 2\xi^*\nabla\xi \cdot \nabla\xi, \quad (1)$$

where ∇ denotes the usual three dimensional spatial operator and ξ is a complex potential being function of λ and μ which are the prolate spheroidal coordinates. By considering, as solution of (1),

$$\xi = p_K\lambda + iq_K\mu, \quad (2)$$

where p_K and q_K are real constants satisfying

$$p_K^2 + q_K^2 = 1, \quad (3)$$

we can obtain the Kerr solution.

The theorem of Robinson-Carter ([2] see p. 292) demonstrates the uniqueness of this vacuum stationary axisymmetric solution with an asymptotically flat behaviour and smooth convex event horizon without naked singularity. This solution is characterized by only two independent parameters, being the mass M and the angular momentum $J = aM$, where a is angular momentum per unit mass. The Kerr solution has an event horizon (or outer horizon), a Cauchy horizon (or inner horizon), an outer ergosphere (or stationary limit surface), an inner ergosphere and a ring singularity.

In this paper we propose to generalize the solution (2) by considering

$$\xi = p\lambda + \beta\mu + i(\gamma\lambda + q\mu), \quad (4)$$

where p, β, γ and q are real constants. The expression (4) is also solution of the Ernst equation and corresponds to the Demiański-Newman (DN) solution [3] as we shall see in section 2. The DN solution obtained through a complex transformation from the Kerr solution, namely the introduction of a constant phase factor (see (7) hereafter), has three independent parameters. The interpretation of the third parameter, usually denoted l , compared to just two parameters in the Kerr solution, is still not clear. In its place we introduce another parameter, which we call s , which parametrizes the singularities in a particularly simple way, as we shall see in section 4. We write the corresponding metric in section 3.

2 Choice of the parameters

The expression (4) is a solution of the equation (1) if the following conditions are satisfied

$$p^2 + \beta^2 + \gamma^2 + q^2 = 1, \quad (5)$$

$$p\beta = -\gamma q. \quad (6)$$

Let us recall that the DN solution [3] is usually presented [4] as

$$\xi = e^{ia_1}(p_{DN}\lambda + iq_{DN}\mu), \quad (7)$$

with

$$p_{DN}^2 + q_{DN}^2 = 1, \quad (8)$$

where p_{DN}, q_{DN} and a_1 are real constants. The comparison between (4) and (7) imposes

$$p = p_{DN} \cos a_1, \quad q = q_{DN} \cos a_1, \quad (9)$$

$$\beta = -q_{DN} \sin a_1, \quad \gamma = p_{DN} \sin a_1. \quad (10)$$

Then (5)–(6) are automatically satisfied, i.e., there is an identity between (4) and (7). In order to write the DN metric in Boyer-Lindquist (BL) coordinates the following relations are usually considered ([4] see p. 387),

$$\lambda = \frac{r - M}{k_{DN}}, \quad \mu = \cos \theta, \quad (11)$$

$$p_{DN} = \frac{k_{DN}}{\sqrt{M^2 + l^2}}, \quad q_{DN} = \frac{a}{\sqrt{M^2 + l^2}}, \quad (12)$$

$$\cos a_1 = \frac{M}{\sqrt{M^2 + l^2}}, \quad \sin a_1 = \frac{l}{\sqrt{M^2 + l^2}}, \quad (13)$$

with

$$k_{DN}^2 = M^2 + l^2 - a^2, \quad (14)$$

where l is a third parameter with mass dimension and r and θ are spherical coordinates. When $l = 0$, the DN metric becomes the Kerr metric.

Here, we shall not proceed in this way and the relations (7)–(14) will not be used. Instead, we shall present the DN solution as follows.

In place of (7), we consider the following complex transformation carried out on the Kerr solution,

$$\xi = e^{ia_1} (p_K \lambda + i q_K \mu), \quad (15)$$

where the variables λ and μ are the Kerr's ones.

This interpretation of the DN solution, as the Kerr solution with a phase factor, seems simpler and more natural to us. Indeed (4) is, for us, a simple linear extension of the Kerr solution of the Ernst equation.

Then, by identification between (15) and (4) we obtain relations identical to (9)–(10), where p_{DN} and q_{DN} are simply replaced by p_K and q_K :

$$p = p_K \cos a_1, \quad q = q_K \cos a_1, \quad (16)$$

$$\beta = -q_K \sin a_1, \quad \gamma = p_K \sin a_1. \quad (17)$$

These two Kerr parameters

$$p_K = \frac{k}{M}, \quad q_K = \frac{a}{M}, \quad (18)$$

with

$$k^2 = M^2 - a^2, \quad (19)$$

are linked to the two physical parameters M and a . Note that (p_K, q_K) are different from (p_{DN}, q_{DN}) given in (12), which are linked to the three parameters M , a and l , though each couple obeys the same relation (8) or (3).

To the third parameter a_1 , present in the DN solution (15), we associate the dimensionless parameter s defined by

$$s^2 = \frac{1}{\cos^2 a_1}, \quad (s \geq 1). \quad (20)$$

So, the relations (16)–(17) become

$$p = \frac{p_K}{s} = \frac{k}{Ms}, \quad q = \frac{q_K}{s} = \frac{a}{Ms}, \quad (21)$$

$$\frac{\gamma}{p} = -\frac{\beta}{q} = \sqrt{s^2 - 1}. \quad (22)$$

Note that relation (22) also holds from (9)–(10), i.e., in the usual interpretation of the DN solution. With (22), relation (5) becomes

$$s^2(p^2 + q^2) = 1, \quad (23)$$

which is also in agreement with (21).

Finally, instead of (11) with (14), we introduce BL coordinates,

$$\lambda = \frac{r - M}{k}, \quad \mu = \cos \theta. \quad (24)$$

where k is defined in (19), which is the BL transformation used for the Kerr solution. In particular, the coordinates are the Kerr's ones, coherently with (15), which is not the case in the usual DN solution for which λ depends on l (compare (11) and (24)).

3 Metric coefficients

The stationary axisymmetric metric, being in the Papapetrou form [4, 5], reads

$$ds^2 = f(dt - \omega d\phi)^2 - \frac{k^2}{f} \left[e^{2\gamma}(\lambda^2 - \mu^2) \left(\frac{d\lambda^2}{\lambda^2 - 1} + \frac{d\mu^2}{1 - \mu^2} \right) + (\lambda^2 - 1)(1 - \mu^2)d\phi^2 \right], \quad (25)$$

where f, ω and γ are functions of λ and μ only. The real part, P , and the imaginary part, Q , of ξ given by (4) become, using (22),

$$P = p\lambda - q\sqrt{s^2 - 1}\mu, \quad Q = p\sqrt{s^2 - 1}\lambda + q\mu. \quad (26)$$

The Ernst method [1, 4] to determine the metric consists to make a homographic transformation

$$\zeta = \frac{\xi - 1}{\xi + 1}, \quad (27)$$

where ζ is of the form

$$\zeta = f + i\Omega, \quad (28)$$

with f being the metric coefficient of (25) and Ω being the so called twist potential linked to the dragging of spacetime, ω , through the differential relations

$$\omega'_\lambda = \frac{k(\mu^2 - 1)}{f^2} \Omega'_\mu, \quad \omega'_\mu = \frac{k(\lambda^2 - 1)}{f^2} \Omega'_\lambda, \quad (29)$$

where indexes indicate to which variable the differentiation, indicated by primes, is to be taken. Once (29) is integrated we obtain ω . Applying this method we have

$$f = \frac{s^2[q^2(\mu^2 - 1) + p^2(\lambda^2 - 1)]}{s^2(p^2\lambda^2 + q^2\mu^2) + 2[p\lambda - \sqrt{s^2 - 1}q\mu] + 1}, \quad (30)$$

$$\omega = \frac{2k}{s^2p} \left[\frac{q(1 + p\lambda)(1 - \mu^2) + p^2\sqrt{s^2 - 1}(1 - \lambda^2)\mu}{q^2(\mu^2 - 1) + p^2(\lambda^2 - 1)} \right]. \quad (31)$$

From the field equations [1, 4] we also find

$$e^{2\gamma} = \frac{s^2(p^2\lambda^2 + q^2\mu^2) - 1}{p^2(\lambda^2 - \mu^2)}. \quad (32)$$

Substituting the expressions for p, q and λ , corresponding to (21), (24), into (30)–(32), we obtain

$$f = \frac{s(r^2 - 2Mr + a^2\mu^2)}{s(r^2 + a^2\mu^2) + 2(s - 1)M^2 - 2M[(s - 1)r + \sqrt{s^2 - 1}a\mu]}, \quad (33)$$

$$\omega = \frac{2M}{s(r^2 - 2Mr + a^2\mu^2)} \left\{ a(1 - \mu^2)[r + (s - 1)M] - (r^2 - 2Mr + a^2)\sqrt{s^2 - 1}\mu \right\}, \quad (34)$$

$$e^{2\gamma} = \frac{s^2(r^2 - 2Mr + a^2\mu^2)}{(M - r)^2 - (M^2 - a^2)\mu^2}. \quad (35)$$

4 Singularities of the DN spacetime

We can write (33) like

$$f = \frac{N}{D}, \quad (36)$$

where

$$N(r, \mu) = s[r - r_-(\mu)][r - r_+(\mu)], \quad (37)$$

$$D(r, \mu) = sr^2 - 2(s - 1)Mr + sa^2\mu^2 - 2\sqrt{s^2 - 1}Ma\mu + 2(s - 1)M^2, \quad (38)$$

with

$$r_-(\mu) = M - \sqrt{M^2 - a^2\mu^2}, \quad r_+(\mu) = M + \sqrt{M^2 - a^2\mu^2}. \quad (39)$$

The equations $r = r_-(\mu)$ and $r = r_+(\mu)$, producing $N = 0$, define respectively the inner and outer ergospheres ([2] see p. 316). The inner horizon or Cauchy horizon, and the outer horizon or event horizon are given, respectively, by $r_- = M - \sqrt{M^2 - a^2}$ and $r_+ = M + \sqrt{M^2 - a^2}$ ([2] see p. 278). $D = 0$ defines the singularities of spacetime (see Appendix). In order to have this second order polynomial equation in r satisfied, we need

$$r = \frac{1}{s} \left\{ (s-1)M \pm \sqrt{-s^2 a^2 (\mu - \mu_s)^2} \right\}, \quad (40)$$

with

$$\mu_s = \sqrt{s^2 - 1} \frac{M}{sa}. \quad (41)$$

Hence, (40) to produce real roots, one needs

$$\mu = \mu_s, \quad (42)$$

which defines the equation of a cone. In this case μ is fixed, for a given s , and we can now write D with (42) like

$$D(r) = s(r - r_s)^2, \quad (43)$$

where

$$r_s = \frac{(s-1)M}{s}. \quad (44)$$

Then, according to (43), the only possible singular points for a given s are distributed on a sphere with radius

$$r = r_s. \quad (45)$$

But at the same time, the singular points have to satisfy (42), and the intersection of the two folds of the cone with the sphere (45) produce two rings. Hence the singularities for DN spacetime are spread along two symmetrical rings centered at the z axis. Therefore we can consider only the upper fold $\theta \in [0, \pi/2]$, remembering that we have to complete the picture by symmetry. Now, if we eliminate the parameter s between the two equations (42) and (45), we obtain

$$r = M - \sqrt{M^2 - a^2 \mu^2}, \quad (46)$$

which we recognize as the equation of the inner ergosphere given just after (39). Hence, each value of s corresponds to a ring singularity, which is a circle, being the intersection of the inner ergosphere with the cone (42), see Fig. 1. The two circular rings are centered on the z axis at $\pm z_s$, where

$$z_s = r_s \mu_s = (s-1) \sqrt{s^2 - 1} \frac{M^2}{s^2 a}, \quad (47)$$

and the radius R_s of the two rings is

$$R_s = r_s \sin \theta_s = r_s \sqrt{1 - \mu_s^2} = \frac{(s-1)M}{s} \sqrt{1 - \frac{(s^2 - 1)M^2}{s^2 a^2}}. \quad (48)$$

We can say that s parametrizes the singular rings, intersections of the inner ergosphere with a continuous foliation of planes orthogonal to the z axis. For each value of s there is

a different spacetime, see (33)–(35). In particular for $s = 1$ we have the Kerr spacetime. However the inner ergosphere is the same, as well as the outer ergosphere and the two horizons, for all these metrics, i.e., for any s (see Fig. 2). Only the ring singularity changes with s , and each ring singularity belongs to the inner ergosphere.

By definition $0 \leq \mu_s \leq 1$, hence from (41) we have

$$1 \leq s \leq s_{max} = \frac{M}{\sqrt{M^2 - a^2}}, \quad (49)$$

and from (44) we have

$$0 \leq r_s \leq r_{smax} = r_- = M - \sqrt{M^2 - a^2}. \quad (50)$$

For $s = 1$, which produces Kerr metric, we have from (44) $r_s = 0$, or from (48) $R_s = 0$. Hence the Kerr limit minimizes the radius of the ring singularity and reduces the two ring singularities to just one. The other metric that minimizes the radius (but produces two symmetrical ring singularities) happens when

$$s = s_{max} = \frac{M}{\sqrt{M^2 - a^2}}, \quad (51)$$

giving, from (41), $\mu_s = 1$ and, from (48), $R_s = 0$.

We can calculate the maximum radius, R_{smax} of the ring singularities given by (48) $R_s(s)$. Calculating dR/ds we find the maximum which is for s given by

$$s = \frac{-1 + \sqrt{1 + 8(1 - a^2/M^2)}}{2(1 - a^2/M^2)}. \quad (52)$$

If $a \ll M$ we see from (49) that $1 \leq s \leq s_{max} \approx 1 + a^2/(2M^2)$, hence up to first order $O(a/M)$ the spacetime reduces to Kerr spacetime. On the other hand, if $a = 0$ and $s \neq 1$ we see that the spacetime reduces to Taub-NUT spacetime ([4] see p. 387) and $D(r) = 0$ has no real roots for r , demonstrating that there are no singularities in this case.

Finally, for the so-called “extreme black hole” ($a = M$), equations (41), (47)–(49) give

$$\begin{aligned} \mu_s &= \frac{\sqrt{s^2 - 1}}{s}, & z_s &= \frac{M(s - 1)\sqrt{s^2 - 1}}{s^2}, \\ R_s &= \frac{M(s - 1)}{s^2}, & 1 &\leq s < \infty \end{aligned} \quad (53)$$

respectively, see Fig. 3 and 4. The metric which maximizes the radius of the ring singularity is obtained from (52) for $s = 4$, and we obtain in this case

$$\mu_s = \frac{\sqrt{15}}{4} \text{ or } \theta_s \simeq 14,48^\circ, \quad r_s = \frac{3M}{4}, \quad z_s = \frac{3\sqrt{15}M}{16}, \quad R_s = \frac{3M}{16}. \quad (54)$$

5 Conclusion

The usual presentation of the DN solution obtained from the Ernst equation introduces another BL transformation (11) instead of the Kerr's one (24), and new parameters (p_{DN}, q_{DN}) which are functions of (M, a, l) and linked by the relation (8) of Kerr's type.

In our interpretation of the DN solution, we keep the BL coordinate transformation of Kerr (24), but we introduce new parameters (p, q) depending on (M, a, s) and linked by relation (23) which is no longer of Kerr's type.

Hence, the DN solution of the field equations for a given source (a, M) constitutes a family of metrics which can be parametrized by a dimensionless parameter s defined by equation (20). The Kerr solution ($s = 1$) belongs to this family.

We call generically "DN black hole" the set of ergospheres, horizons and singularities of this family.

Then, the only change that s introduces on the DN black hole structure concerns the singularities. The ergospheres and horizons are the same for each metric of the DN family, i.e., whatever s , in particular for the Kerr metric ($s = 1$). s parametrizes only each ring singularity, which always belongs to the inner ergosphere, including the limiting case of Kerr ($s = 1$).

So the Kerr metric appears as the one which minimizes the ring singularity.

Appendix

Here we present the components of $R_{\alpha\beta\gamma\delta}$ for the metric (25), transformed in spherical coordinates, with (33)–(35). The convention used for the Riemann tensor is $(R^\alpha{}_{\beta\gamma\delta} = -\Gamma^\alpha{}_{\beta\gamma,\delta} + \dots)$. Since the expressions become too long, we restrict to present only the denominators $d[R_{\alpha\beta\gamma\delta}]$ of its non null components. We use the definitions (37)–(38) and $\Delta = r^2 - 2Mr + a^2$ producing the following components :

$$\begin{aligned}
 d[R_{t\phi t\phi}] &= 4sN^2D^3, \\
 d[R_{t\phi\theta r}] &= 4\Delta ND^2, \\
 d[R_{t\theta t\theta}] &= d[R_{t\theta tr}] = d[R_{trtr}] = 4\Delta ND^3, \\
 d[R_{t\theta\phi\theta}] &= d[R_{t\theta\phi r}] = d[R_{tr\phi\theta}] = d[R_{tr\phi r}] = 4\Delta N^2D^3, \\
 d[R_{\phi\theta\phi\theta}] &= d[R_{\phi\theta\phi r}] = d[R_{\phi r\phi r}] = 4\Delta N^3D^3, \\
 d[R_{\theta r\theta r}] &= 4\Delta D.
 \end{aligned} \tag{55}$$

We see that the denominators of all components of the Riemann tensor become null only if $D = 0$, i.e., when the relations (42)–(43), (45)–(46) are satisfied.

When $s = 1$ we reobtain the components of the Riemann tensor for Kerr spacetime.

The calculation of the invariants of curvature (R^2 , $R^{\alpha\beta}R_{\alpha\beta}$, $R^{\alpha\beta\gamma\delta}R_{\alpha\beta\gamma\delta}$, etc) confirms that the condition $D = 0$ determines the singularities of DN spacetime.

Acknowledgments

R. Colistete Jr. would like to thank CAPES of Brazil for financial support. N. O. Santos would like to thank the Laboratoire de Gravitation et Cosmologie Relativistes of Université Pierre et Marie Curie where part of this work has been done, for financial aid and hospitality.

References

- [1] F. J. Ernst, *Phys. Rev.* **167**, 166 (1968).
- [2] S. Chandrasekhar, *The Mathematical Theory of Black Holes* (Oxford University Press, Oxford 1983).
- [3] M. Demiański and E.T. Newman, *Bull. Acad. Polon. Sci. Ser. Math. Astro. Phys.* **14**, 653 (1966).
- [4] M. Carmeli, *Classical Fields: General Relativity and Gauge Theory* (John Wiley and Sons, New York 1982).
- [5] A. Papapetrou, *Ann. der Phys.* **12**, 309 (1953).




Figure 1: Inner ergosphere of DN spacetime for the following values, $a = 4$ and $M = 4.2$, of the parameters. The vertical axis of revolution is the z axis. The orthogonal plane xy contains the angle ϕ , and θ is the angle between the z axis and the position vector of a point in the space. The intersections of the planes $z = cte$ with the inner ergosphere are the ring singularities. For each value of s there are two ring singularities, symmetrical with respect to the plane xy . For the Kerr metric ($s = 1$) the ring singularity is reduced to the origin O . For s_{max} , the two ring singularities are the poles on the z axis.




Figure 2: Ergospheres (with solid lines) and horizons (with dashed lines) of DN spacetime for the following values, $a = 4$ and $M = 4.2$, of the parameters. We plot here the intersections of the plane $\phi = cte$ with the surfaces of revolution, obtained from the rotation of these curves around the z axis. The radius of Cauchy horizon is here $r_- \simeq 2.92$, and the radius of the event horizon $r_+ \simeq 5.48$.




Figure 3: Inner ergosphere of a extreme DN black-hole for the following values, $a = M = 4$, of the parameters. The intersections of the planes $z = cte$ with the inner ergosphere are the ring singularities. For each value of s there are two ring singularities, symmetrical with respect to the plane xy .




Figure 4: Ergospheres (with solid lines) and horizons (with dashed lines) of a extreme DN black-hole for the following values, $a = M = 4$, of the parameters. We plot here the intersections of the plane $\phi = cte$ with the surfaces of revolution, obtained from the rotation of these curves around the z axis. Notice the continuity of the two ergospheres and of their slopes. The two horizons are the same, $r_{\pm} = M = 4$.




Figure 5: Ergospheres and horizons of DN spacetime for the following values, $a = 4$ and $M = 4.2$, of the parameters.




Figure 6: Ergospheres and horizons of a extreme DN black-hole for the following values, $a = M = 4$, of the parameters.