

CBPF-NF-012/85

CRITICALITY OF THE D=2 QUANTUM HEISENBERG  
FERROMAGNET WITH QUENCHED RANDOM ANISOTROPIC

by

Ananias M. Mariz<sup>\*\*</sup> and C. Tsallis<sup>\*\*</sup>

<sup>+</sup>Permanent address:

Centro Brasileiro de Pesquisas Físicas - CNPq/CBPF  
Rua Dr. Xavier Sigaud, 150  
22290 - Rio de Janeiro, RJ - Brasil

<sup>\*</sup>Departamento de Física  
Universidade Federal do Rio Grande do Norte  
59000 - Natal, RN - Brasil

ABSTRACT

We consider the square-lattice spin 1/2 anisotropic Heisenberg ferromagnet with interactions whose symmetry can independently (quenched model) and randomly be of two competing types, namely the isotropic Heisenberg type and the Ising one. Within a real space renormalization group framework, we perform a quite precise numerical calculation of the critical frontier, and establish its main asymptotic behaviors. We also characterize the relevant universality classes, through the analysis of the correlation length critical exponent.

Key-words: Heisenberg ferromagnet; magnetic anisotropy; phase diagram; universality classes.

In recent years, several attempts have been made to study critical properties of magnetic systems characterized by interactions belonging to competing symmetries (detailed theoretical and experimental information can be respectively found in Refs. [1, 2] and [3] and references therein). A particularly interesting case is that where spins on a regular lattice might be coupled through uniaxial, planar or spherical interactions, whose respective prototypes are the Ising, isotropic XY and isotropic Heisenberg models. Such situations have already been experimentally encountered in antiferromagnetic systems like  $\text{Fe}_{1-x}\text{Co}_x\text{Br}_2$  [4] (Ising - XY competition) and  $\text{Rb}_2\text{Co}_x\text{Mn}_{1-x}$  [5, 6] (Ising-Heisenberg competition). From the theoretical standpoint, Ising-Heisenberg mixtures have been studied, for quenched random site systems, within effective field frameworks [7], and, for  $D=2$  quenched random bond systems, with high-temperature series techniques [2, 8]. For this second case, a continuous variation of the susceptibility critical exponent  $\gamma$  with concentration was obtained. As pointed out by Pekalski himself [2, 8], this result is clearly unsatisfactory; indeed, symmetry arguments strongly suggest that the universality class corresponding to the system under analysis should be, almost everywhere in the critical frontier, that of the Ising model.

In the present paper we study the phase diagram and universality classes of the quenched bond-random spin 1/2 anisotropic Heisenberg ferromagnet in square lattice, each bond of which being either an isotropic Heisenberg interaction or an Ising-like one. The formalism we use is a real space renormalization group (RG) one, which has recently been developed [9, 10] for quantum spin systems, and whose performance has proved to be quite reliable (both qualitatively and quantitatively for square lattice).

We consider the following dimensionless Hamiltonian:

$$\mathcal{X} = \sum_{\langle i, j \rangle} K_{ij} [\sigma_i^Z \sigma_j^Z + (1 - \Delta_{ij}) (\sigma_i^X \sigma_j^X + \sigma_i^Y \sigma_j^Y)] \quad (1)$$

where  $\langle i, j \rangle$  denotes first neighbors on a square lattice, the  $\sigma$ 's are the Pauli operators,  $K_{ij} \equiv J_{ij}/k_B T > 0$  ( $J_{ij}$  is the coupling constant) is the same for all bonds, and  $\Delta_{ij} \in [0, 1]$  is the random anisotropy parameter. For the limiting value  $\Delta_{ij} = 1$  ( $\Delta_{ij} = 0$ ) we recover the Ising (isotropic Heisenberg) model. The randomness of the problem is described by the following probability law:

$$P(K_{ij}, \Delta_{ij}) = [p \delta(\Delta_{ij} - \Delta) + (1 - p) \delta(\Delta_{ij})] \delta(K_{ij} - K) \quad (2)$$

with  $0 \leq p \leq 1$ ,  $0 \leq \Delta \leq 1$  and  $K \equiv J/k_B T > 0$ . The particular case  $\Delta = 1$  corresponds to the Ising-Heisenberg mixture analyzed in Refs. [2] and [8].

To construct the RG we follow along the lines of Refs. [9] and [10], renormalizing the cluster of Fig. 1(a) into that of Fig. 1(b) (the linear scale factor being consequently  $b=2$ ); both clusters are self-dual, therefore particularly performant for the square lattice.

We associate the binary distribution (2) with each one of the 5 bonds of cluster 1(a). Consequently  $2^5$  different configurations are possible (some of them being topologically equivalent). Each configuration is characterized by the set  $(\{K_{ij}^{(\ell)}\}, \{\Delta_{ij}^{(\ell)}\})$  with  $\ell = 1, 2, \dots, 5$ . With each configuration we associate  $K_H(\{K_{ij}^{(\ell)}\}, \{\Delta_{ij}^{(\ell)}\})$  and  $\Delta_H(\{K_{ij}^{(\ell)}\}, \{\Delta_{ij}^{(\ell)}\})$  by imposing [9,10]

$$e^{\mathcal{X}'_{12}} = \text{Tr}_{3,4} e^{\mathcal{X}_{1234}} \quad (3)$$

Where  $\mathcal{X}_{1234}$  and  $\mathcal{X}'_{12}$  are the Hamiltonians corresponding to the clusters of Figs. 1(a) and 1(b) respectively. These two Hamilto-

nians are precisely of the same type indicated in Eq.(1) (i.e., the present RG generates no new types of terms). Although imposition (3) is a very natural one, its operational implementation is rather complex as it involves the computational treatment of a 16 X 16 matrix (associated with  $\mathcal{C}_{1234}$ ); practical details on the procedure can be found in Refs.[9,10].

The renormalized parameters  $K_{ij}$  and  $\Delta_{ij}$  are now associated with a distribution law  $P_H$  which is no more binary. It has in fact 14 different  $\delta$ 's, and is given by

$$P_H(K_{ij}, \Delta_{ij}) = \int \prod_{\ell=1}^5 [dK_{ij}^{(\ell)} d\Delta_{ij}^{(\ell)} P(K_{ij}^{(\ell)}, \Delta_{ij}^{(\ell)})] \delta(K_{ij} - K_H) \delta(\Delta_{ij} - \Delta_H) \quad (4)$$

Under successive renormalizations the distribution law becomes more and more complex. It is in principle possible to keep track of its evolution up to an eventual stabilization, but, following along the lines of previous similar theories (e.g., Ref.[11]), we shall instead approximate it by the following binary one:

$$P'(K_{ij}, \Delta_{ij}) = [p' \delta(\Delta_{ij} - \Delta') + (1-p') \delta(\Delta_{ij})] \delta(K_{ij} - K') \quad (5)$$

where  $p'$ ,  $K'$  and  $\Delta'$  have to be found as functions of  $p$ ,  $K$  and  $\Delta$ . To do this we impose that the main momenta are preserved through renormalization. More specifically we demand

$$\langle K_{ij} \rangle_{P'} = \langle K_{ij} \rangle_{P_H} \equiv g_1(p, K, \Delta) \quad (6)$$

$$\langle \Delta_{ij} \rangle_{P'} = \langle \Delta_{ij} \rangle_{P_H} \equiv g_2(p, K, \Delta) \quad (7)$$

$$\langle \Delta_{ij}^2 \rangle_{P'} = \langle \Delta_{ij}^2 \rangle_{P_H} \equiv g_3(p, K, \Delta) \quad (8)$$

where  $\langle \dots \rangle$  denotes the standard mean values. While Eqs. (6) and (7) are quite natural choices, Eq. (8) has been adopted in order to decouple  $p$  and  $\Delta$ . The set of Eqs.(6)-(8) immediately

yield

$$K' = g_1 \quad (9)$$

$$\Delta' = g_3/g_2 \quad (10)$$

$$F' = g_2/\Delta' \quad (11)$$

which constitute the RG recursive relations which close the formalism. Iteration (in the  $(p, 1/K, \Delta)$  space, for instance) provides the para-ferromagnetic critical surface (see Figs. 2(a) and 2(b) for selected cuts of this surface) as well as the set of universality classes. For  $p=1$  we recover the results obtained in Ref. [9]. When  $p$  and/or  $\Delta$  increase, the lower symmetry (Ising) becomes dominant, consequently the critical temperature is expected to increase, as exhibited in Fig. 2. In the neighborhood of  $p = \Delta = 1$ , we obtain the following asymptotic behaviors:

$$\frac{T_c(p=1, \Delta) - T_c(p, \Delta)}{T_c(p=1, \Delta)} \sim A(\Delta) (1-p) \quad (p \rightarrow 1) \quad (12)$$

and

$$\frac{T_c(p, \Delta=1) - T_c(p, \Delta)}{T_c(p, \Delta=1)} \sim B(p) (1-\Delta) + C(p) (1-\Delta)^2 \quad (\Delta \rightarrow 1) \quad (13)$$

where  $A(\Delta)$ ,  $B(p)$  and  $C(p)$  are shown in Figs. 3 and 4. Numerical difficulties prevented us from a reliable description of the  $T \rightarrow 0$  asymptotic behaviors.

Two non trivial fixed points belong to the critical surface, namely the isotropic Heisenberg one (at  $(p, k_B T/J, \Delta) = (1, 0, 0)$ ), and the Ising one (at  $(p, k_B T/J, \Delta) = (1, 2.269\dots, 1)$ ): both are located at the exact values [12, 13]. These points charac-

terize the unique two universality classes of this problem; indeed the RG flow shows that the universality class is that of the Ising (isotropic Heisenberg) model for all points of the critical surface at finite (vanishing) temperature. These results disagree with those obtained (for the antiferromagnetic system) by high-temperature series [2, 8], and confirm the symmetry-based intuitive expectation.

The correlation length critical exponents are given by

$$\nu_i = \ln b / \ln \lambda_i \quad (i = T, \Delta) \quad (14)$$

where  $\lambda_i$  is the relevant eigenvalue ( $\lambda_i \geq 1$ ) of the Jacobian matrix  $\partial(p', K', \Delta') / \partial(p, K, \Delta)$  calculated at the corresponding fixed point. We obtain  $\nu_T \approx 1.15$  (the exact value equals 1 [14]) for the Ising fixed point, and  $\nu_T = \infty$  (which reproduces the exact value [15]) and  $\nu_\Delta \approx 1.22$  (we found no other value in the literature for comparison) for the isotropic Heisenberg fixed point.

To conclude let us say that we believe that the present approximation of the critical surface should be a numerically quite reliable one.

We acknowledge fruitful discussions with A.O. Caride; one of us (A.M.M.) also acknowledges useful remarks from C.A.P. Galvão.

REFERENCES

1. A. Aharony, J. Magn. Mater. 31-34, 1432 (1983).
2. A. Pekalski, in "Static Critical Phenomena in Inhomogeneous Systems", Lectures Notes in Physics 206, pp. 158, Springer-Verlag. (1984).
3. K. Katsumata, J. Magn. Mater. 31-34, 1435 (1983).
4. K. Katsumata, J. Tuchendler and S. Legrand, Solid State Comm. 49, 1, 83 (1984).
5. H. Ikeda, T. Riste and G. Shirane, J. of Phys. Soc. of Japan, 49, 2, 504 (1981).
6. Y. Ajiro, K. Adachi and M. Maketa, Solid State Comm. 37, 2, 449 (1981).
7. A. Komoda and A. Pekalski, J. Phys. C. 14, L1067 (1981).
8. A. Pekalski, J. Phys. C. 10, 4785 (1977).
9. A.O. Caride, C. Tsallis and S.I. Zanette, Phys. Rev. Lett 51, 145 (1983); 51, 616 (1983).
10. A.M. Mariz, C. Tsallis and A.O. Caride (Submitted to J. Phys. C)
11. R.B. Stinchcombe and B.P. Watson, J. Phys. C. 9, 3221 (1976).
12. N.D. Mermin and H. Wagner, Phys. Rev. Lett. 17, 1133 (1966).
13. L. Onsager, Phys. Rev. 65, 117 (1944).
14. T.T. Wu, Phys. Rev. 149, 380 (1966).
15. A.M. Polyakov, Phys. Lett. 59B, 79 (1975).



FIGURE CAPTIONS

- FIG. 1 - Self-dual two-terminal clusters. The terminal and internal nodes are respectively denoted by 0 and .
- FIG. 2 - (a) Cuts of the critical frontier for selected values of  $\Delta$ ; (b) Cuts of the critical frontier for selected values of  $p$ . P and F are respectively the paramagnetic and ferro magnetic phases.
- FIG. 3 -  $\Delta$ -dependence of the asymptotic coefficient  $A(\Delta)$  given by Eq. (12), [ $A(1) \approx 0.32$  and  $A(0) \approx 0.18$ ].
- FIG. 4 -  $p$ -dependence of the coefficients  $B(p)$  and  $C(p)$  given by Eq. (13); [ $B(1) \approx 0$ ,  $C(1) \approx 0.29$  and  $B(0) \approx 0.17$ ,  $C(0) \approx 0.05$ ].




FIG. 1




FIG. 2(a)




FIG. 2(b)

-11-




FIG. 3




FIG. 4