

CBPF-NF-044/89

HIGHER ORDER EQUATIONS OF MOTION

by

C.G. BOLLINI<sup>1\*</sup> and J.J. GIAMBIAGI<sup>1+</sup>

<sup>1</sup>Centro Brasileiro de Pesquisas Físicas - CBPF/CNPq  
Rua Dr. Xavier Sigaud, 150  
22290 - Rio de Janeiro, RJ - Brasil

\*Departamento de Física, Facultad de Ciencias Exactas y  
Naturales, Universidad Nacional de La Plata  
La Plata, Argentina, and Comisión de  
Investigaciones Científicas de la Provincia de  
Buenos Aires, Argentina

<sup>+</sup>Centro Latinoamericano de Física - CLAF  
Av. Wenceslau Braz 71 (fundos)  
22290 - Rio de Janeiro, RJ - Brasil

## ABSTRACT

We discuss the possibility that the motion of elementary particles be described by higher order differential equations induced by supersymmetry in higher dimensional space-time. We take the specific example of six dimensions writing the corresponding Lagrangian and equations of motion.

Key-words: Field theory; Supersymmetry; Higher order equations.

One of the main sources of ideas and discussions in the last twenty years or so, has been supersymmetry. Specially since the proof of the Haag, Lopuszanski, Sohnius - theorem [1]. In particular the Wess-Zumino model [2] has served as a basis for the construction of "relativistic" lagrangian theories that could describe the physical world.

The usual rule for writing a supersymmetric Lagrangian for a chiral superfield [3], for example, is to take

$$\phi(\theta, \bar{\theta}, x) = e^{\frac{i}{2}\theta\partial\bar{\theta}} \phi_0(\theta, x) \quad (1)$$

and write the kinetic lagrangian as:

$$\mathcal{L} = (\bar{\phi}\phi)_D \quad (2)$$

where  $( )_D$  means the highest component (maximum possible number of Grassmann variables).

We have

$$\begin{aligned} \bar{\phi}\phi &= \bar{\phi}_0 e^{i\theta\partial\bar{\theta}} \phi_0 + \text{total divergence} \\ &\cong \bar{\phi}_0 \sum_{s=0} \frac{(i\theta\partial\bar{\theta})^s}{s!} \phi_0 \end{aligned} \quad (3)$$

Now, the Grassmann variable  $\theta_\alpha$  is a Weyl spinor having

$$\omega = 2^{\frac{v}{2}-1} \quad (4)$$

independent components in  $\nu$  dimensions. (In what follows we will only consider  $\nu = \text{even number}$ ).

The sum over  $S$  in eq. (3) runs then between  $S = 0$  and  $S = \omega$ .

In four dimensions  $S$  runs from zero to two and this leads us to Lagrangians which implies at most, second order wave equations. But this is not so for higher number of dimensions. In fact the same construction leads again to expression (3) but now in six dimensions, for example where  $\omega = 4$ , the equations of motion are of the fourth order and of even greater order for  $\nu > 6$ . The mass term is introduced via a Lagrangian which is proportional to the square of the chiral field (3).

$$\mathcal{L}_m = c \phi_0^2 \Big|_F + hc \quad (5)$$

where  $F$  means the coefficient of the maximum number of  $\theta$ -variables.

It is easy to see that by defining the components  $\psi_{\alpha_1 \dots \alpha_S}$  as:

$$\phi_0 = \sum_{S=0}^{\omega} \frac{1}{S!} \theta^{\alpha_1} \dots \theta^{\alpha_S} \psi_{\alpha_1 \dots \alpha_S}(x) \quad (6)$$

the mass Lagrangian is given by:

$$\mathcal{L}_m = c \sum_{S=0}^{\omega} \psi_{\alpha_1 \dots \alpha_S} \psi_{\alpha_{S+1} \dots \alpha_{\omega}} + hc \quad (7)$$

(2) and (5) (or (7)) leads, for each component  $X$  to an equation of motion [4]

-3-

$$\left(\square^{2\omega} - m^\omega\right) X = 0 \quad (8)$$

where  $C$ , for dimensional reasons, has been substituted by  $m^\omega$ .

In four dimensions, (8) is the Klein-Gordon equation, and in six, for example, we have:

$$\left(\square^2 - m^4\right)X = 0 \quad \left(\square - m^2\right)\left(\square + m^2\right)X = 0 \quad (9)$$

The massive propagator is of course:

$$\frac{1}{(p^2)^{\frac{\omega}{2}} - m^\omega} \quad \text{and} \quad \frac{1}{(p^2)^{\frac{\omega}{2}}} \quad \text{for massless particles.} \quad (10)$$

The Fourier transforms for the massless case have the form (see Ref. [4]).

$$G = \frac{1}{R^2} \quad \text{for} \quad \nu = 4 \quad \text{or} \quad \nu = 6 \quad (11)$$

$$G = \ln R \quad \nu = 8 \quad (12)$$

$$G = R^\alpha \ln R \quad \text{with} \quad \alpha = \omega - \nu \geq 0 \quad \text{for} \quad \nu > 8 \quad (13)$$

we see that due to the fact that  $\alpha > 0$  for  $\nu > 8$  the convolution of two of these functions has not ultraviolet divergences. The same happens for the massive cases as for  $p \rightarrow \infty$  the mass term has no importance.

In this case (massive particle) there is no infrared divergences, neither. This absence of singularities is due to the fact that in a convolution between two Green-functions the number of integration variables grows linearly with the number of dimensions while the denominators grow exponentially (see (10)), so that the power of  $p$  in the denominator outnumber those in the numerator (for  $\nu > 8$ ). For example, while in eight dimensions we have  $\omega = \nu = 8$ , in ten dimensions we have  $\omega = 16$  ( $\nu = 10$ ). (not to speak of  $\nu = 26!$ ). These facts show that it is worth our while to look carefully into theories containing higher order equations such as that given by eq. (9).

However, it is well known that this kind of equations presents considerable difficulties both of a mathematical nature and of physical interpretation (see Hawking [5]). It is advisable then to study first higher order equation in the case of only one significative variable, where it is possible to address the question of unitarity in a controlled way, describing the scattering data in a precise mathematical language [6,7] and looking at the physical implications of the scattering processes.

The general linear differential equation for only one significative variable takes the form:

$$\frac{d^n \phi}{dx^n} + q_{n-2} \frac{d^{n-2} \phi}{dx^{n-2}} + \dots + q_0 \phi = z^n \phi \quad (14)$$

Where a possible  $q_{n-1} \frac{d^{n-1} \phi}{dx^{n-1}}$  has been eliminated by means of a transformation  $\psi \rightarrow f\psi$ .

This equation has  $n - 1$  independent potentials  $q_i$ , ( $i = 0 \dots n - 2$ ) when they are well-behaved they tend to zero sufficiently

-5-

rapid for  $x \rightarrow \pm \infty$ ; so that asymptotically (15) tend to

$$\frac{d^2 \phi^{(0)}}{dx^n} = z^n \phi^{(0)} \quad (15)$$

whose solution are :

$$\phi_i^{(0)} = e^{\alpha_i z x} \quad (16)$$

with  $\alpha_i^n = 1$ ,  $\alpha_i$  being thus a nth root of unity.

Eq. (15) of course, has n independent solutions that can be expressed in terms of a set of basic linearly independent solutions defined by conditions on  $x = \pm \infty$ .

We define the first Jost function  $f_1(zx)$  as the solution of (15) with the greatest rate of decrease for  $x \rightarrow -\infty$ .

The second Jost function  $f_2(zx)$  is that solution of (14) which, for  $x \rightarrow -\infty$  has the second rate of decrease (and for  $x \rightarrow \infty$  has the second rate of increase), etc., etc. See Ref. [8].

In other words.

$$f_j(zx) \rightarrow e^{\alpha_j z x} \quad (17)$$

where the roots are ordered in such a way that

$$\text{Real } \alpha_1 z > \text{Real } \alpha_2 z > \dots > \text{Real } \alpha_n z \quad (18)$$

It is clear that (18) divides the z-plane in  $2n$  regions, within each region the inequality (18) is well defined but as

argZ is varied, there are lines for which  $\text{Real } \alpha_\ell Z = \text{Real } \alpha_{\ell+1} Z$  and this defines a ray on which the order of roots is ill defined. There are  $2n$  rays dividing the  $Z$ -plane in regions with an angle  $\frac{2\pi}{2n} = \frac{\pi}{n}$ . In particular for fourth order differential equations, the  $Z$ -plane is divided in eight "octants" each one with an angle of  $\frac{\pi}{4}$  radians [9].

The Jost functions have discontinuities at those rays and furthermore they can have poles with corresponding residues. The set of all discontinuities, including poles and residues, form the so called "scattering data" of the differential equation (14). It is shown by mathematicians that the knowledge of the scattering data is equivalent to the knowledge of the differential equation. In other words the set of discontinuities (including poles and residues) determines the  $n-1$  potentials  $q_i$  of the equation.

However from the point of view of a physicist, not all the scattering data are physical. For instance, in a fourth order differential equation, only the real and imaginary axis correspond to physical data (i.e., a plane wave going to plane wave). (see ref. 9). The other rays at  $\frac{\pi}{4}$  and  $\pm \frac{3\pi}{4}$  correspond to scattering of waves that grow exponentially (or decrease) for  $x \rightarrow \pm \infty$ . These are unphysical data. Therefore in order that the equation be a physical one, the Jost functions must not present discontinuities on these rays. This implies relations to be satisfied by the potentials of equation (14) in order for it to be physically acceptable.

Summarizing: Any higher order equation of motion can not



in principle be thought as having physical significance unless some specific relations exist among the coefficients. One hopes that supersymmetric theories may provide the clue to physically meaningful higher order equations. Any way, we want to mention that by using the method  $v \rightarrow \infty$  of reference [10] in a higher order equation, we get the static limit in leading approximation and, as a second approximation a second order equation is obtained. The higher order derivatives appear in the following approximations (in a  $v^{-1}$  development). In this sense we can say that a higher order equation has a second order equation as an approximation.

With these motivating ideas we start looking for a supersymmetric theory in six dimensions which is the simplest higher order case. In this respect we like to point out that a more realistic theory that the one we are presenting here can be found in works by P. Fayet [11].

We want to find the coupling between the chiral superfield, equation (1), and a gauge superfield [12].

$$V = \sum_{s,t=2}^4 \bar{\theta}_{\dot{\alpha}_1 \dots \dot{\alpha}_4} \bar{\theta}_{\dot{\alpha}_1 \dots \dot{\alpha}_4} A_{\alpha_1 \dots \alpha_t}^{\dot{\alpha}_1 \dots \dot{\alpha}_5} \theta^{\alpha_1 \dots \alpha_t} \quad (19)$$

where we have chosen the Wess-Zumino gauge. (See [12]).

A chiral superfield strength for this field is given by

$$W_{\alpha_1 \alpha_2} = \bar{D} \bar{D}_{\alpha_1} D_{\alpha_2} V \quad (20)$$

and the corresponding Lagrangian is

$$\mathcal{L}_w = \varepsilon^{\alpha_1 \alpha_2 \alpha_3 \alpha_4} W_{\alpha_1 \alpha_2} W_{\alpha_3 \alpha_4} + \text{hc}$$

By reducing the spinor components of the gauge superfield defined by (20), using Elie Cartan's reduction recipes, we found the following tensor components.

$A_{(\mu\nu)}$ : (Graviton field)  $(\mu\nu)$  means symmetric part and the gauge transformation with parameters  $\lambda_\mu$  are:

$$A'_{(\mu\nu)} = A_{(\mu\nu)} + \partial_\mu \lambda_\nu + \partial_\nu \lambda_\mu - \eta_{\mu\nu} \partial^\alpha \lambda_\alpha \quad (21)$$

This gauge transformation can be used to simplify the Lagrangian by choosing the De Donder gauge in which  $\partial_\mu A^\mu_\nu = 2\partial^\nu A_{(\mu\nu)}$ . In this gauge:

$$\mathcal{L}'_{22} \equiv \square A_{(\mu\nu)} \square A_{(\mu\nu)} \quad (22)$$

$A_{[\mu\nu]}$  ( $[\mu\nu]$  = antisymmetric part) has the gauge transformation

$$A'_{[\mu\nu]} = A_{[\mu\nu]} + \partial^\rho \lambda_{\rho\mu\nu} \quad (23)$$

$\lambda_{\rho\mu\nu}$  = self dual and completely antisymmetric three-vector. The corresponding Lagrangian only contains the divergence of this tensor, which is gauge-invariant

$$\mathcal{L}''_{22} \approx \partial_\rho \partial^\mu A_{[\mu\nu]} \partial^\rho \partial_\sigma A^{[\sigma\nu]} \quad (24)$$

-9-

$-A_\mu^\alpha$ : (gravitino field). Under a gauge transformation it transforms as.

$$A_\mu^{\prime\alpha} = A_\mu^\alpha + (C\tilde{\delta}\gamma_\mu\tilde{\delta})^{\alpha\beta}\lambda_\beta \quad (25)$$

where  $\lambda_\beta$  is a spinor parameter, which can be adjusted so as to have a "zero gamma trace" gauge.

$$(\gamma^\mu C)_{\alpha\beta} A_\mu^\beta = 0 \quad (C = \text{transposition matrix})$$

In this gauge, the Lagrangian is:

$$\mathcal{L}_{2,3} = i\bar{\psi}^\mu \partial_\mu^\alpha \psi_\alpha^\nu + i\bar{\psi}^\mu \partial_\alpha^\nu \psi_\mu^\alpha \quad (26)$$

$-B_\mu$ : Complex gauge invariant vector field. Its Lagrangian is:

$$\mathcal{L}_{2,4} = 2\bar{B}^\mu \partial_\mu^\nu B_\nu - \bar{B}^\mu \square B_\mu \quad (27)$$

$-A_\mu$ : real vector field with gauge transformation

$$A_\mu' = A_\mu + \partial_\mu \bar{\lambda} \quad (28)$$

and the Maxwell type Lagrangian.

$$\mathcal{L}'_{33} = F^{\mu\nu} F_{\mu\nu}; F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu \quad (29)$$

$A_{\mu\rho}$  real gauge invariant self-dual three-vector field.

The Lagrangian is:

$$\mathcal{L}_{33} = G^{\mu\nu} G_{\mu\nu}; G_{\mu\nu} = \partial^\rho A_{\rho\sigma\lambda}$$

$B^\alpha$ : photino field. A complex gauge invariant spinor field with Lagrangian

$$\mathcal{L}_{3,4} = iB^\alpha \partial_\alpha \bar{B}_{\dot{\alpha}}$$

Finally: an auxiliary gauge invariant scalar field  $D$  with Lagrangian

$$\mathcal{L}_{44} = D^2.$$

Of course, for a more realistic theory it is necessary to work with non-abelian gauge group, in particular Yang-Mills type theories for the standard model and also introducing super gravity; but our aim is not so much to construct a realistic theory, but rather to show the plausibility of using higher order equations in a theory that could be renormalizable and unitary.

- [1] R. Haag, J. Lopuszanski, and M. Sohnius: Nucl. Phys. B88 (1975) 257.
- [2] J. Wess and B. Zumino: Nucl. Phys. B70 (1974) 39.
- [3] J. Wess and J. Bagger: Supersymmetry and Supergravity. The Princeton University Press, 1983.
- [4] C.G. Bollini and J.J. Giambiagi: Phys. Rev. D V32 (1985) 3316.
- [5] S.W. Hawking: Preprint. Univ. of Cambridge. Dept. of Applied Math. and Theor. Phys. September (1985).
- [6] R. Beals, P. Deift and C. Tomei: Atas da VI<sup>a</sup> ELAM, IMPA, Rio de Janeiro, Brasil (1986).
- [7] P. Deift, C. Tomei and E. Trubowicz: Comm. Pure and Appl. Math. XXXV (1982) 567.
- [8] C. Tomei: Notes for a book on the general linear differential equation. PUC. Rio de Janeiro Brasil (1986).  
We are indebted to the author for letting us know the notes previous to publication.
- [9] C.G. Bollini and J.J. Giambiagi: Il Nuovo Cimento. 98A (1987) 151.
- [10] E. Witten: Physics Today, July (1980) 38.
- [11] P. Fayet: Phys. Scripta T15 (1987) 46.  
B. Delamotte and P. Fayet: Phys. Letters 195 (1987) 563.  
P. Fayet: Phys. Letters 192 (1987) 395.  
P. Fayet: Phys. Letters 159B (1985) 121.
- [12] C.G. Bollini and J.J. Giambiagi: Phys. Rev. D 39 (1989) 1169.