

Nonlinear Schroedinger equation in the presence of uniform acceleration

A. R. Plastino and C. Tsallis

Citation: *J. Math. Phys.* **54**, 041505 (2013); doi: 10.1063/1.4798999

View online: <http://dx.doi.org/10.1063/1.4798999>

View Table of Contents: <http://jmp.aip.org/resource/1/JMAPAQ/v54/i4>

Published by the [American Institute of Physics](#).

Additional information on J. Math. Phys.


Journal Homepage: <http://jmp.aip.org/>

Journal Information: http://jmp.aip.org/about/about_the_journal

Top downloads: http://jmp.aip.org/features/most_downloaded

Information for Authors: <http://jmp.aip.org/authors>

ADVERTISEMENT



Maple 17
The Essential Tool for Mathematics and Modeling

The most comprehensive support for Physics in any mathematical software package

- State-of-the-art environment for algebraic computations in physics
- Access to Maple's full mathematical power, programming language, visualization routines, and documentation creation tools
- The only system with the ability to handle a wide range of physics computations as well as pencil-and-paper style input and textbook-quality display of results
- A programming library that gives access to almost 100 internal commands to write programs or extend the capabilities of the Physics package

InertiaTensor := $\sum_{k=1}^n m_k r_k^2$
 $(1 + 2 E C) \frac{\partial}{\partial r} R C$

[Click to learn more >>](#) World-leading tools for performing calculations in theoretical physics

Nonlinear Schroedinger equation in the presence of uniform acceleration

A. R. Plastino^{1,2} and C. Tsallis^{3,4}

¹*CeBio y Secretaria de Investigaciones, Universidad Nacional del Noroeste de la Prov. de Buenos Aires -UNNOBA y CONICET, R. Saenz Peña 456, Junin, Argentina*

²*Instituto Carlos I de Fisica Teorica y Computacional, Universidad de Granada, 18071 Granada, Spain*

³*Centro Brasileiro de Pesquisas Físicas and National Institute of Science and Technology for Complex Systems, Rua Xavier Sigaud 150, 22290-180 Rio de Janeiro-RJ, Brazil*

⁴*Santa Fe Institute, 1399 Hyde Park Road, Santa Fe, New Mexico 87501, USA*

(Received 24 August 2012; accepted 15 March 2013; published online 8 April 2013)

We consider a recently proposed nonlinear Schroedinger equation exhibiting soliton-like solutions of the power-law form $e_q^{i(kx-wt)}$, involving the q -exponential function which naturally emerges within nonextensive thermostatics [$e_q^z \equiv [1 + (1 - q)z]^{1/(1-q)}$, with $e_1^z = e^z$]. Since these basic solutions behave like free particles, obeying $p = \hbar k$, $E = \hbar\omega$, and $E = p^2/2m$ ($1 \leq q < 2$), it is relevant to investigate how they change under the effect of uniform acceleration, thus providing the first steps towards the application of the aforementioned nonlinear equation to the study of physical scenarios beyond free particle dynamics. We investigate first the behaviour of the power-law solutions under Galilean transformation and discuss the ensuing Doppler-like effects. We consider then constant acceleration, obtaining new solutions that can be equivalently regarded as describing a free particle viewed from an uniformly accelerated reference frame (with acceleration a) or a particle moving under a constant force $-ma$. The latter interpretation naturally leads to the evolution equation $i\hbar \frac{\partial}{\partial t} \left(\frac{\Phi}{\Phi_0} \right) = -\frac{1}{2-q} \frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} \left[\left(\frac{\Phi}{\Phi_0} \right)^{2-q} \right] + V(x) \left(\frac{\Phi}{\Phi_0} \right)^q$ with $V(x) = max$. Remarkably enough, the potential V couples to Φ^q , instead of coupling to Φ , as happens in the familiar linear case ($q = 1$). © 2013 American Institute of Physics. [<http://dx.doi.org/10.1063/1.4798999>]

The importance of symmetries in physics can hardly be overestimated. A proper understanding of the relevant symmetries exhibited by a physical phenomenon encapsulates some of its most essential features, and usually leads to the most elegant and compact mathematical formulation of the associated physical laws. Many important symmetries are directly related to changes in the reference frame employed to describe physical systems. Such is the case of the dynamical symmetries associated with the Galilean transformation between inertial frames, playing a central role in Newtonian mechanics and non-relativistic quantum mechanics; the Lorentz transformation plays an analogous role in special relativity, both at the classical and at the quantum levels. We can also mention the invariance of statistical mechanics and thermodynamics under uniform translation of the energy spectrum, which leads to the full freedom of choosing the zero level of energies in any physical system (the above list of examples is, of course, far from complete). A remarkable feature of symmetry or invariance principles is that they constitute powerful heuristic tools to extend knowledge about particular or restricted instances of the behaviour of physical systems to more general situations or scenarios. The most spectacular example of this approach was probably Einstein's use of his Principle of Equivalence in the development of General Relativity.

Symmetries with important physical implications are also observed in pure mathematics. We may focus, for example, on the Central Limit Theorem: the stability of Gaussians in the space of distributions with regard to adding (into the system) a finite or even infinite number of nearly

independent random variables with finite variance is at the heart of their robustness, and therefore of their ubiquity in nature. The expansibility property of an entropic functional (that is, its invariance when adding new microscopic configurations with zero probability) constitutes another relevant example. It plays a special role within the (Shannon-Khinchine) set of axioms that uniquely determine the mathematical form of the Boltzmann-Gibbs (BG) additive entropy, basis of BG statistical mechanics.

Symmetry considerations arise naturally in the study of both linear and nonlinear dynamics. In the latter case, however, the analysis of the relevant symmetries is usually much more difficult. The physically relevant nonlinear evolution equations are highly diverse and describe wide classes of physical systems: see for instance Refs. 1–4. Among the most studied nonlinear differential equations we have nonlinear versions of the Schroedinger^{1,5,6} and Fokker-Planck^{2,7,8} ones. Here we focus primarily on the soliton-like solutions of the recently proposed nonlinear Schroedinger equation⁵ inspired by nonextensive statistical mechanics and the associated nonadditive entropies.^{9–11} The nonlinear evolution equation advanced in Ref. 5 (and the above mentioned soliton-like solution) is implicitly assumed to hold with respect to an inertial reference frame. Our main aim in the present work is to investigate the form of these solutions when described with respect to a uniformly accelerated reference frame. We consider first the behaviour of the soliton-like solutions under Galilean transformations connecting inertial frames. We then tackle the case of accelerated frames.

The aforementioned soliton-like solutions are referred to as q -plane waves and may be relevant in diverse areas of physics, including nonlinear optics, superconductivity, plasma physics, and dark matter.^{6,12} The theory within which q -plane waves emerged generalizes the BG entropy and statistical mechanics, through the introduction of an index q ($q \rightarrow 1$ recovers the BG case). Along this line, considerable progress was achieved, leading to generalized functions, distributions, various equations of physics, and new forms of the Central Limit Theorem.¹³ In particular, the q -Gaussian distribution, which generalizes the standard Gaussian, appears naturally by extremizing the q -entropy,⁹ or from the solution of the corresponding nonlinear Fokker-Planck equation,¹⁴ and has been successfully applied to the analysis of recent experimental results in various fields.¹⁰ Among others, we may mention: (i) The velocities of cold atoms in dissipative optical lattices;¹⁵ (ii) The velocities of particles in quasi-two dimensional dusty plasma;¹⁶ (iii) Single ions in radio frequency traps interacting with a classical buffer gas;¹⁷ (iv) The relaxation curves of Ruderman-Kittel-Kasuya-Yosida (RKKY) spin glasses, like CuMn and AuFe;¹⁸ (v) Transverse momenta distributions at Large Hadron Collider (LHC) experiments.¹⁹

Herein we discuss the following q -generalized Schroedinger equation for a d -dimensional free particle of mass m :⁵

$$i\hbar \frac{\partial}{\partial t} \left[\frac{\Phi(\vec{x}, t)}{\Phi_0} \right] = -\frac{1}{2-q} \frac{\hbar^2}{2m} \nabla^2 \left[\frac{\Phi(\vec{x}, t)}{\Phi_0} \right]^{2-q} \quad (q \geq 1), \quad (1)$$

where Φ_0 guarantees the correct physical dimensionalities for all terms (this scaling becomes irrelevant only for the linear equation, i.e., $q = 1$). Notice that, in spite of the apparent singularity in $q = 2$, it can be verified after some algebra that Eq. (1) is well defined for this value of the parameter q . It should be emphasized that Eq. (1) has important structural differences with respect to other nonlinear Schroedinger equations appearing in the literature, which contain extra nonlinear terms (usually a cubic nonlinearity) involving only the wave function itself. In contrast with those equations, the evolution equation (1) considered here has a power-law nonlinearity inside the Laplacian term. This gives rise to q -plane wave solutions that are compatible with the Planck and de Broglie relations and behave like free particles. These solutions are expressed in terms of the q -exponential function $\exp_q(u)$ which, for a pure imaginary iu , is defined as the principal value of

$$\exp_q(iu) = [1 + (1 - q)iu]^{1/(1-q)}; \quad \exp_1(iu) \equiv \exp(iu). \quad (2)$$

The above function satisfies,²⁰

$$\begin{aligned} \exp_q(\pm iu) &= \cos_q(u) \pm i \sin_q(u), \\ \cos_q(u) &= \rho_q(u) \cos \left\{ \frac{1}{q-1} \arctan[(q-1)u] \right\}, \end{aligned}$$

$$\begin{aligned} \sin_q(u) &= \rho_q(u) \sin \left\{ \frac{1}{q-1} \arctan[(q-1)u] \right\}, \\ \rho_q(u) &= [1 + (1-q)^2 u^2]^{1/[2(1-q)]}, \end{aligned} \quad (3)$$

$$\exp_q(iu) \exp_q(-iu) = [\rho_q(u)]^2 = \exp_q(-(q-1)u^2),$$

$$\exp_q(iu_1) \exp_q(iu_2) \neq \exp_q[i(u_1 + u_2)], \quad (q \neq 1). \quad (4)$$

As a consequence of Eqs. (3) and (4), a q -exponential with a pure imaginary argument, $\exp_q(iu)$, presents an oscillatory behaviour with a u -dependent amplitude $\rho_q(u)$. The function $\exp_q(iu)$ complies with the physically important property of square integrability for $1 < q < 3$, whereas the concomitant integral diverges in both limits $q \rightarrow 1$ and $q \rightarrow 3$ and also for $q < 1$.²¹

The d -dimensional q -plane wave is given by

$$\Phi(\vec{x}, t) = \Phi_0 \exp_q \left[i(\vec{k} \cdot \vec{x} - \omega t) \right]. \quad (5)$$

If we take into account that $d \exp_q(z)/dz = [\exp_q(z)]^q$ and $d^2 \exp_q(z)/dz^2 = q[\exp_q(z)]^{2q-1}$ we obtain, for the d -dimensional Laplacian,

$$\nabla^2 \left(\frac{\Phi}{\Phi_0} \right) = -q \left(\sum_{n=1}^d k_n^2 \right) \left(\frac{\Phi}{\Phi_0} \right)^{2q-1}. \quad (6)$$

Now, inserting the q -plane wave ansatz (5) into the NL Schroedinger equation (1), we verify that the q -plane wave is indeed a solution provided that the frequency ω and the momentum k satisfy the relation $\omega = \frac{\hbar k^2}{2m}$. Equivalently, if one makes,⁵ according to the celebrated de Broglie and Planck relations, the identifications $\vec{k} \rightarrow \vec{p}/\hbar$ and $\omega \rightarrow E/\hbar$, one verifies that the q -plane wave is a solution of Eq. (1) with $E = p^2/2m$, thus *preserving the energy spectrum of the free particle for all values of q* .

As already mentioned, Eq. (1) differs from previous formulations^{1,22} where new nonlinear terms are added to the two existing linear terms. The main differences between Eq. (1) and other proposals for NL Schroedinger equations are: (i) Instead of adding an extra term in which the nonlinearity is introduced, we generalize the spatial second-derivative term; (ii) The equation, together with the proposed solution, is easily extended from one to d dimensions; (iii) The corresponding solution of Eq. (1) manifests nonlinearity in both space and time, through a modulation in these two variables, which keeps the norm finite for all (\vec{x}, t) ; (iv) The well-known energy spectrum of a free particle is preserved for all q . Therefore Eq. (1), together with its solution Eq. (5), can be considered as candidates for describing interesting types of physical phenomena.

Let us now investigate how the q -plane wave solution is transformed under two basic types of changes in the reference frame. As already mentioned, we consider first a Galilean transformation connecting inertial reference frames. Then, we analyze the aspect of the q -plane wave solutions when “viewed” from an uniformly accelerated reference frame. We shall consider the one-dimensional nonlinear Schroedinger equation. The extension to the d -dimensional case is straightforward.

We shall assume that the nonlinear Schroedinger equation,

$$i\hbar \frac{\partial}{\partial t'} \left[\frac{\Psi(x', t')}{\Psi_0} \right] = -\frac{1}{2-q} \frac{\hbar^2}{2m} \frac{\partial^2}{\partial x'^2} \left[\frac{\Psi(x', t')}{\Psi_0} \right]^{2-q} \quad (q \geq 1), \quad (7)$$

holds in an inertial reference frame characterized by the spatio-temporal coordinates (x', t') and that the system under consideration is described by a q -plane wave solution

$$\Psi(x', t') = \Psi_0 \exp_q [i(kx' - \omega t')]. \quad (8)$$

Let us consider now a Galilean transformation

$$t = t'; \quad x = x' - vt', \quad (9)$$

relating the original inertial frame (x', t') with a second inertial frame (x, t) that moves with respect to the former one with a uniform velocity v . To obtain a “new” solution $\Phi(x, t)$ to the nonlinear

Schroedinger equation expressed in terms of the spatio-temporal coordinates (x, t) , that corresponds to the “old” solution (8), one may naively just re-express (8) in terms of the new variables (that is, substitute in (8) the old variables (x', t') by their expressions in terms of the new variables (x, t)),

$$\begin{aligned}\Psi(x', t') &\rightarrow \Phi(x, t) = \Psi(x + vt, t) \\ &= \Phi_0 \exp_q [i(k(x + vt) - \omega t)], \quad \Phi_0 = \Psi_0.\end{aligned}\quad (10)$$

However, this procedure leads to a function of x and t that does not satisfy the nonlinear Schroedinger equation (as expressed in terms of the new variables (x, t)). In order to obtain a valid solution it is necessary to add an extra term to the argument of the q -exponential. Indeed, it can be verified after some algebra that

$$\frac{\Phi}{\Phi_0} = \left[1 - i(1 - q) \left\{ \omega t - k(x + vt) + \frac{1}{\hbar} \left(mvx + \frac{1}{2} mv^2 t \right) \right\} \right]^{\frac{1}{1-q}} \quad (11)$$

does satisfy the nonlinear Schroedinger equation. The extra term $\frac{1}{\hbar} (mvx + \frac{1}{2} mv^2 t)$ admits a clear physical interpretation. If we recast (11) under the guise,

$$\frac{\Phi}{\Phi_0} = \left[1 - i(1 - q) \left\{ \left(\omega - kv + \frac{mv^2}{2\hbar} \right) t - \left(k - \frac{mv}{\hbar} \right) x \right\} \right]^{\frac{1}{1-q}}, \quad (12)$$

it is plain that (12) has the form of a q -plane wave with frequency $\tilde{\omega}$ and wave number \tilde{k} , respectively, given by

$$\tilde{\omega} = \omega - kv + \frac{mv^2}{2\hbar}; \quad \tilde{k} = k - \frac{mv}{\hbar}. \quad (13)$$

Now, as shown in Ref. 5, the q -plane wave solutions to the nonlinear Schroedinger equation are compatible with the Planck and de Broglie relations connecting, respectively, frequency and wave number with energy and momentum: $E = \hbar\omega$ and $p = \hbar k$. Combining equations (13) with the Planck and de Broglie relations we obtain $\tilde{E} = E - pv + \frac{mv^2}{2}$ and $\tilde{p} = p - mv$, which are the correct Galilean transformations for the kinetic energy and momentum of a particle of mass m obeying the (nonrelativistic) energy-momentum relation $E = p^2/2m$. These considerations reinforce the validity of the Planck and de Broglie relations for the q -plane wave solutions of Eq. (1).

Taking the limit $q \rightarrow 1$ of the transformed solution (11), one sees that the relation between the original solution $\Psi(x', t')$ and the transformed one $\Phi(x, t)$ becomes,

$$\begin{aligned}\Psi(x', t') &\rightarrow \Phi(x, t) \\ &= \exp \left[-\frac{i}{\hbar} \left(\frac{mv^2}{2} t + mvx \right) \right] \Psi(x + vt, t),\end{aligned}\quad (14)$$

thus recovering the transformation rule corresponding to the linear Schroedinger equation.²³

Let us now consider a uniformly accelerated reference frame. The corresponding spatio-temporal coordinates (x, t) are

$$t = t'; \quad x = x' - \frac{1}{2} at'^2 = x' - \frac{1}{2} \frac{F}{m} t'^2, \quad (15)$$

where (x', t') are the variables associated with an inertial frame, a is the constant acceleration of reference frame (x, t) , and $a = \frac{F}{m}$. As in the previous discussion, we assume that the nonlinear Schroedinger equation (7) holds in the inertial frame (x', t') , and also that in this frame our system is described by the q -plane wave solution (8). Again, simply re-writing the q -plane wave solution (8) in terms of the new variables (x, t) does not yield a solution of the nonlinear Schroedinger equation. As in the above Galilean transformation case, new terms are needed in the argument of the q -exponential to obtain a valid solution. Let us consider the ansatz,

$$\frac{\Phi}{\Phi_0} = \left[1 - i(1 - q) \left\{ \omega t - k \left(x + \frac{Ft^2}{2m} \right) + \frac{F}{\hbar} \left(xt + \frac{Ft^3}{6m} \right) \right\} \right]^{\frac{1}{1-q}}. \quad (16)$$

Inserting (16) in the right- and the left-hand sides of the nonlinear Schroedinger equation yields

$$i\hbar \frac{\partial}{\partial t} \left(\frac{\Phi}{\Phi_0} \right) = \left[\hbar\omega - \frac{\hbar k F t}{m} + Fx + \frac{F^2 t^2}{2m} \right] \left(\frac{\Phi}{\Phi_0} \right)^q \quad (17)$$

and

$$-\frac{1}{2-q} \frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} \left[\left(\frac{\Phi}{\Phi_0} \right)^{2-q} \right] = \left[\frac{\hbar^2 k^2}{2m} - \frac{\hbar k F t}{m} + \frac{F^2 t^2}{2m} \right] \left(\frac{\Phi}{\Phi_0} \right)^q. \quad (18)$$

Comparing (17) with (18) one verifies that the ansatz (16) satisfies the nonlinear equation,

$$i\hbar \frac{\partial}{\partial t} \left(\frac{\Phi}{\Phi_0} \right) = -\frac{1}{2-q} \frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} \left[\left(\frac{\Phi}{\Phi_0} \right)^{2-q} \right] + V(x) \left(\frac{\Phi}{\Phi_0} \right)^q, \quad (19)$$

where $V(x) = Fx$. The nonlinear equation (19) can be interpreted as describing the motion of a particle of mass m under a constant force $-F$ (with the associated potential function $V = Fx$). This is consistent with the well-known fact that the behavior of a free particle with respect to a uniformly accelerated reference frame is equivalent to the behaviour of a particle in an inertial reference frame moving under the effect of a constant force. In the limit $F \rightarrow 0$, Eq. (19) reduces to the nonlinear Schroedinger equation for a free particle introduced in Ref. 5, and solution (16) reduces to the corresponding q -plane wave solution. Also, $q \rightarrow 1$ in Eq. (19) corresponds to the standard linear Schroedinger equation for a particle of mass m moving under a constant force $-F$. An interesting feature of Eq. (19) is that the potential V couples to Φ^q , instead of coupling to Φ , as happens in the standard linear case ($q = 1$). Consistently, the q -plane wave $\Phi(x, t) = \Phi_0 \exp_q[i(kx - \omega t)]$ is not only a solution of the free-particle nonlinear Schroedinger equation (when $\hbar\omega = \frac{\hbar^2 k^2}{2m}$), but also of the nonlinear equation

$$i\hbar \frac{\partial}{\partial t} \left(\frac{\Phi}{\Phi_0} \right) = -\frac{1}{2-q} \frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} \left[\left(\frac{\Phi}{\Phi_0} \right)^{2-q} \right] + V_0 \left(\frac{\Phi}{\Phi_0} \right)^q, \quad (20)$$

with a constant potential V_0 , provided that $\hbar\omega = \frac{\hbar^2 k^2}{2m} + V_0$, which, using the Planck and de Broglie relations, becomes $E = \frac{p^2}{2m} + V_0$, as expected.

Considering now the limit $q \rightarrow 1$ of the transformed solution (16), we verify that the original solution $\Psi(x', t')$ and the transformed one $\Phi(x, t)$ are linked through

$$\begin{aligned} \Psi(x', t') &\rightarrow \Phi(x, t) \\ &= \exp \left[-\frac{i}{\hbar} \left(Fxt + \frac{F^2 t^3}{6m} \right) \right] \Psi \left(x + \frac{Ft^2}{2m}, t \right), \end{aligned} \quad (21)$$

thus recovering the transformation rule associated with the linear Schroedinger equation.²³ Other nonlinear equations previously studied in the literature also behave like the standard linear Schroedinger equation, in the sense that they “pick up” phases (like in (21)) when accelerated (or under Galilean transformations). In contrast, in the case of the evolution equation studied here the extra terms appearing in the transformed solutions, strictly speaking, are not “phases” because they cannot (except in the limit $q \rightarrow 1$) be written as a (complex) multiplicative factor of modulus 1 (as happens in (21)). These extra terms are, however, of special relevance in connection with the Planck and de Broglie relations discussed in the present work.

We have investigated the effect of uniform acceleration on q -plane waves, soliton-like solutions of a recently proposed nonlinear Schroedinger equation associated with non-extensive thermostatics. We first studied the behaviour of these solutions under Galilean transformations relating different inertial frames and obtained the transformation rule satisfied by the q -plane waves. This rule turns out to be fully consistent with the de Broglie and Planck relations, thus providing further support to the validity of these relations for the q -plane wave solutions. Then we derived the transformation law yielding new solutions corresponding to the aforementioned q -plane waves when “viewed” from uniformly accelerated frames. In the limit $q \rightarrow 1$ the transformation laws advanced here reduce to those associated with time dependent solutions of the standard, linear Schroedinger equation. The

accelerated q -plane wave solutions investigated here admit two possible interpretations: they can be viewed as describing a free particle as “seen” from a uniformly accelerated frame or, alternatively, as describing a particle moving under the effect of a constant force. In fact, the nonlinear Schrödinger equation governing these solutions (when expressed in terms of the accelerated frame’s coordinates) incorporates a new term involving a potential function $V(x)$. This equation indicates that, within the present generalization of Schrödinger equation, the potential $V(x)$ “couples” to a power of the wave function, Φ^q , instead of coupling just to the wave function Φ , as happens with the linear Schrödinger equation. This result opens the door for the study of a variety of basic potentials such as, among others, potential steps, square wells and barriers, and harmonic potentials which, although simple, may provide valuable insights concerning possible physical applications.

Partial financial support from CNPq and FAPERJ (Brazilian agencies), and from the Projects FQM-2445 and FQM-207 of the Junta de Andalucía is acknowledged.

- ¹ C. Sulem and P.-L. Sulem, *The Nonlinear Schrödinger Equation: Self-Focusing and Wave Collapse* (Springer, New York, 1999).
- ² T. D. Frank, *Nonlinear Fokker-Planck Equations: Fundamentals and Applications* (Springer, Berlin, 2005).
- ³ A. C. Scott, *The Nonlinear Universe* (Springer, Berlin, 2007).
- ⁴ A. D. Polyaniin and V. F. Zaitsev, *Handbook of Nonlinear Partial Differential Equations* (Chapman and Hall/CRC, Boca Raton, 2004).
- ⁵ F. D. Nobre, M. A. Rego-Monteiro, and C. Tsallis, “Nonlinear relativistic and quantum equations with a common type of solution,” *Phys. Rev. Lett.* **106**, 140601 (2011).
- ⁶ F. D. Nobre, M. A. Rego-Monteiro, and C. Tsallis, “A generalized nonlinear Schrödinger equation: Classical field-theoretic approach,” *EPL* **97**, 41001 (2012).
- ⁷ J. S. Andrade, Jr., G. F. T. da Silva, A. A. Moreira, F. D. Nobre, and E. M. F. Curado, “Thermostatistics of overdamped motion of interacting particles,” *Phys. Rev. Lett.* **105**, 260601 (2010); Y. Levin and R. Pakter, “Comment on: Thermostatistics of overdamped motion of interacting particles,” *Phys. Rev. Lett.* **107**, 088901 (2011); J. S. Andrade, Jr., G. F. T. da Silva, A. A. Moreira, F. D. Nobre, and E. M. F. Curado, “Andrade *et al.* reply,” *Phys. Rev. Lett.* **107**, 088902 (2011).
- ⁸ M. S. Ribeiro, F. D. Nobre, and E. M. F. Curado, “Time evolution of interacting vortices under overdamped motion,” *Phys. Rev. E* **85**, 021146 (2012).
- ⁹ C. Tsallis, “Possible generalization of Boltzmann-Gibbs statistics,” *J. Stat. Phys.* **52**, 479 (1988).
- ¹⁰ C. Tsallis, *Introduction to Nonextensive Statistical Mechanics* (Springer, New York, 2009).
- ¹¹ C. Beck, “Generalised information and entropy measures in physics,” *Contemp. Phys.* **50**, 495 (2009).
- ¹² A. Carati, S. L. Cacciatori, and L. Galgani, “Discrete matter, far fields, and dark matter,” *EPL* **83**, 59002 (2008).
- ¹³ S. Umarov, C. Tsallis, and S. Steinberg, *Milan J. Math.* **76**, 307 (2008); S. Umarov, C. Tsallis, M. Gell-Mann, and S. Steinberg, “Generalization of symmetric α -stable Lévy distributions for $q > 1$,” *J. Math. Phys.* **51**, 033502 (2010).
- ¹⁴ A. R. Plastino and A. Plastino, “Non-extensive statistical mechanics and generalized Fokker-Planck equation,” *Physica A* **222**, 347 (1995); C. Tsallis and D. J. Bukman, “Anomalous diffusion in the presence of external forces: Exact time-dependent solutions and their thermostatical basis,” *Phys. Rev. E* **54**, R2197 (1996).
- ¹⁵ P. Douglas, S. Bergamini, and F. Renzoni, “Tunable Tsallis distributions in dissipative optical lattices,” *Phys. Rev. Lett.* **96**, 110601 (2006).
- ¹⁶ B. Liu and J. Goree, “Superdiffusion and non-Gaussian statistics in a driven-dissipative 2D dusty plasma,” *Phys. Rev. Lett.* **100**, 055003 (2008).
- ¹⁷ R. G. DeVoe, “Power-law distributions for a trapped ion interacting with a classical buffer gas,” *Phys. Rev. Lett.* **102**, 063001 (2009).
- ¹⁸ R. M. Pickup, R. Cywinski, C. Pappas, B. Farago, and P. Fouquet, “Generalized spin-glass relaxation,” *Phys. Rev. Lett.* **102**, 097202 (2009).
- ¹⁹ V. Khachatryan *et al.* (CMS Collaboration), “Transverse-momentum and pseudorapidity distributions of charged hadrons in pp collisions at $\sqrt{s} = 7$ TeV,” *Phys. Rev. Lett.* **105**, 022002 (2010).
- ²⁰ E. P. Borges, “On a generalization of circular and hyperbolic functions,” *J. Phys. A* **31**, 5281 (1998).
- ²¹ M. Jauregui and C. Tsallis, “New representations of π and Dirac delta using the nonextensive-statistical-mechanics q -exponential function,” *J. Math. Phys.* **51**, 063304 (2010); A. Chevreuil, A. Plastino, and C. Vignat, “On a conjecture about Dirac’s delta representation using q -exponentials,” *J. Math. Phys.* **51**, 093502 (2010).
- ²² G. Kaniadakis, A. Lavagno, and P. Quarati, “Wigner equation for particles obeying an exclusion-inclusion principle,” *Phys. Rev. E* **57**, 1395 (1998).
- ²³ A. Peres, *Quantum Theory: Concepts and Methods* (Kluwer, Dordrecht, 1993).