

THERMODYNAMICS AND STATISTICAL MECHANICS
FOR COMPLEX SYSTEMS
– FOUNDATIONS AND APPLICATIONS*

CONSTANTINO TSALLIS

Centro Brasileiro de Pesquisas Físicas
and National Institute of Science and Technology for Complex Systems
Rua Xavier Sigaud 150, 22290-180 Rio de Janeiro, Brazil
and

Santa Fe Institute, 1399 Hyde Park Road, Santa Fe, New Mexico 87501, USA

The complexity (or simplicity) of a system can be characterized through very many paths. A quite interesting and operational one, which we follow here, consists in making use of entropic concepts. The thermodynamical and statistical-mechanical foundations of this approach, as well as its (analytical, computational, observational, and experimental) applications for natural, artificial and social systems will be briefly reviewed.

1. Introduction

In the 1860s Clausius introduced [1] the concept of entropy (noted S) within thermodynamics. This most relevant step basically concluded the foundations of this science, and enabled its expression in terms of the so called principles of thermodynamics (nowadays referred to as zeroth, first, second, third principles). One decade later Boltzmann postulated [2, 3] an entropic functional (noted S_{BG} where BG stands for Boltzmann-Gibbs, in order to also incorporate the refinements introduced by Gibbs [4]), which connects the macroscopic entropy introduced by Clausius with the microscopic world. The entropy S_{BG} is essentially consistent with the entire structure of classical thermodynamics, and constitutes the basis of the theory (very important in contemporary physics) referred to as statistical mechanics. During nearly 140 years, BG statistical mechanics keeps providing uncountable excellent predictions for many kinds of thermal and similar phenomena. This has somehow sedimented – wrongly! – in the mind of

* Invited review presented at the XXVIIth Marian Smoluchowski Symposium on Statistical Physics, held in Zakopane, Poland, in 22-26 September 2014.

many physicists that no other statistical-mechanical theory is possible, and that S_{BG} is not only sufficient but also necessary for thermodynamics to be satisfied. The scenario has gradually and substantially evolved during the last 25 years, especially through the proposal of nonadditive entropies as a basis for possibly generalizing BG statistical mechanics [5].

The whole process exhibits a strong analogy with the history of mechanics. At the end of the XIXth century, mechanics was a synonym of Newtonian or classical mechanics. But soon after, quantum and relativistic mechanics were formulated, and there are now several theories within mechanics, which contain Newtonian mechanics as particular case. In the present review we argue that more than one theory exists within statistical mechanics, which nevertheless remain related to thermodynamics, the BG statistical mechanics playing of course the basic reference role.

2. Nonadditive entropies and nonextensive statistical mechanics

2.1. Nonadditive entropies

For a system constituted by N elements that globally admits W microscopic configurations (*complexions* in the language of Boltzmann's time), we have the BG entropy given by

$$S_{BG}(N) \equiv -k \sum_{i=1}^{W(N)} p_i \ln p_i \quad \left(\sum_{i=1}^{W(N)} p_i = 1 \right), \quad (1)$$

where k is a conventional constant (typically Boltzmann constant k_B), and $\{p_i\}$ are the probabilities. When all probabilities are equal, i.e., $p_i = 1/W(N)$, we recover the celebrated Boltzmann formula

$$S_{BG}(N) = k \ln W(N). \quad (2)$$

If the system is such that, in the $N \rightarrow \infty$ limit,

$$W(N) \propto \mu^N, \quad (3)$$

where $\mu > 1$ is a constant, then Eq. (2) implies $S_{BG}(N) \propto N$, which provides an *extensive* entropy (i.e., $0 < \lim_{N \rightarrow \infty} S(N)/N < \infty$), in conformity with thermodynamics. For example, if we have N independent two-faced fair coins, we have that $W(N) = 2^N$, hence $S_{BG} = kN \ln 2$.

But if the elements of the system are strongly correlated, it frequently happens that the number of admissible microscopic configurations decreases drastically, and it might essentially become, in the $N \rightarrow \infty$ limit, a zero-Lebesgue measure set. For example, a wide class of systems yield

$$W(N) \propto N^\rho \quad (\rho > 0), \quad (4)$$

hence $S_{BG}(N) \propto \ln N$, thus violating the thermodynamic extensivity of the entropy. A natural way out appears to be to define *nonadditive* entropies such as [5]

$$\begin{aligned} S_q(N) &\equiv k \frac{1 - \sum_{i=1}^{W(N)} p_i^q}{q-1} = k \sum_{i=1}^{W(N)} p_i \ln_q \frac{1}{p_i} = -k \sum_{i=1}^{W(N)} p_i^q \ln_q p_i \\ &= -k \sum_{i=1}^{W(N)} p_i \ln_{2-q} p_i \quad \left(\sum_{i=1}^{W(N)} p_i = 1; q \in \mathcal{R}; S_1 = S_{BG} \right), \end{aligned} \quad (5)$$

where the q -logarithmic function is defined as $\ln_q z \equiv \frac{z^{1-q} - 1}{1-q}$ ($z > 0$; $\ln_1 z = \ln z$). If all admissible events are equally probable, i.e., $p_1 = 1/W(N)$, we have

$$S_q(N) = k \ln_q W(N). \quad (6)$$

If $W(N)$ is given by Eq. (4), we straightforwardly verify that $S_{q=1-1/\rho}(N) \propto N$, thus satisfying the thermodynamic extensivity of the entropy as desirable.

It is easy to verify that, if A and B are two probabilistically independent systems, the entropy of the total system $A + B$ satisfies

$$\frac{S_q(A+B)}{k} = \frac{S_q(A)}{k} + \frac{S_q(B)}{k} + (1-q) \frac{S_q(A)}{k} \frac{S_q(B)}{k}. \quad (7)$$

In other words, by using Penrose's definition of entropic additivity [6], we have that the entropic functional S_q is *nonadditive* for all values of q excepting for $q = 1$, value for which it recovers the *additive* BG functional S_{BG} .

We can summarize these facts by saying that, if the system is such that $W(N)$ is given by Eq.(3) (which basically means independence or weak correlations), the additive entropy S_{BG} is extensive and the nonadditive entropy S_q is nonextensive for all $q \neq 1$, whereas, if $W(N)$ is given by Eq. (4) (which is a frequent class of strong correlations), S_q is extensive for $q = 1 - \frac{1}{\rho}$ and nonextensive for all other values of q , including $q = 1$. As we see, additivity only depends on the entropic functional and is easy to determine. Extensivity, in contrast, depends on both the entropic functional and the system, and can generically be quite hard to analytically determine.

Another interesting case occurs if

$$W(N) \propto \nu^{N^\gamma} \quad (\nu > 1; 0 < \gamma < 1), \quad (8)$$

which also essentially corresponds to zero Lebesgue measure for the occupancy of phase space (or of the analogous spaces for quantum systems). To

tackle this case, we may use the following entropy [8, 106]:

$$S_\delta(N) \equiv k \sum_{i=1}^{W(N)} p_i \left(\ln \frac{1}{p_i} \right)^\delta \quad \left(\sum_{i=1}^{W(N)} p_i = 1; S_1 = S_{BG} \right). \quad (9)$$

This entropy becomes, for equal probabilities, i.e., $p_i = 1/W(N)$,

$$S_\delta(N) = k(\ln W)^\delta. \quad (10)$$

We can verify that $S_{\delta=1/\gamma}(N) \propto N$, in conformity with thermodynamics. Also, if A and B are probabilistically independent, we have that, for $\delta \neq 1$,

$$S_\delta(A+B) \neq S_q(A) + S_q(B), \quad (11)$$

hence S_δ is nonadditive.

It is worthy to emphasize that $\mu^N \gg \nu^{N^\delta} \gg N^\rho$, which exhibits that both nonexponential cases (4) (power law) and (8) (stretched exponential) correspond essentially to zero Lebesgue measure problems in what concerns the occupancy of phase space. Let us finally mention that all three entropies that have been discussed here can be unified through

$$S_{q,\delta}(N) \equiv k \sum_{i=1}^{W(N)} p_i \left(\ln_q \frac{1}{p_i} \right)^\delta \quad \left(\sum_{i=1}^{W(N)} p_i = 1; S_{q,1} = S_q; S_{1,\delta} = S_\delta; S_{1,1} = S_{BG} \right). \quad (12)$$

A summary of the present Subsection is presented in Table 1.

2.2. Extensive entropies

Let us answer here an interesting question: *Since we admit entropic functionals that might be nonadditive, why do we not admit nonextensive entropies as well?* (This question was first pointed to me in these terms by A. Coniglio).

To discuss thermodynamics and its Legendre-transform structure on general grounds, let us remind a typical form of the thermodynamical energy G (Gibbs energy) of a generic d -dimensional system [9]:

$$G(V, T, p, \mu, H, \dots) = U(V, T, p, \mu, H, \dots) - TS(V, T, p, \mu, H, \dots) \quad (13)$$

$$+ pV - \mu N(V, T, p, \mu, H, \dots) - HM(V, T, p, \mu, H, \dots) - \dots, \quad (14)$$

where T, p, μ, H are the temperature, pressure, chemical potential, external magnetic field respectively, and U, S, V, N, M are the internal energy,

Table 1. Entropic functionals and classes of systems (*exponential*, *power-law* and *stretched-exponential*, see text) for which the entropy is extensive, *i.e.*, proportional to the number N of elements. $W(N)$ is the number of admissible microscopic configurations of a system with N elements (only configurations with nonvanishing occurrence probability are considered admissible). We also see the specific values of q and δ for which respectively S_q and S_δ are extensive. From [10].

	ENTROPY		
$W(N)$ ($N \rightarrow \infty$)	S_{BG} (ADDITIVE)	S_q ($q \neq 1$) (NONADDITIVE)	S_δ ($\delta \neq 1$) (NONADDITIVE)
$\sim \mu^N$ ($\mu > 1$)	EXTENSIVE	NONEXTENSIVE	NONEXTENSIVE
$\sim N^\rho$ ($\rho > 0$)	NONEXTENSIVE	EXTENSIVE ($q = 1 - 1/\rho$)	NONEXTENSIVE
$\sim \nu^{N^\gamma}$ ($\nu > 1$; $0 < \gamma < 1$)	NONEXTENSIVE	NONEXTENSIVE	EXTENSIVE ($\delta = 1/\gamma$)

entropy, volume, number of particles (in turn proportional to the number of degrees of freedom), magnetization respectively. From the Legendre structure we identify three classes of variables, namely (i) those that are expected to *always be extensive* like N itself (S, V, N, M, \dots), *i.e.*, scaling with the (d -dimensional) volume $V = L^d \propto N$, where L is a characteristic linear dimension of the system (clearly, $V \propto A^{d/(d-1)}$, where A is the d -dimensional area)¹, (ii) those that characterize the external conditions under which the system is placed (T, p, μ, H, \dots), scaling with L^θ , and (iii) those that represent energies (G, U), scaling with L^ϵ .

It trivially follows

$$\epsilon = \theta + d. \quad (15)$$

If we divide Eq. (14) by $L^{\theta+d}$ and consider the large L limit (*i.e.*, the

¹ Within the thermodynamical Legendre-transform structure, it is of course natural that S, V, N, M belong to the same class. The variable N is *extensive* by definition. Therefore, so must clearly also be the entropy S , as well as V, M .

thermodynamical limit), we obtain

$$g\left(\frac{T}{L^\theta}, \frac{p}{L^\theta}, \frac{\mu}{L^\theta}, \frac{H}{L^\theta}, \dots\right) = u\left(\frac{T}{L^\theta}, \frac{p}{L^\theta}, \frac{\mu}{L^\theta}, \frac{H}{L^\theta}, \dots\right) - \frac{T}{L^\theta} s\left(\frac{T}{L^\theta}, \frac{p}{L^\theta}, \frac{\mu}{L^\theta}, \frac{H}{L^\theta}, \dots\right) \\ + \frac{p}{L^\theta} - \frac{\mu}{L^\theta} n\left(\frac{T}{L^\theta}, \frac{p}{L^\theta}, \frac{\mu}{L^\theta}, \frac{H}{L^\theta}, \dots\right) - \frac{H}{L^\theta} m\left(\frac{T}{L^\theta}, \frac{p}{L^\theta}, \frac{\mu}{L^\theta}, \frac{H}{L^\theta}, \dots\right) - \dots \quad (16)$$

where $g \equiv \lim_{L \rightarrow \infty} G/L^{\theta+d}$, $u \equiv \lim_{L \rightarrow \infty} U/L^{\theta+d}$, $s \equiv \lim_{L \rightarrow \infty} S/L^d$, $n \equiv \lim_{L \rightarrow \infty} N/L^d$, $m \equiv \lim_{L \rightarrow \infty} M/L^d$. The three types of thermodynamical scalings that have emerged are represented in Fig. 1. Physical illustrations, including for black holes, of these various behaviours can be found in [11].

The correctness of these scalings has been repeatedly checked in the literature for many types of fluid, magnetic, percolating, polymerized systems [12–23]: details can be found in those papers.

Besides the above general Legendre structure which mandates the thermodynamical entropy to be extensive in all cases, there exist other consistent arguments as well, which come from large deviation theory considerations (see [24] and references therein). But the presentation here of those arguments remains out of the present scope.

2.3. Nonextensive statistical mechanics

The optimization of nonextensive entropies such as S_q with appropriate constraints [25, 26] yields what is referred in the literature as *nonextensive statistical mechanics*. The word *nonextensive* comes here from the (internal) energy $U(N)$, which typically indeed is nonextensive (typically $\lim_{N \rightarrow \infty} U(N)/N$ diverges), *not* from the entropy, which is not necessarily taken to be additive in order precisely to be always extensive.

In other words, in BG statistical mechanics, both the energy and the entropy are extensive; in nonextensive statistical mechanics, the energy is nonextensive (superextensive for long-range-interacting many-body Hamiltonian systems), but the entropy remains extensive. The entropic extensivity within BG thermostatistics is guaranteed by the additive BG entropic functional; the entropic extensivity within nonextensive thermostatistics is guaranteed by nonadditive entropic functionals such as S_q and S_δ . Let us mention at this point a regrettable misname that has been source of not few confusions in the literature: the title of the book [27] should have been *Nonadditive Entropy*, and not *Nonextensive Entropy*.

If the linear mean value of a random variable x is finite, the optimization of $S_q[p(x)]$ yields the q -exponential function $p(x) \propto e_q^{-\beta x}$, where β is related to the Lagrange parameter, and the function e_q^z is defined as follows:

$$e_q^z \equiv [1 + (1 - q)z]^{\frac{1}{1-q}}, \quad (17)$$




Fig. 1. Representation of the different scaling regimes of the Eq. (16) for classical d -dimensional systems. For attractive long-range interactions (*i.e.*, $0 \leq \alpha/d \leq 1$, α characterizes the interaction range in a potential with the form $1/r^\alpha$) we may distinguish three classes of thermodynamic variables, namely, those scaling with L^θ , named *pseudo-intensive* (L is a characteristic linear length, θ is a system-dependent parameter), those scaling with $L^{d+\theta}$, the *pseudo-extensive* ones (the energies), and those scaling with L^d (which are always extensive). For short-range interactions (*i.e.*, $\alpha > d$) we have $\theta = 0$ and the energies recover their standard L^d extensive scaling, falling in the same class of S , N , V , etc, whereas the previous pseudo-intensive variables become truly intensive ones (independent of L); this is the region, with two classes of variables, that is covered by the traditional textbooks of thermodynamics. From [10].

if $1 + (1 - q)z > 0$, and zero otherwise. This function is the inverse of $\ln_q z$ previously defined; also, $e_1^z = e^z$.

If the linear mean value vanishes and the quadratic mean value is nonvanishing, the optimization of $S_q[(p(x))]$ yields $p(x) \propto e_q^{-\beta x^2}$, currently referred to in the literature as q -Gaussian.

In what concerns the thermostatics corresponding to S_δ and $S_{q,\delta}$, the basic results are already available, in [28] and [29] respectively.

To end this Section let us remind that the indices q and δ are to be determined from first principles, *i.e.*, essentially from mechanics. One illustrative such calculation can be found in [30, 31]: see Fig. 2.




Fig. 2. The index q has been determined [30] from first principles, namely from the universality class of the Hamiltonian, characterized by the central charge c . The values $c = 1/2$ and $c = 1$ respectively correspond to the Ising and XY ferromagnetic chains in the presence of transverse field at $T = 0$ criticality. For other models see [32, 33]. In the $c \rightarrow \infty$ limit we recover the Boltzmann-Gibbs (BG) value, i.e., $q = 1$. For arbitrary value of c , the subsystem *nonadditive* entropy S_q is thermodynamically *extensive* for, and only for, $q = \frac{\sqrt{9+c^2}-3}{c}$.

3. Applications

Nonadditive entropies and nonextensive statistical mechanics have been relevant to a wide number of predictions, verifications and applications in natural, artificial and social systems [34]. We briefly mention here some of them.

There have been various applications to long-range-interacting many-body classical Hamiltonians such as the XY ferromagnet and the Fermi-Pasta-Ulam models ([35, 36] and references therein), self-organized criticality [37], turbulence [38, 39], low-dimensional dissipative maps [40–45], low-dimensional conservative maps [46, 47], anomalous diffusion [48, 49], the Heisenberg model with long-range interactions [50, 51] (by the way the quantum analog with long-range interactions remains an open problem; for the short-range model see, for instance, [52] and references therein), spin glass relaxation [53], cold atoms in optical lattices [54, 55], ion trapping [56], overdamped motion of vortices in type-II superconductors [57–61], dusty plasma [62], the so called q -triplet in solar wind and similar phenomena [63–71], asymptotically scale-free networks [72], black-hole physics [11, 73–75], non-




Fig. 3. Experimental distributions of the transverse moments in hadronic jets at the CMS, ALICE and ATLAS detectors at LHC. The data are from [103]. They can be remarkably well fitted (along fourteen decades) with the q -exponential function $e_q^x \equiv [1 + (1 - q)x]^{1/(1-q)}$, which, under appropriate constraints, extremizes the entropy S_q . See details in [104].

linear relativistic and quantum equations [76]. A special mention is deserved by the q -generalization of central limit theorems and related properties [77–80].

Let us specifically add here recent applications in high-energy physics at the LHC/CERN and elsewhere [81–102]. A typical illustration is presented in [31, 103, 104]: see for instance Fig. 3 (some tiny log-periodic oscillations are observed on top of the q -exponentials, which might be related to some microscopic fractality [105]).

Many other applications (in economics, linguistics, geophysics, neurosciences) can be found in [27, 106].

4. Final remarks

The logical path we have followed here is to satisfy in all cases the extensivity of the entropy, since this property is mandated by classical thermodynamics. For standard systems (essentially ergodic, which implies an occupation of phase space with a nonzero Lebesgue measure) this requirement

is satisfied by the Boltzmann-Gibbs entropic functional, which is additive. For nonstandard systems (essentially nonergodic, which typically emerges through a zero-Lebesgue-measure occupation of phase space) this requirement is, for wide classes of systems, satisfied for nonadditive entropic functionals such as S_q and S_δ . The indices q and δ are to be determined from first principles, namely from mechanics (classical, quantum, relativistic). When this is analytically intractable, they may be determined from mesoscopic arguments (e.g., from Langevin-like and Fokker-Planck-like equations), or simply through fitting. The extremization of these nonadditive entropies under appropriate constraints yields a generalization of Boltzmann-Gibbs statistical mechanics, currently referred to as nonextensive statistical mechanics. The amazing number of phenomena in natural, artificial and social systems which do follow (not rarely through many experimental or numerical decades) q -exponential and q -Gaussian behaviours together with their analytical connections provide, as far as we can see, a neat validation of these concepts. Various interesting questions remain open, such as – to name just one – the analytical connection between the indices q and the exponent characterizing the range of long-range-interacting many-body Hamiltonians.

Acknowledgments

Partial financial support from the Brazilian agencies CNPq and Faperj and from the John Templeton Foundation is acknowledged.

REFERENCES

- [1] R. Clausius, *Über verschiedene für die Anwendung bequeme Formen der Hauptgleichungen der mechanischen Wärmetheorie*, Annalen der Physik **125**, 353 (1865).
- [2] L. Boltzmann, *Weitere Studien über das Wärmegleichgewicht unter Gas molekülen* [*Further Studies on Thermal Equilibrium Between Gas Molecules*], Wien, Ber. **66**, 275 (1872).
- [3] L. Boltzmann, *Über die Beziehung eines allgemeine mechanischen Satzes zum zweiten Hauptsatz der Wärmetheorie*, Sitzungsberichte, K. Akademie der Wissenschaften in Wien, Math.-Naturwissenschaften **75**, 67-73 (1877); English translation (*On the Relation of a General Mechanical Theorem to the Second Law of Thermodynamics*) in S. Brush, *Kinetic Theory*, Vol. **2: Irreversible Processes**, 188-193 (Pergamon Press, Oxford, 1966).
- [4] J.W. Gibbs, *Elementary Principles in Statistical Mechanics – Developed with Especial Reference to the Rational Foundation of Thermodynamics* (C. Scribner's Sons, New York, 1902; Yale University Press, New Haven, 1948); OX Bow Press, Woodbridge, Connecticut, 1981) (page 35).

- [5] C. Tsallis, *Possible generalization of Boltzmann-Gibbs statistics*, J. Stat. Phys. **52**, 479 (1988) [First appeared as preprint in 1987: CBPF-NF-062/87, ISSN 0029-3865, Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro].
- [6] O. Penrose, *Foundations of Statistical Mechanics: A Deductive Treatment* (Pergamon, Oxford, 1970), p. 167.
- [7] C. Tsallis, *Introduction to Nonextensive Statistical Mechanics - Approaching a Complex World* (Springer, New York, 2009).
- [8] M.R. Ubriaco, *Entropies based on fractional calculus*, Phys. Lett. A **373**, 2516 (2009).
- [9] H.B. Callen, *Thermodynamics and an Introduction to Thermostatistics*, 2nd Edition (John Wiley, 1985).
- [10] C. Tsallis and L.J.L. Cirto, *Thermodynamics is more powerful than the role to it reserved by Boltzmann-Gibbs statistical mechanics*, Eur. Phys. J. Special Topics **223**, 2161 (2014).
- [11] C. Tsallis and L.J.L. Cirto, *Black hole thermodynamical entropy*, Eur. Phys. J. C **73**, 2487 (2013).
- [12] P. Jund, S.G. Kim and C. Tsallis, *Crossover from extensive to nonextensive behavior driven by long-range interactions*, Phys. Rev. B **52**, 50 (1995).
- [13] J.R. Grigera, *Extensive and non-extensive thermodynamics. A molecular dynamic test*, Phys. Lett. A **217**, 47 (1996).
- [14] S.A. Cannas and F.A. Tamarit, *Long-range interactions and non-extensivity in ferromagnetic spin systems*, Phys. Rev. B **54**, R12661 (1996).
- [15] L.C. Sampaio, M.P. de Albuquerque and F.S. de Menezes, *Nonextensivity and Tsallis statistics in magnetic systems*, Phys. Rev. B **55**, 5611 (1997).
- [16] C. Anteneodo and C. Tsallis, Phys. Rev. Lett. **80**, 5313 (1998).
- [17] Curilef and C. Tsallis, *Critical temperature and nonextensivity in long-range-interacting Lennard-Jones-like fluids*, Phys. Lett. A **264**, 270 (1999).
- [18] R.F.S. Andrade and S.T.R. Pinho, *Tsallis scaling and the long-range Ising chain: A transfer matrix approach*, Phys. Rev. E **71**, 026126 (2005).
- [19] C.A. Condat, J. Rangel and P.W. Lamberti, *Anomalous diffusion in the nonasymptotic regime*, Phys. Rev. E **65**, 026138 (2002).
- [20] H.H.A. Rego, L.S. Lucena, L.R. da Silva and C. Tsallis, *Crossover from extensive to nonextensive behavior driven by long-range $d = 1$ bond percolation*, Physica A **266**, 42 (1999).
- [21] U.L. Fulco, L.R. da Silva, F.D. Nobre, H.H.A. Rego and L.S. Lucena, *Effects of site dilution on the one-dimensional long-range bond-percolation problem*, Phys. Lett. A **312**, 331 (2003).
- [22] L. Borland and J.G. Menchero, *Nonextensive effects in tight-binding systems with long-range hopping*, in *Nonextensive Statistical Mechanics and Thermodynamics*, eds. S.R.A. Salinas and C. Tsallis, Braz. J. Phys. **29**, 169 (1999).
- [23] L. Borland, J.G. Menchero and C. Tsallis, *Anomalous diffusion and nonextensive scaling in a one-dimensional quantum model with long-range interactions*, Phys. Rev. B **61**, 1650 (2000).

- [24] G. Ruiz and C. Tsallis, *Reply to Comment on "Towards a large deviation theory for strongly correlated systems"*, Phys. Lett. A **377**, 491-495 (2013).
- [25] E.M.F. Curado and C. Tsallis, *Generalized statistical mechanics: connection with thermodynamics*, J. Phys. A **24**, L69 (1991); Corrigenda: **24**, 3187 (1991) and **25**, 1019 (1992).
- [26] C. Tsallis, R.S. Mendes and A.R. Plastino, *The role of constraints within generalized nonextensive statistics*, Physica A **261**, 534 (1998).
- [27] M. Gell-Mann and C. Tsallis, *Nonextensive Entropy - Interdisciplinary Applications*, (Oxford University Press, New York, 2004).
- [28] M.S. Ribeiro, C. Tsallis and F.D. Nobre, *Probability distributions extremizing the nonadditive entropy S_δ and stationary states of the corresponding nonlinear Fokker-Planck equation*, Phys. Rev. E **88**, 052107 (2013).
- [29] M.S. Ribeiro, F.D. Nobre and C. Tsallis, *Probability distributions and associated nonlinear Fokker-Planck equation for the two-index entropic form $S_{q,\delta}$* , Phys. Rev. E **89**, 052135 (2014).
- [30] F. Caruso and C. Tsallis, *Nonadditive entropy reconciles the area law in quantum systems with classical thermodynamics*, Phys. Rev. E **78**, 021102 (2008).
- [31] C. Tsallis and H.J. Haubold, *Boltzmann-Gibbs entropy is sufficient but not necessary for the likelihood factorization required by Einstein*, 1407.6052 [cond-mat.stat-mech] (2014).
- [32] F.C. Alcaraz, *The critical behaviour of self-dual $Z(N)$ spin systems: finite-size scaling and conformal invariance*, J. Phys. A: Math. Gen. **20**, 2511 (1987).
- [33] F.C. Alcaraz and M.J. Martins, *The operator content of the exactly integrable $SU(N)$ magnets*, J. Phys. A: Math. Gen. **23**, L1079 (1990).
- [34] A regularly updated Bibliography can be found at <http://tsallis.cat.cbpf.br/biblio.htm>
- [35] L.J.L. Cirto, V.R.V. Assis and C. Tsallis, *Influence of the interaction range on the thermostatics of a classical many-body system*, Physica A **393**, 286 (2014).
- [36] H. Christodoulidi, C. Tsallis and T. Bountis, *Fermi-Pasta-Ulam model with long-range interactions: Dynamics and thermostatics*, EPL **108**, 40006 (2014).
- [37] F.A. Tamarit, S.A. Cannas and C. Tsallis, *Sensitivity to initial conditions in the Bak-Sneppen model of biological evolution*, Eur. phys. J. B **1**, 545 (1998).
- [38] B.M. Boghosian, *Thermodynamic description of the relaxation of two-dimensional turbulence using Tsallis statistics*, Phys. Rev. E **53**, 4754 (1996).
- [39] C. Anteneodo and C. Tsallis, *Two-dimensional turbulence in pure-electron plasma: A nonextensive thermostatical description*, J. Molecular Liquids **71**, 255 (1997).
- [40] F. Baldovin and A. Robledo, *Sensitivity to initial conditions at bifurcations in one-dimensional nonlinear maps: Rigorous nonextensive solutions*, Europhys. Lett. **60**, 518 (2002).

- [41] F. Baldovin and A. Robledo, *Universal renormalization-group dynamics at the onset of chaos in logistic maps and nonextensive statistical mechanics*, Phys. Rev. E **66**, R045104 (2002).
- [42] F. Baldovin and A. Robledo, *Nonextensive Pesin identity. Exact renormalization group analytical results for the dynamics at the edge of chaos of the logistic map*, Phys. Rev. E **69**, 045202(R) (2004).
- [43] G.F.J. Ananos and C. Tsallis, *Ensemble averages and nonextensivity at the edge of chaos of one-dimensional maps*, Phys. Rev. Lett. **93**, 020601 (2004).
- [44] E. Mayoral and A. Robledo, *Tsallis' q index and Mori's q phase transitions at edge of chaos*, Phys. Rev. E **72**, 026209 (2005).
- [45] U. Tirnakli, C. Beck and C. Tsallis, *Central limit behavior of deterministic dynamical systems*, Phys. Rev. E **75**, 040106 (2007).
- [46] G. Ruiz, T. Bountis and C. Tsallis, *Time-evolving statistics of chaotic orbits of conservative maps in the context of the Central Limit Theorem*, Int. J. Bifurcation and Chaos **22**, 1250208 (2012).
- [47] U. Tirnakli and E.P. Borges, *Standard map: From Boltzmann-Gibbs statistics to Tsallis statistics*, 1501.02459 [cond-mat.stat-mech] (2015).
- [48] C. Tsallis and D.J. Bukman, *Anomalous diffusion in the presence of external forces: exact time-dependent solutions and their thermostistical basis*, Phys. Rev. E **54**, R2197 (1996).
- [49] G. Drazer, H.S. Wio and C. Tsallis, *Anomalous diffusion with absorption: Exact time-dependent solutions*, Phys. Rev. E **61**, 1417 (2000).
- [50] F.D. Nobre and C. Tsallis, *Classical infinite-range-interaction Heisenberg ferromagnetic model: Metastability and sensitivity to initial conditions*, Phys. Rev. E **68**, 036115 (2003).
- [51] F.D. Nobre and C. Tsallis, *Metastable states of the classical inertial infinite-range-interaction Heisenberg ferromagnet: Role of initial conditions*, Physica A **344**, 587 (2004).
- [52] A.O. Caride, C. Tsallis and S.I. Zanette, *Criticality of the anisotropic quantum Heisenberg-model on a self-dual hierarchical lattice*, Phys. Rev. Lett. **51**, 145 (1983).
- [53] R.M. Pickup, R. Cywinski, C. Pappas, B. Farago and P. Fouquet, *Generalized spin glass relaxation*, Phys. Rev. Lett. **102**, 097202 (2009).
- [54] P. Douglas, S. Bergamini and F. Renzoni, *Tunable Tsallis distributions in dissipative optical lattices*, Rev. Lett. **96**, 110601 (2006).
- [55] E. Lutz and F. Renzoni, *Beyond Boltzmann-Gibbs statistical mechanics in optical lattices*, Nature Physics **9**, 615 (2013).
- [56] R.G. DeVoe, *Power-law distributions for a trapped ion interacting with a classical buffer gas*, Phys. Rev. Lett. **102**, 063001 (2009).
- [57] J. S. Andrade Jr., G.F.T. da Silva, A.A. Moreira, F.D. Nobre and E.M.F. Curado, *Thermostatistics of overdamped motion of interacting particles*, Phys. Rev. Lett. **105**, 260601 (2010).
- [58] M.S. Ribeiro, F.D. Nobre and E.M.F. Curado, *Time evolution of interacting vortices under overdamped motion*, Phys. Rev. E **85**, 021146 (2012).

- [59] M.S. Ribeiro, F.D. Nobre and E.M.F. Curado, *Overdamped motion of interacting particles in general confining potentials: Time-dependent and stationary-state analyses*, Eur. Phys. J. B **85**, 399 (2012).
- [60] E.M.F. Curado, A.M.C. Souza, F.D. Nobre and R.F.S. Andrade, *Carnot cycle for interacting particles in the absence of thermal noise*, Phys. Rev. E **89**, 022117 (2014).
- [61] R.F.S. Andrade, A.M.C. Souza, E.M.F. Curado and F.D. Nobre, *A thermodynamical formalism describing mechanical interactions*, EPL **108**, 20001 (2014).
- [62] B. Liu and J. Goree, *Superdiffusion and non-Gaussian statistics in a driven-dissipative 2D dusty plasma*, Phys. Rev. Lett. **100**, 055003 (2008).
- [63] C. Tsallis, *Dynamical scenario for nonextensive statistical mechanics*, Physica A **340**, 1 (2004).
- [64] L.F. Burlaga, A.F. Vinas, N.F. Ness and M.H. Acuna, *Tsallis statistics of the magnetic field in the heliosheath*, Astrophys. J. **644**, L83 (2006).
- [65] L.F. Burlaga and N.F. Ness, *Compressible “turbulence” observed in the heliosheath by Voyager 2*, Astrophys. J. **703**, 311 (2009); also, Astrophys. J. **725**, 1306 (2010), and Astrophys. J. **737**, 35 (2011).
- [66] L.F. Burlaga, N.F. Ness and M.H. Acuna, *Magnetic field strength fluctuations and temperature in the heliosheath*, Astrophys. J. **691**, L82 (2009).
- [67] J. Cho and A. Lazarian, *Simulations of electron magnetohydrodynamic turbulence*, Astrophys. J. **701**, 236 (2009).
- [68] A. Esquivel and A. Lazarian, *Tsallis statistics as a tool for studying interstellar turbulence*, Astrophys. J. **710**, 125 (2010).
- [69] G. Livadiotis and D.J. McComas, *Beyond kappa distributions: Exploiting Tsallis Statistical Mechanics in space plasmas*, J. Geophys. Res. - Space Physics **114**, A11105 (2009).
- [70] G. Livadiotis and D.J. McComas, *Exploring transitions of space plasmas out of equilibrium*, Astrophys. J. **714** (1), 971-987 (2010).
- [71] G. Livadiotis and D.J. McComas, *Measure of the departure of the q-metastable stationary states from equilibrium*, Phys. Scripta **82**, 035003 (2010).
- [72] D.J.B. Soares, C. Tsallis, A.M. Mariz and L.R. da Silva, *Preferential attachment growth model and nonextensive statistical mechanics*, EPL **70**, 70 (2005).
- [73] N. Komatsu and S. Kimura, *Entropic cosmology for a generalized black-hole entropy*, Phys. Rev. D **88**, 083534 (2013).
- [74] N. Komatsu, *Entropic cosmology from a thermodynamics viewpoint*, JPS Conf. Proc. **1**, 013112 (2014).
- [75] N. Komatsu and S. Kimura, *Evolution of the universe in entropic cosmologies via different formulations*, Phys. Rev. D **89**, 123501 (2014).
- [76] F.D. Nobre, M.A. Rego-Monteiro and C. Tsallis, *Nonlinear relativistic and quantum equations with a common type of solution*, Phys. Rev. Lett. **106**, 140601 (2011).
- [77] S. Umarov, C. Tsallis and S. Steinberg, *On a q-central limit theorem consistent with nonextensive statistical mechanics*, Milan J. Math. **76**, 307 (2008).

- [78] S. Umarov, C. Tsallis, M. Gell-Mann and S. Steinberg, *Generalization of symmetric α -stable Lévy distributions for $q > 1$* , J. Math. Phys. **51**, 033502 (2010).
- [79] C. Tsallis, M. Gell-Mann and Y. Sato, *Asymptotically scale-invariant occupancy of phase space makes the entropy S_q extensive*, Proc. Natl. Acad. Sc. USA **102**, 15377 (2005).
- [80] L.G. Moyano, C. Tsallis and M. Gell-Mann, *Numerical indications of a q -generalised central limit theorem*, EPL **73**, 813 (2006).
- [81] CMS Collaboration, *Transverse-momentum and pseudorapidity distributions of charged hadrons in pp collisions at $\sqrt{s} = 7$ TeV*, Phys. Rev. Lett. **105**, 022002 (2010); also, J. High Energy Phys. **09**, 091 (2010) and J. High Energy Phys. **05**, 064 (2011).
- [82] CMS Collaboration, *Observation of long-range, near-side angular correlations in proton-proton collisions at the LHC*, J. High Energy Phys. **09**, 091 (2010).
- [83] CMS Collaboration, *Strange particle production in pp collisions at $\sqrt{s} = 0.9$ and 7 TeV*, J. High Energy Phys. **05**, 064 (2011).
- [84] ALICE Collaboration, *Transverse momentum spectra of charged particles in proton-proton collisions $\sqrt{s} = 900$ GeV with ALICE at the LHC*, Phys. Lett. B **693**, 53 (2010); also, Eur. Phys. J. C **71**, 1594 (2011) and Eur. Phys. J. C **71**, 1655 (2011).
- [85] ALICE Collaboration, *Strange particle production in protonproton collisions at $\sqrt{s} = 0.9$ TeV with ALICE at the LHC*, Eur. Phys. J. C **71**, 1594 (2011).
- [86] ALICE Collaboration, *Production of pions, kaons and protons in pp collisions at $\sqrt{s} = 900$ GeV with ALICE at the LHC*, Eur. Phys. J. C **71**, 1655 (2011).
- [87] A. Tawfik, *Antiproton-to-proton ratios for ALICE heavy-ion collisions*, Nuclear Phys. A **859**, 63 (2011).
- [88] ATLAS Collaboration, *Charged-particle multiplicities in pp interactions measured with the ATLAS detector at the LHC*, New J. Physics **13**, 053033 (2011).
- [89] PHENIX Collaboration, *Measurement of neutral mesons in $p + p$ collisions at $\sqrt{s} = 200$ GeV and scaling properties of hadron production*, Phys. Rev. D **83**, 052004 (2011); also, Phys. Rev. C **83**, 024909 (2011), and Phys. Rev. C **83**, 064903 (2011).
- [90] M. Shao, L. Yi, Z.B. Tang, H.F. Chen, C. Li and Z.B. Xu, *Examination of the species and beam energy dependence of particle spectra using Tsallis statistics*, J. Phys. G **37**, 085104 (2010).
- [91] F.S. Navarra, O.V. Utyuzh, G. Wilk and Z. Wlodarczyk, *Estimating inelasticity with the information theory approach*, Phys. Rev. D **67**, 114002 (2003).
- [92] T. Osada, O.V. Utyuzh, G. Wilk and Z. Wlodarczyk, *Extended Gaussian ensemble or q -statistics in hadronic production processes?*, Eur. Phys. J. B **50**, 7 (2006).
- [93] M. Biyajima, T. Mizoguchi, N. Nakajima, N. Suzuki and G. Wilk, *Modified Hagedorn formula including temperature fluctuation - Estimation of temperature at RHIC experiments*, Eur. Phys. J. C **48**, 597 (2007).
- [94] T. Osada and G. Wilk, *Nonextensive hydrodynamics for relativistic heavy-ion collision*, Phys. Rev. C **77**, 044903 (2008).

- [95] T.S. Biro and E. Molnar, *Non-extensive statistics, relativistic kinetic theory and fluid dynamics*, Eur. Phys. J. A **48**, 172 (2012)
- [96] K. Urmosy, T.S. Biro and G.G. Barnafoldi, *Pion production via resonance decay in a non-extensive quark-gluon medium with non-additive energy composition rule*, EPJ **13**, 5003 (2011)
- [97] J. Cleymans, G. Hamar, P. Levai and S. Wheaton, *Near-thermal equilibrium with Tsallis distributions in heavy ion collisions*, J. Phys. G **36**, 064018 (2009).
- [98] J. Cleymans, *Recent developments around chemical equilibrium*, J. Phys. G: Nucl. Part. Phys. **37**, 094015 (2010)
- [99] L. Marques, E. Andrade-II and A. Deppman, *Nonextensivity of hadronic systems*, Phys. Rev. D **87**, 114022 (2013).
- [100] A. Deppman, *Self-consistency in non-extensive thermodynamics of highly excited hadronic states*, Physica A **391**, 6380 (2012); Corrigendum: Physica A **400**, 207 (2014).
- [101] A. Deppman, *Properties of hadronic systems according to the nonextensive self-consistent thermodynamics*, J. Phys. G: Nucl. Part. Phys. **41**, 055108 (2014) .
- [102] E. Megias, D.P. Menezes and A. Deppman, *Nonextensive thermodynamics for hadrons with finite chemical potentials*, Physica A **421**, 15 (2015).
- [103] C.Y. Wong and G. Wilk, *Tsallis fits to p_T spectra and relativistic hard scattering in pp collisions at LHC*, Phys. Rev. D **87**, 114007 (2013).
- [104] L.J.L. Cirto, C. Tsallis, C.Y. Wong and G. Wilk, *The transverse-momenta distributions in high-energy pp collisions – A statistical-mechanical approach*, arxiv 1409.3278 [hep-ph] (2014).
- [105] C. Tsallis, L.R. da Silva, R. S. Mendes, R.O. Vallejos and A.M. Mariz, *Specific heat anomalies associated with Cantor-set energy spectra*, Phys. Rev. E **56**, R4922 (1997).
- [106] C. Tsallis, *Nonadditive entropy and nonextensive statistical mechanics - An overview after 20 years*, Brazilian J. Physics **39**, 337 (2009).