

# On the Classification of the Lévy-Leblond Spinors

Luiza Miranda<sup>(1)</sup>, Isaquie P. de Freitas<sup>(2)</sup> and Francesco Toppan<sup>(3)</sup>

<sup>(1,2,3)</sup>COTEO, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro (RJ), Brazil.

E-mail: <sup>(1)</sup>luiza16@cbpf.br, <sup>(2)</sup>isaquiefreitas@cbpf.br, <sup>(3)</sup>toppan@cbpf.br

**Abstract.** The first-order Lévy-Leblond differential equations (LLEs) are the non-relativistic analogous of the Dirac equation: they are the “square roots” of the Schrödinger equation in  $(1 + d)$  dimensions and admit spinor solutions. In this paper we show how to extend to the Lévy-Leblond spinors the real/complex/quaternionic classification of the relativistic spinors (which leads to the notions of Dirac, Weyl, Majorana, Majorana-Weyl, Quaternionic spinors). Besides the free equations, we also consider the presence of potential terms. Applied to a conformal potential, the simplest  $(1 + 1)$ -dimensional LLE induces a new differential realization of the  $osp(1|2)$  superalgebra in terms of first-order differential operators depending on the time and space coordinates.

## 1 Introduction

The Dirac equation which introduces the notion of relativistic spinors can be regarded, see [1], as a “square root equation” of the relativistic Klein-Gordon equation. In a similar spirit, Lévy-Leblond introduced in 1967 a  $1+3$ -dimensional equation [2] which is the square root of the nonrelativistic (invariant under the Galilei group) Schrödinger equation. The equation implies the presence of nonrelativistic Lévy-Leblond spinors. Quite often the Lévy-Leblond equation (“LLE” for short) appears when taking, for the speed of light  $c$  of a relativistic system, the  $c \rightarrow \infty$  limit. On the other hand, LLE has its own virtue; it can be applied to investigate nonrelativistic systems, as those studied in condensed matter, which do not necessarily arise as a limit of relativistic systems.

As a clarification note, Lévy-Leblond correctly criticized the use of the term “nonrelativistic” to refer to theories satisfying the Galilei’s relativity principle described in *Dialogo sopra i due massimi sistemi del mondo*; despite of that, in this paper we adhere to the standard terminology, the adjective “relativistic” being exclusively applied to Einstein’s special relativity.

The theory of relativistic spinors is rather rich. Relativistic equations can be generalized to arbitrary  $(s, t)$  spacetimes, not only the  $1+3$  Minkowski spacetime. Depending on the spacetime, different solutions can be found. Besides complex spinors (as the original Dirac spinors), real Majorana spinors can be found, Weyl chiral/antichiral spinors can exist which, in certain spacetimes, can be combined to give Majorana-Weyl spinors. Spinors defined for the division algebra of quaternions produce different types of relativistic equations. The classification of the relativistic spinors is based on the [3] classification of the associated Clifford algebras. Relevant works (for this current paper) which present the classification of relativistic spinors are [4–6].

Generalized Lévy-Leblond equations can be introduced as square roots of Schrödinger equations in  $1 + d$ -dimensions, where the number  $d$  of the space components is an arbitrary integer. The Schrödinger equations under consideration can be free, but can also admit potential terms. The theories of LLEs and their associated spinors have not been systematically investigated as their relativistic counterparts (see, e.g., [7]).



Some words should be added to justify the interest in classifying the theories of nonrelativistic spinors. Obviously, information about relativistic theories should be recovered in the  $c \rightarrow \infty$  nonrelativistic theory and the mapping between the relativistic spinors and their nonrelativistic counterparts is an essential tool. Similarly, in recent years, the opposite  $c \rightarrow 0$  (so-called “Carroll”) limit of relativistic theories is widely investigated (the connection between Galilei and Carroll limits is discussed in [8]). The Galilei algebra is not semisimple and admits the Bargmann’s central extension [9]; due to that its representation theory is quite nontrivial. It also implies, as shown below in Section 5, a different role for the time and for the space coordinates. Last but not least, in applications to condensed matter emergent nonrelativistic spinors can arise in theories which do not necessarily result from the  $c \rightarrow \infty$  limit of relativistic models.

In this paper we show how to extend, to the generalized LLEs, the methods presented in [5] and [6] for relativistic spinors. It must be said that LLEs turned out to have surprisingly rich structures. The dynamical symmetry of the free equations possess [10, 11]  $Z_2 \times Z_2$ -graded Lie superalgebra structures, fitting the class of superalgebras introduced by Rittenberg-Wyler in [12, 13]. Furthermore it was recently shown [14] that, in the presence of a harmonic potential, the  $Z_2 \times Z_2$ -graded Lie superalgebras act as spectrum-generating superalgebras.

In this paper we show how to systematically classify the  $1 + d$ -dimensional Lévy-Leblond spinors as real (that is, of Majorana-type), complex (Dirac-type), quaternionic, chiral/antichiral (Weyl-type) and in determining their vector dimensions. We proceed at first with the free equations; later we show how to introduce potentials. The specialization to the inverse square potentials of the Conformal mechanics [15] implies new differential realizations of the superconformal algebras ( $osp(1|2)$  in the simplest case) in terms of differential operators in the time and space coordinates.

As a technical tool we introduce the convenient [16] “alphabetic presentation” of Clifford algebras which allows to represent the four  $2 \times 2$  building-block matrices of [6] as letters and the generic Clifford algebras gamma-matrices as words in a 4-letter alphabet.

## 2 Fundamental ingredients

The Clifford algebra  $Cl(p, q)$  is the enveloping algebra produced by the  $n \times n$  matrices  $\gamma_i$  which satisfy, for  $i, j = 1, 2, \dots, p + q$ , the anticommutation relations

$$\{\gamma_i, \gamma_j\} = 2\eta_{ij}\mathbb{I}_{n \times n}, \quad (1)$$

where  $\mathbb{I}_{n \times n}$  is the Identity matrix and  $\eta_{ij}$  is a  $(p + q) \times (p + q)$  diagonal matrix with  $p$  entries  $+1$  and  $q$  entries  $-1$ . In application to relativistic theories  $p$  is the number of space-like coordinates and  $q$  is the number of time-like coordinates. The irreducible representations of the  $Cl(p, q)$  algebras are recovered [5, 6] from tensoring four  $2 \times 2$  real matrices. They can be identified with four letters according to

$$X = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad Y = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}. \quad (2)$$

By dropping the symbol “ $\otimes$ ” in the tensor products expressing the generic  $\gamma_i$  matrices entering (1), we can associate these gamma matrices with words written in a 4-letter alphabet, see [25] for details. The three single-letter words  $X, Y, A$  define the three gamma matrices entering the  $Cl(2, 1)$  Clifford algebra. In the [6] recursive construction the five  $4 \times 4$  gamma matrices defining  $Cl(3, 2)$  can be expressed as the two-letter words  $XX, XY, XA, YI, AI$  (the three matrices  $XX, XY, YI$  are space-like, while the two matrices  $XA, AI$  are time-like). The general construction is presented in [6].

The length of the words defines the size of the corresponding gamma-matrices. The possibility of having Weyl-type spinors is ensured if all gamma matrices are of block-antidiagonal form. In the alphabetic presentation this requirement implies that all corresponding words should start with an  $Y$  or  $A$  letter.

The  $\mathbb{R}/\mathbb{C}/\mathbb{H}$  real/complex/quaternionic structures of the Clifford algebras and of their associated relativistic spinors are obtained, via Schur’s lemma, by looking at the most general matrices which, at given  $p, q$ , commute with all gamma matrices:

- in the  $\mathbb{R}$  case the most general matrix is proportional to the identity  $\mathbb{I}_{n \times n}$ ,
- in the  $\mathbb{C}$  case it is given, for  $a, b$  reals, by  $a\mathbb{I}_{n \times n} + bJ$ , where  $J^2 = -\mathbb{I}_{n \times n}$ ,
- in the  $\mathbb{H}$  case it is given, for  $a, b_1, b_2, b_3$  reals, by  $a\mathbb{I}_{n \times n} + \sum_i b_i J_i$ , where the three matrices  $J_i$  are imaginary quaternions satisfying the equations  $J_i J_j = -\delta_{ij}\mathbb{I}_{n \times n} + \varepsilon_{ijk} J_k$ .

The three above cases find an adequate description in terms of the alphabetic presentation.

The extension of the classification to the Lévy-Leblond nonrelativistic spinors requires introducing, in

the alphabetic presentation, a “fifth letter” denoted by  $Q$ . It correspond to the  $2 \times 2$  differential matrix

$$Q = \begin{pmatrix} 0 & 1 \\ i\partial_t & 0 \end{pmatrix} \Rightarrow Q^2 = i\partial_t \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}. \quad (3)$$

Essentially,  $Q$  is the square root of the time derivative entering the nonrelativistic Schrödinger equation.

An important property in the construction of LLEs is the anticommutation of the matrices associated with the letters  $Q$  and  $X$ :

$$\{Q, X\} = 0. \quad (4)$$

### 3 Lévy-Leblond free equations and types of spinors

By setting  $\hbar = 1$  and  $m = \frac{1}{2}$  the free (matrix) Schrödinger equation in  $1 + d$  dimensions reads as

$$i\hbar\partial_t\Psi(t, \vec{x}) = -\frac{\vec{\nabla}^2}{2m}\Psi(t, \vec{x}) \Rightarrow i\partial_t\Psi(t, \vec{x}) = -\vec{\nabla}^2\Psi(t, \vec{x}), \quad (5)$$

where  $\Psi(t, \vec{x})$  is an  $n$ -component vector (for  $n = 1, 2, 3, 4, \dots$ ). Its simplest square root equation is obtained for  $d = 1$  and  $n = 2$ . By using the (2,3) notation it is written as

$$Q\Psi(t, x) = X\partial_x\Psi(t, x) \Rightarrow i\partial_t\Psi = Q^2\Psi = QX\partial_x\Psi = -X\partial_xQ\Psi = -X^2\partial_x^2\Psi = -\partial_x^2\Psi. \quad (6)$$

This construction admits generalizations. The next simplest case is the square root of the  $n = 4$ -components  $d = 2$  free Schrödinger equation with space coordinates  $x, y$ . By taking an alphabetic basis of the five gamma matrices of the Clifford algebra  $Cl(3, 2)$ , given by  $XX, XY, XA, YI, AI$ , one can express the square root equation as

$$QI\Psi = (XX\partial_x + XY\partial_y)\Psi. \quad (7)$$

The corresponding 4-component spinors are real (that is, they are the nonrelativistic analogues of the Majorana spinors).

The 4-component equation

$$QY\Psi(t, x) = XY\partial_x\Psi(t, x) \quad (8)$$

admits spinors which are both real and (due to the presence of the block-antidiagonal matrix  $Y$ ) chiral; they are the nonrelativistic analogues of the Majorana-Weyl spinors. Due to the chirality constraint its fundamental components, in real counting, are  $\frac{1}{2} \times 4 = 2$ .

The next level of square root equations is obtained for  $n = 8$ -component spinors. The building blocks are the gamma matrices of the  $Cl(4, 3)$  Clifford algebras. Two equivalent alphabetic presentations are given by the following two sets of seven 3-letter words.

The first set is given by  $XXX, XXY, XXA, XYI, XAI, YII, AII$ .

The second set by  $XYX, XYY, XYA, XXI, XAI, YII, AII$ .

Three inequivalent square root equations are obtained.

The first one is

$$QII\Psi = (XXX\partial_x + XXY\partial_y + XYI\partial_z)\Psi, \quad (9)$$

producing 8-component Majorana-type spinors for a  $1 + 3$ -dimensional free LLE (the space coordinates being  $x, y, z$ ).

The second equation is

$$QII\Psi = XYI\partial_x\Psi. \quad (10)$$

The  $IIA$  complex structure (which commutes with both  $QYI$  and  $XYI$ ) implies complex (nonrelativistic analogues of Dirac) spinors for an 8-component  $(1 + 1)$ -dimensional free LLE.

The third equation is

$$QYI\Psi = (XYX\partial_x + XYY\partial_y)\Psi. \quad (11)$$

The presence of the  $Y$  letter in the second position implies that it describes nonrelativistic Majorana-Weyl type spinors which, due to chirality, possess  $\frac{1}{2} \times 8 = 4$  components. The equation (11) is an 8-component  $(1+2)$ -dimensional free LLE.

The next level is obtained for  $n = 16$ . At this level one recovers 5 inequivalent free Lévy-Leblond equations. They are:

- the 16-component “Majorana-type” spinor for the  $(1+4)$ -dimensional free LLE, given by

$$QIII\Psi = (XXXX\partial_x + XXXY\partial_y + XXYI\partial_z + XYII\partial_w)\Psi; \quad (12)$$

- the “Dirac-type” complex spinor for the  $(1+2)$ -dimensional free LLE with  $IIIA$  complex structure,

$$QIII\Psi = (XXYI\partial_x + XYII\partial_y)\Psi; \quad (13)$$

- the “Majorana-Weyl-type” spinor (with 8 real components) for the  $(1+3)$ -dimensional free LLE

$$QYII\Psi = (XYXX\partial_x + XYXY\partial_y + XYYI\partial_z)\Psi; \quad (14)$$

- the “Weyl-type” complex spinor for the  $(1+1)$ -dimensional free LLE with  $IIIA$  complex structure,

$$QYII\Psi = XYYI\partial_x\Psi; \quad (15)$$

- the quaternionic spinor for the  $(1+1)$ -dimensional free LLE with  $IIIA, IIAX, IIAY$  quaternionic structure,

$$QIII\Psi = XYII\partial_x\Psi. \quad (16)$$

We presented here the inequivalent free LLEs for  $n = 2, 4, 8, 16$ , that is  $n = 2^k$  for  $k = 1, 2, 3, 4$ . The construction can be easily extended to any  $k$ . It is based on the properties of the associated  $Cl(p, q)$  Clifford algebras. Up to  $k \leq 4$  we can present the following table illustrating the Majorana-type ( $M$ ), Weyl-type ( $W$ ), Majorana-Weyl-type ( $MW$ ), Dirac-type ( $D$ ), Quaternionic ( $H$ ) nonrelativistic spinors of the free  $(1+d)$ -dimensional LLEs. We get

$(2 \times 2)$ matrices:	$M$ ,	$(1+1)$ ,	2 real components,
$(4 \times 4)$ matrices:	$M$ ,	$(1+2)$ ,	4 real components,
$(4 \times 4)$ matrices:	$MW$ ,	$(1+1)$ ,	$4/2 = 2$ real components,
$(8 \times 8)$ matrices:	$M$ ,	$(1+3)$ ,	8 real components,
$(8 \times 8)$ matrices:	$MW$ ,	$(1+2)$ ,	$8/2 = 4$ real components,
$(8 \times 8)$ matrices:	$D$ ,	$(1+1)$ ,	$4_{\mathbf{C}} \equiv 8$ real components,
$(16 \times 16)$ matrices:	$M$ ,	$(1+4)$ ,	16 real components,
$(16 \times 16)$ matrices:	$MW$ ,	$(1+3)$ ,	$16/2 = 8$ real components,
$(16 \times 16)$ matrices:	$D$ ,	$(1+2)$ ,	$8_{\mathbf{C}} \equiv 16$ real components,
$(16 \times 16)$ matrices:	$W$ ,	$(1+1)$ ,	$4_{\mathbf{C}} \equiv 8$ real components,
$(16 \times 16)$ matrices:	$H$ ,	$(1+1)$ ,	$4_{\mathbf{H}} \equiv 16$ real components.

(17)

## 4 The introduction of potential terms

In this Section we show how our scheme can be enlarged to introduce Lévy-Leblond equations which are square roots of matrix Schrödinger equations in the presence of potential terms. We illustrate the simplest example of the nonrelativistic Majorana spinors in  $1+1$  dimensions.

The introduction of potential terms requires larger matrix realizations with respect to the free case. While the free Lévy-Leblond equation in  $1+1$  dimensions can be realized by  $2 \times 2$  matrices, we need at least  $4 \times 4$  matrices to include the potentials. The construction goes as follows. The basic equation is

$$QI\Psi = XY\partial_x\Psi + XAf(x)\Psi, \quad (18)$$

where the “words”  $QI$ ,  $XY$ , and  $XA$  represent  $4 \times 4$  matrices; the symbol  $\partial_x$  denotes the derivative with respect to the space coordinate  $x$  and the function  $f(x)$  is the prepotential which allows to reconstruct the potential terms of the matrix Schrödinger equation. The column vector spinor  $\Psi(x, t)$  has 4 components ( $\Psi^T = (\psi_1, \psi_2, \psi_3, \psi_4)$ ). The following equations are satisfied for its  $\psi_i(x, t)$  components with  $i = 1, 2, 3, 4$ :

$$\begin{aligned}\psi_3 &= \partial_x \psi_2 + f(x) \psi_2, \\ \psi_4 &= \partial_x \psi_1 - f(x) \psi_1,\end{aligned}\tag{19}$$

together with the time-dependent equations

$$\begin{aligned}i\partial_t \psi_1 &= -\partial_x \psi_4 - f(x) \psi_4, \\ i\partial_t \psi_2 &= -\partial_x \psi_3 + f(x) \psi_3.\end{aligned}\tag{20}$$

The (19) algebraic equations for  $\psi_3, \psi_4$  imply two independent Schrödinger equations for  $\psi_1, \psi_2$ , given by

$$\begin{aligned}i\partial_t \psi_1 &= -\partial_x^2 \psi_1 + (f^2(x) + f'(x)) \psi_1, \\ i\partial_t \psi_2 &= -\partial_x^2 \psi_2 + (f^2(x) - f'(x)) \psi_2,\end{aligned}\tag{21}$$

where  $f'(x) = \frac{d}{dx} f(x)$  and

$$V_{\pm}(x) = f^2(x) \pm f'(x)\tag{22}$$

are the respective potentials.

By applying the  $i\partial_t$  derivative to  $\psi_3, \psi_4$  entering (19) and taking into account the (20) equations, we obtain that the  $\psi_3, \psi_4$  components satisfy the Schrödinger equations

$$\begin{aligned}i\partial_t \psi_3 &= -\partial_x^2 \psi_3 + V_+(x) \psi_3, \\ i\partial_t \psi_4 &= -\partial_x^2 \psi_4 + V_-(x) \psi_4.\end{aligned}\tag{23}$$

The extension of the procedure to introduce potentials for nonrelativistic spinors in  $(1+d)$ -dimensions, complex structures, etc., is rather straightforward; it is based on the previously discussed construction of the free equations.

The specialization to certain types of potentials, like the inverse square potential proportional to  $\frac{1}{x^2}$ , produces interesting differential realizations of the superalgebras associated with the Lévy-Leblond square root of the matrix conformal mechanics. The simplest example is illustrated in the next Section.

## 5 The $osp(1|2)$ superalgebra induced by a conformal potential

In  $1+1$  dimensions, the Lévy-Leblond “square root” of a matrix conformal mechanics with potential term proportional to  $1/x^2$  is given by the  $4 \times 4$  matrix differential equation

$$\Omega \Psi(x, t) = 0,\tag{24}$$

where the  $\Psi(x, t)$  Lévy-Leblond spinor is a 4-component column vector and the matrix differential operator  $\Omega$  belongs to a 5-generator  $osp(1|2)$  superalgebra. The explicit form of the five  $osp(1|2)$  generators is given in terms of the  $2 \times 2$  matrices  $I, X, Y, A$  and  $Q$  introduced in (2) and (3), plus two extra  $2 \times 2$  auxiliary matrices  $\Lambda$  and  $R$  which depend on an (arbitrary) real scaling parameter  $\lambda$ :

$$\Lambda = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda + \frac{1}{2} \end{pmatrix}, \quad R = \begin{pmatrix} 0 & 0 \\ \lambda & 0 \end{pmatrix}.\tag{25}$$

The consistent scaling assignments of the time/space coordinates  $t, x$ , of their derivatives and of the auxiliary  $2 \times 2$  matrices  $Q, \Lambda, R$  are:

$$[t] = -1, \quad [\partial_t] = +1, \quad [x] = -\frac{1}{2}, \quad [\partial_x] = +\frac{1}{2}, \quad [Q] = +\frac{1}{2}, \quad [\Lambda] = 0, \quad [R] = -\frac{1}{2}.\tag{26}$$

The five  $osp(1|2)$  generators (denoted as  $H, \Omega, D, \Xi, K$ ) are given by

$$\begin{aligned} H &= I \otimes I \cdot (i\partial_t + \partial_x^2 - \frac{g^2}{x^2}) + I \otimes X \cdot \frac{g}{x^2}, \\ \Omega &= Q \otimes I - X \otimes Y \cdot \partial_x - X \otimes A \cdot \frac{g}{x}, \\ D &= I \otimes I \cdot (\frac{1}{4} + \frac{1}{2}x\partial_x + t\partial_t) + \Lambda \otimes I, \\ \Xi &= Q \otimes I \cdot (-it) - X \otimes Y \cdot \frac{x}{2} + R \otimes I, \\ K &= I \otimes I \cdot (-it^2\partial_t + \frac{x^2}{4}) - \Lambda \otimes I \cdot 2it, \end{aligned} \quad (27)$$

where  $g$  is a dimensionless ( $[g] = 0$ ) coupling constant.

The scaling dimensions of the above five generators are

$$[H] = +1, \quad [\Omega] = +\frac{1}{2}, \quad [D] = 0, \quad [\Xi] = -\frac{1}{2}, \quad [K] = -1. \quad (28)$$

The three generators  $H, D, K$  are even (bosonic) and close an  $sl(2)$  subalgebra having  $D$  as the Cartan generator. The two generators  $\Omega, \Xi$  are odd (fermionic); they respectively correspond to the positive and negative simple roots of  $osp(1|2)$ .

The closure of the  $osp(1|2)$  superalgebra is given by the following (anti)commutators:

$$\begin{aligned} [D, H] &= -H, & [D, K] &= K, & [H, K] &= 2D, \\ [D, \Omega] &= -\frac{1}{2}\Omega, & [D, \Xi] &= \frac{1}{2}\Xi, \\ [H, \Omega] &= 0, & [K, \Omega] &= -\Xi, & [K, \Xi] &= 0, \\ \{\Omega, \Omega\} &= 2H, & \{\Omega, \Xi\} &= 2D, & \{\Xi, \Xi\} &= 2K. \end{aligned} \quad (29)$$

Therefore,  $\Omega$  is the square root of the  $4 \times 4$  matrix conformal mechanics Schrödinger equation

$$H\Psi(x, t) = 0. \quad (30)$$

Usually, a Schrödinger equation such as (30) is split into a left part which depends on a first-order time-derivative and a right part which depends on a second-order differential equation in  $x$  (the Hamiltonian, which we denote in boldface as  $\mathbf{H}$ ); the (30) equation can therefore be rewritten as

$$I \otimes I \cdot i\partial_t \Psi(x, t) = \mathbf{H}\Psi(x, t) \equiv I \otimes I(-\partial_x^2 + \frac{g^2}{x^2}) - I \otimes X \cdot \frac{g}{x^2}. \quad (31)$$

Superconformal algebra realizations of first-order in time “ $D$ -module” differential realizations have been investigated in [17]. For the free Schrödinger equation, a second-order differential realization of  $osp(1|2)$  in terms of the space coordinate  $x$  appears naturally (see [18], which connects the [19] Wigner’s approach to the quantization of the harmonic oscillator with the Niederer’s analysis [20–22] of the maximal kinematical invariance group of the Schrödinger equation for various potentials).

The Lévy-Leblond equation  $\Omega\Psi = 0$  induces, for the conformal potential proportional to  $\frac{1}{x^2}$ , a new interesting and non-trivial differential realization of  $osp(1|2)$  which involves both time and space coordinates.

The construction of the admissible Hilbert spaces for the conformal mechanics, depending on the range of the coupling constant  $g$ , has been discussed in [23, 24] and, in the matrix case, [25].

An open question should be pointed out. The addition in conformal mechanics of a harmonic potential term to the  $\frac{1}{x^2}$  potential produces the de Alfaro-Fubini-Furlan deformed oscillator [26] having the same conformal algebra as spectrum-generating algebra. The new Hamiltonian is the sum of  $\mathbf{H}$  and of its conformal partner. Concerning the (29) realization, one should note that the sum  $H + K$  does not produce a Lévy-Leblond equation. It is an open question how to produce a de Alfaro-Fubini-Furlan construction for the LLE.

## 6 Conclusions

This paper outlines the classification of the Lévy-Leblond nonrelativistic spinors and of their associated equations using tools and methods borrowed from the classification of relativistic spinors. It further

illustrates in simple examples how to introduce potential terms to Lévy-Leblond equations and how to derive the “square root” of conformal mechanics with their induced superconformal algebras. This work is part of a research program where an extended version and systematic presentation of the constructions for real/complex/quaternionic and Weyl-type Lévy-Leblond spinors is under preparation.

## Acknowledgments

This work was supported by CNPq (PQ grant 308846/2021-4).

## References

- 1 P. A. M. Dirac, *The quantum theory of the electron*, Proc. R. Soc. **A 117 (778)**, 610 (1928).
- 2 J. M. Lévy-Leblond, *Nonrelativistic particles and wave equations*, Comm. Math. Phys. **6**, 286 (1967).
- 3 M. F. Atiyah, R. Bott and A. Shapiro, *Clifford modules*, Topology **3 Supp. 1**, 3 (1964).
- 4 T. Kugo and P. Twonsend, *Supersymmetry and the division algebras*, Nucl. Phys. **B 221**, 357 (1983).
- 5 S. Okubo, *Real representations of finite Clifford algebras. I. Classification*, J. Math. Phys. **32**, 1657 (1991).
- 6 H. L. Carrion, M. Rojas and F. Toppan, *Quaternionic and Octonionic Spinors. A Classification*, JHEP **0304**, 040 (2003); arXiv:hep-th/0302113.
- 7 A. Chakraborty, B. Debnath, R. Datta and P. Banerjee, *Construction of a Few Quantum Mechanical Hamiltonians via Lévy-Leblond Type Linearization: Clifford Momentum, Spinor States and Supersymmetry*, Adv. Appl. Clifford Algebras **32**, 56 (2022).
- 8 J. Figueroa O’Farrill, *Lie algebraic Carroll/Galilei duality*, J. Math. Phys. **64**, 013503 (2023); arXiv:2210.13924.
- 9 V. Bargmann, *On Unitary Ray Representations of Continuous Groups*, Ann. of Math. **59**, 1 (1954).
- 10 N. Aizawa, Z. Kuznetsova, H. Tanaka and F. Toppan,  *$\mathbb{Z}_2^2$ -graded Lie symmetries of the Lévy-Leblond equations*, Prog. Theor. Exp. Phys. **2016**, 123A01 (2016); arXiv:1609.08224[math-ph].
- 11 N. Aizawa, Z. Kuznetsova, H. Tanaka and F. Toppan, *Generalized supersymmetry and Lévy-Leblond equation*, in S. Duarte et al (eds), Physical and Mathematical Aspects of Symmetries, Springer, Cham, p. 79 (2017); arXiv:1609.08760[math-ph].
- 12 V. Rittenberg and D. Wyler, *Generalized Superalgebras*, Nucl. Phys. **B 139**, 189 (1978).
- 13 V. Rittenberg and D. Wyler, *Sequences of  $\mathbb{Z}_2 \otimes \mathbb{Z}_2$  graded Lie algebras and superalgebras*, J. Math. Phys. **19**, 2193(1978).
- 14 M. Ryan, *Graded colour Lie superalgebras for solving Lévy-Leblond equations*, arXiv:2407.19723[math-ph].
- 15 F. Calogero, *Solution of a three-body problem in one dimension*, J. Math. Phys. **10**, 2191 (1969).
- 16 F. Toppan and P. W. Verbeek, *On Alphabetic Presentations of Clifford Algebras and Their Possible Applications*, J. Math. Phys. **50**, 123523 (2009); arXiv:0903.0940 [math-ph].
- 17 Z. Kuznetsova and F. Toppan, *D-module Representations of  $N = 2, 4, 8$  Superconformal Algebras and Their Superconformal Mechanics*, J. Math. Phys. **53**, 043513 (2012); arXiv:1112.0995[hep-th].
- 18 F. Toppan, *Symmetries of the Schrödinger Equation and Algebra/Superalgebra Duality*, J. of Phys: Conf. Series **597**, 012071 (2015); arXiv:1411.7867[math-ph].
- 19 E. P. Wigner, *Do the Equations of Motion Determine the Quantum Mechanical Commutation Relations?*, Phys. Rev. **77**, 711 (1950).
- 20 U. Niederer, *The Maximal Kinematical Invariance Group of the Free Schrödinger Equation*, Helv. Phys. Acta **45**, 802 (1972).

- 21 U. Niederer, *The Maximal Kinematic Invariance Group of the Harmonic Oscillator*, Helv. Phys. Acta **46**, 191 (1973).
- 22 U. Niederer, *The maximal kinematic invariance groups of the Schrödinger equations with arbitrary potentials*, Helv. Phys. Acta **47**, 167 (1974).
- 23 H. Miyazaki and I. Tsutsui, *Quantum Tunneling and Caustics under Inverse Square Potential*, Ann. Phys. **299**, 78 (2002); arXiv:quant-ph/0202037.
- 24 L. Fehér, I. Tsutsui and T. Fülöp, *Inequivalent quantizations of the three-particle Calogero model constructed by separation of variables*, Nucl. Phys. **B 715**, 713 (2005); arXiv:math-ph/0412095.
- 25 F. Toppan and M. Valenzuela, *Higher Spin Symmetries and Deformed Schrödinger Algebra in Conformal Mechanics*, Adv. Math. Phys. **6263150** (2018); arXiv:1705.04004[hep-th].
- 26 V. de Alfaro, S. Fubini and G. Furlan, *Conformal invariance in quantum mechanics*, Nuovo Cim. **A 34**, 569 (1976).
- 27 N. L. Holanda and F. Toppan, *Four types of (super)conformal mechanics: D-module reps and invariant actions*, J. Math. Phys. **55**, 061703 (2014); arXiv:1402.7298[hep-th].