

NOTAS DE FÍSICA

VOLUME VI

Nº 10

OBSERVATION OF A SHORT-LIVED COSMIC-RAY SOLAR FLARE INCREASE
WITH A HIGH-COUNTING-RATE MESON DETECTOR

by

K. G. McCracken and R. A. R. Palmeira

CENTRO BRASILEIRO DE PESQUISAS FÍSICAS

Av. Wenceslau Braz, 71

RIO DE JANEIRO

1960

OBSERVATION OF A SHORT-LIVED COSMIC-RAY SOLAR FLARE INCREASE
WITH A HIGH-COUNTING-RATE MESON DETECTOR

K. G. McCracken

Physics Department and Laboratory for Nuclear Science,
Massachusetts Institute of Technology, Cambridge, Massachusetts

and

R. A. R. Palmeira*

Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil

(Published in Physical Review Letters, Vol. 5, Nº 1, 15, (1960))

(Received October 13, 1960)

We report here observations of a very abrupt and short-lived cosmic-ray increase which was observed at M.I.T. (geographic coordinates $41^{\circ}23'N$, $71^{\circ}08'W$, sea level) on May 4, 1960. These data were obtained from three large meson telescopes, of

* This work was performed when one of the authors (R.P.) was at the M.I.T.

total sensitive area 10 m^2 , and total counting rate ≈ 900 counts sec^{-1} . The telescopes have been described elsewhere,¹ and it suffices to state that there are now 7.5 cm of lead between the scintillators. The minimum μ -meson momentum accepted by the telescope is now approximately 202 Mev/c.

The data from the telescopes are recorded as follows.

(1) The counting rate from each telescope is accumulated over intervals of six minutes, and punched on tape in the form of binary numbers. (2) A printing register records the hourly totals from each telescope as decimal numbers. (3) The outputs from the three telescopes are combined, accumulated over intervals of 30 seconds, and the accumulated count registered on a chart recorder.

Figure 1 displays the 30-second counting rate data obtained between 1010 and 1110 U.T. on May 4, 1960. The total




Fig. 1. The 30-second meson data observed at M.I.T. subsequent to the production of cosmic rays by the sun on May 4, 1960.

number of counts recorded in each 30-second interval is plotted as a percentage relative to the mean counting rate for the period 1010 to 1030. The standard deviation which is appropriate to any 30-second measurement is shown. To check the reliability

of the data shown in the figure, we have examined the 6-minute and hourly counting rate data, finding that there is no

significant difference between the variations observed by the three telescopes.

Comparison with WWV revealed that our time standard was lagging 8 seconds behind Universal Time: for example, the count total plotted between 1015 and 1015 + 30 seconds, was obtained between 10:15:08 U.T. and 10:15:38 U.T. No corrections have been made for changes of barometric pressure, as it remained constant over this hour to within 0.005 inch of mercury.

It is seen from the figure that the counting rate measurements remained at the pre-event value until 10:31:08. The next measurement, that is, the one obtained between 10:31:08 and 10:31:38, was 4 standard deviations above this value, and we therefore estimate that the first arrival of solar cosmic rays of energies ≈ 6 Bev at M.I.T. occurred no later than 10:31:38 and, probably, not before about 10:30:53. The maximum counting rate (6% above the pre-event value) was attained about 10:35 U.T., and after remaining at this value for about 5 minutes, the counting rate rapidly returned to the pre-event value, our data showing no significant continuation of the solar flare effect after 10:55 U.T.

According to a preliminary report issued by the High Altitude Observatory at Boulder, Colorado,² a flare of importance 2 was observed at 10:20 U.T. in an active region on the western solar limb. As no other solar flares were reported to have occurred at this time, we conclude that the cosmic radiation was generated in this flare.

The most outstanding feature of the cosmic-ray effect reported in this communication is the very short time scale - the rising intensity and decaying intensity phases being much shorter than those observed during any previous case of this type event. A study of the temporal dependence of all recorded cases of the cosmic-ray flare effect has been completed recently, and a model of the magnetic configurations in the sun-earth region has been constructed. These results will be published elsewhere in the near future. It suffices to say here that, on the basis of the model applicable on May 4, the short time scale would be explained in terms of the earth being within an essentially radial field which was rooted in the active region on the sun. The very short time scale suggests that there would be relatively few irregularities in this field and, consequently, the radiation arriving at the earth would be largely collimated along the lines of the field. This predicts that during the May 4 event (1) impact zones were very pronounced, (2) there was little time dispersion between the high - and low - energy particles, and (3) the effect observed by stations outside the impact zones was relatively small. It is highly desirable that data with good time resolution from many parts of the world be compared to determine whether these predictions were fulfilled.

The meson telescopes, with their fine-detail timing circuits, were designed by Professor R. W. Williams and Mr. R. D'Arcy for the specific purpose of studying cosmic-ray flare increases in detail.

- 1 R. A. R. Palmeira and R. W. Williams, Nuovo Cimento 8, 352 (1958).
- 2 Preliminary Report of Solar Activity Number TR 453, High Altitude Observatory, Boulder, Colorado.
